

PB-229 203

SAVANNAH RIVER STUDIES, JULY 1951 -  
JULY 1952

D. D. Gold, et al

Robert A. Taft Sanitary Engineering Center  
Cincinnati, Ohio

April 1954

DISTRIBUTED BY:

**NTIS**

National Technical Information Service  
U. S. DEPARTMENT OF COMMERCE  
5285 Port Royal Road, Springfield Va. 22151

INTERIM REPORT  
ON THE  
SAVANNAH RIVER STUDIES  
July 1951 - July 1952

by

D. D. Gold  
D. C. Scott  
A. Hirsch  
C. C. Ruchhoft  
L. R. Setter

PROPERTY OF:  
U.S. DEPT. OF THE INTERIOR  
FWPCA - Federal Wildlife Service  
633 Indiana Avenue, N.W.

U. S. DEPARTMENT OF  
HEALTH, EDUCATION, AND WELFARE  
Public Health Service  
Bureau of State Services  
Division of Sanitary Engineering Services  
Robert A. Taft Sanitary Engineering Center  
Cincinnati, Ohio  
April 1954



TABLE OF CONTENTS

Reproduced from  
best available copy.

|                                                                                      | Page |
|--------------------------------------------------------------------------------------|------|
| LIST OF CONTRIBUTORS . . . . .                                                       | 1    |
| ACKNOWLEDGMENTS . . . . .                                                            | ii   |
| CHAPTER I                                                                            |      |
| <u>SUMMARY OF FINDINGS AND RECOMMENDATIONS</u> . . . . .                             | 1    |
| FINDINGS AND CONCLUSIONS . . . . .                                                   | 1    |
| RECOMMENDATIONS . . . . .                                                            | 2    |
| CHAPTER II                                                                           |      |
| <u>GENERAL INFORMATION</u> . . . . .                                                 | 10   |
| OBJECTIVES . . . . .                                                                 | 10   |
| INITIATION OF STUDIES . . . . .                                                      | 11   |
| SAMPLING PROGRAM . . . . .                                                           | 12   |
| LOCATION BY RIVER MILES OF SAMPLING STATIONS,<br>LANDINGS, TRIBUTARIES, ETC. . . . . | 13   |
| LABORATORY ACTIVITIES . . . . .                                                      | 18   |
| SPECIAL STUDIES . . . . .                                                            | 19   |
| CHAPTER III                                                                          |      |
| <u>GENERAL DESCRIPTION</u> . . . . .                                                 | 21   |
| THE SAVANNAH RIVER BASIN . . . . .                                                   | 21   |
| TOPOGRAPHY AND GEOLOGY (Clark Hill Dam to Atlantic Ocean) . .                        | 21   |
| POPULATION . . . . .                                                                 | 23   |
| LAND USE . . . . .                                                                   | 23   |
| WATER USE AND DEVELOPMENT . . . . .                                                  | 23   |
| <u>Fish and Wildlife</u> . . . . .                                                   | 23   |
| <u>Recreation</u> . . . . .                                                          | 24   |
| <u>Reservoirs and Diversion Projects</u> . . . . .                                   | 24   |
| Clark Hill . . . . .                                                                 | 24   |
| Proposed Dams and Reservoirs . . . . .                                               | 25   |
| Diversion of Savannah River Water . . . . .                                          | 26   |
| Navigation . . . . .                                                                 | 26   |
| <u>Water Supplies</u> . . . . .                                                      | 26   |
| <u>Sources of Pollution</u> . . . . .                                                | 27   |
| Domestic Sewage . . . . .                                                            | 27   |
| Industrial Wastes . . . . .                                                          | 29   |

# CHAPTER IV

|                                                         | Page |
|---------------------------------------------------------|------|
| <u>STREAM FLOWS</u> . . . . .                           | 33   |
| DISCHARGE DATA . . . . .                                | 33   |
| <u>Savannah River at Augusta, Georgia</u> . . . . .     | 33   |
| <u>Savannah River at Burtons Ferry Bridge</u> . . . . . | 33   |
| <u>Savannah River Near Clyo</u> . . . . .               | 34   |
| <u>Augusta Canal Near Augusta, Georgia</u> . . . . .    | 34   |
| <u>Briar Creek at Millhaven, Georgia</u> . . . . .      | 34   |

# CHAPTER V

|                                                                                                |    |
|------------------------------------------------------------------------------------------------|----|
| <u>CHEMICAL, BACTERIOLOGICAL AND PHYSICAL STUDIES</u> . . . . .                                | 37 |
| WATER QUALITY . . . . .                                                                        | 38 |
| <u>Dissolved Oxygen</u> . . . . .                                                              | 38 |
| <u>Biochemical Oxygen Demand</u> . . . . .                                                     | 38 |
| <u>Coliform MPN</u> . . . . .                                                                  | 39 |
| <u>Minerals</u> . . . . .                                                                      | 39 |
| <u>Turbidity</u> . . . . .                                                                     | 43 |
| <u>Color</u> . . . . .                                                                         | 43 |
| SPECIAL STUDIES . . . . .                                                                      | 44 |
| <u>Dissolved Oxygen and Salinity Studies in the</u><br><u>Savannah River Estuary</u> . . . . . | 44 |
| <u>Study of Sludge Bank in Lower Savannah River</u><br><u>Section, S-14 to S-8</u> . . . . .   | 46 |
| <u>Clark Hill Reservoir Studies</u> . . . . .                                                  | 58 |
| SAVANNAH RIVER WATER TEMPERATURE IN AUGUSTA CANAL AT<br>AUGUSTA, GEORGIA . . . . .             | 71 |

# CHAPTER VI

|                                                                        |     |
|------------------------------------------------------------------------|-----|
| <u>BIOLOGICAL STUDIES</u> . . . . .                                    | 74  |
| FRESHWATER ENVIRONMENTAL ASSOCIATIONS . . . . .                        | 74  |
| <u>Plankton</u> . . . . .                                              | 76  |
| <u>Benthos</u> . . . . .                                               | 83  |
| <u>Benthic Algal Growths and Associated Organisms</u> . . . . .        | 83  |
| Organisms Associated with Compact Clay. . . . .                        | 96  |
| Organisms Associated with Sand and Mud. . . . .                        | 100 |
| Organisms Associated with Brush, Trees,<br>Rocks, Gravel, etc. . . . . | 106 |
| <u>Nekton</u> . . . . .                                                | 116 |

|                                                | Page |
|------------------------------------------------|------|
| COMMERCIAL AND SPORT FISHING . . . . .         | 116  |
| <u>Commercial Fishing</u> . . . . .            | 116  |
| Basket Fishing for Catfish . . . . .           | 117  |
| Shad Fishing . . . . .                         | 118  |
| <u>Sport Fishing</u> . . . . .                 | 124  |
| SEINE AND NET SAMPLING . . . . .               | 125  |
| <u>Hoop Netting</u> . . . . .                  | 125  |
| <u>Gill Netting</u> . . . . .                  | 130  |
| <u>Seining</u> . . . . .                       | 130  |
| ROTENONE SAMPLING . . . . .                    | 131  |
| AGE AND GROWTH DETERMINATIONS . . . . .        | 134  |
| FISH MOVEMENT STUDIES . . . . .                | 137  |
| ANNOTATED LIST OF FRESH WATER FISHES . . . . . | 138  |
| ESTUARINE STUDIES . . . . .                    | 157  |

## CHAPTER VII

|                                                              |     |
|--------------------------------------------------------------|-----|
| <u>BACKGROUND RADIOACTIVITY OF WATER AND BIOTA</u> . . . . . | 160 |
| RADIOACTIVITY OF WATER . . . . .                             | 161 |
| RADIOACTIVITY OF FRESH WATER BIOTA . . . . .                 | 163 |
| RADIOACTIVITY OF ESTUARINE BIOTA . . . . .                   | 163 |
| MINIMUM DETECTABLE RADIOACTIVITY . . . . .                   | 165 |

## APPENDICES

### APPENDIX "A"

|                                                              |     |
|--------------------------------------------------------------|-----|
| <u>SUMMARY OF INDIVIDUAL RESULTS - SANITARY EXAMINATIONS</u> |     |
| S-230.7 - S-218.5 . . . . .                                  | 177 |
| S-218.5 - S-217.2;HGG 0.01 . . . . .                         | 178 |
| S-217.2;HGG 0.01 - S-213.36;Ho 0.01 . . . . .                | 179 |
| S-212.0 . . . . .                                            | 180 |
| S-203.3 - S-202.9 . . . . .                                  | 181 |
| S-202.65;Bu 0.02 - S-202.65;Bu 2.2 . . . . .                 | 182 |
| S-202.65;Bu 2.2 - S-201.65 . . . . .                         | 183 |
| S-201.65 - S-197.0 . . . . .                                 | 184 |
| S-197.0 - S-182.0 . . . . .                                  | 185 |
| S-177.0 - S-172.0 . . . . .                                  | 186 |
| S-172.0 - S-167.0 . . . . .                                  | 187 |
| S-164.26;Fo 0.01 - S-157.0 . . . . .                         | 188 |
| S-155.2;St 0.01 - S-152.0 . . . . .                          | 189 |
| S-152.0 - S-137.32;LTR 0.01 . . . . .                        | 190 |
| S-132.0 - S-105.0;Bl 0.2 . . . . .                           | 191 |

|                            | Page |
|----------------------------|------|
| S-102.0 - S-52.0 . . . . . | 192  |
| S-52.0 - S-32.0 . . . . .  | 193  |
| S-11.0 . . . . .           | 194  |

APPENDIX "B"

SUMMARY OF INDIVIDUAL RESULTS - MINERAL EXAMINATIONS

|                                      |     |
|--------------------------------------|-----|
| S-218.5 - S-164.26;Fo 0.01 . . . . . | 197 |
| S-155.2;St 0.01 - S-11 . . . . .     | 198 |

APPENDIX "C"

CLARK HILL RESERVOIR - SUMMARY OF INDIVIDUAL RESULTS -  
IRON, MANGANESE, TEMPERATURE,  
TURBIDITY AND DISSOLVED OXYGEN  
DETERMINATIONS; RESERVOIR DISCHARGE  
DATA

|                                                                                                                   |     |
|-------------------------------------------------------------------------------------------------------------------|-----|
| S-255<br>Ga.Hwy. #43 Bridge over Savannah River<br>4/30/52 - 5/22/52 . . . . .                                    | 201 |
| S-253, Li(S.C.)-3.2<br>S.C.Hwy. #43 Bridge over Little River, S.C.<br>4/30/52 - 5/22/52 . . . . .                 | 202 |
| S-246<br>8 miles above Dam - 5/6/52 - 5/21/52 . . . . .                                                           | 203 |
| S-239.1, Li(Ga)-9<br>Ga.Hwy. #47 Bridge over Little River, Ga.<br>4/30/52 - 5/22/52 . . . . .                     | 204 |
| S-239.1<br>1 mile above Dam - Junction of Little River, Ga.<br>with Savannah River<br>3/10/52 - 5/1/52 . . . . .  | 205 |
| S-239.1<br>1 mile above Dam - Junction of Little River, Ga.<br>with Savannah River<br>5/3/52 . . . . .            | 206 |
| S-239.1<br>1 mile above Dam - Junction of Little River, Ga.<br>with Savannah River<br>5/21/52 - 5/26/52 . . . . . | 207 |

|                                                                         | Page |
|-------------------------------------------------------------------------|------|
| S-239.1                                                                 |      |
| 1 mile above Dam - Junction of Little River, Ga.<br>with Savannah River |      |
| 6/13/52 . . . . .                                                       | 208  |
| PROVISIONAL DATA                                                        |      |
| S-237.8                                                                 |      |
| 0.3 mile below Clark Hill Dam.                                          |      |
| Sample taken from construction bridge below the dam . .                 | 209  |
| 5/26/52                                                                 |      |

#### APPENDIX "D"

##### SUMMARY OF INDIVIDUAL ANALYTICAL RESULTS FOR THE SAVANNAH RIVER SECTION - S-23 to S-0

|                                        |     |
|----------------------------------------|-----|
| S-23 to S-2 - March 28, 1952 . . . . . | 212 |
| S-23 to S-0 - June 23, 1952 . . . . .  | 213 |

#### APPENDIX "E"

##### DESCRIPTION OF THE PROCEDURE USED FOR ASSAYING THE RADIOACTIVITY IN FRESH WATER AND FOR BIOTA

|                                                |     |
|------------------------------------------------|-----|
| FRESH WATER PROCEDURE . . . . .                | 217 |
| <u>Suspended Solids</u> . . . . .              | 217 |
| <u>Dissolved Solids</u> . . . . .              | 217 |
| BIOTA PROCEDURE . . . . .                      | 218 |
| <u>Fish Flesh</u> . . . . .                    | 218 |
| <u>Fish Bone</u> . . . . .                     | 219 |
| <u>Brush Box Organisms and Algae</u> . . . . . | 220 |

#### APPENDIX "F"

##### CHEMICAL METHODS OF WATER ANALYSIS

|                                                  |     |
|--------------------------------------------------|-----|
| MANGANESE - COLORIMETRIC DETERMINATION . . . . . | 223 |
| <u>Apparatus</u> . . . . .                       | 223 |
| <u>Reagents</u> . . . . .                        | 223 |
| <u>Preparation of Sample</u> . . . . .           | 224 |
| <u>Procedure</u> . . . . .                       | 224 |
| <u>Calculations</u> . . . . .                    | 225 |

|                                                                     | Page |
|---------------------------------------------------------------------|------|
| IRON - COLORIMETRIC DETERMINATION<br>PHENANTHROLINE METHOD. . . . . | 225  |
| Apparatus . . . . .                                                 | 225  |
| Reagents . . . . .                                                  | 225  |
| Preparation of Sample . . . . .                                     | 226  |
| Procedure . . . . .                                                 | 226  |
| Calculations . . . . .                                              | 226  |
| Suggested Standards . . . . .                                       | 227  |
| Notes and Comments                                                  |      |

APPENDIX "G"

BIOLOGICAL METHODS AND TABLE - (COMPLETE LIST OF  
ORGANISMS FOUND IN BRUSH BOX SAMPLES)

|                                                                              |     |
|------------------------------------------------------------------------------|-----|
| PLANKTON COLLECTION AND COUNTING METHODS . . . . .                           | 231 |
| BOTTOM ORGANISM COLLECTION METHODS. . . . .                                  | 231 |
| <u>Petersen Dredge.</u> . . . . .                                            | 231 |
| <u>Brush Boxes.</u> . . . . .                                                | 232 |
| TABLE I - COMPLETE LIST OF ORGANISMS FOUND<br>IN BRUSH BOX SAMPLES . . . . . | 236 |



LIST OF TABLES

| Table No. |                                                                                                         | Page |
|-----------|---------------------------------------------------------------------------------------------------------|------|
| 1.        | Daily Average Savannah River Discharge in cfs for period July 1, 1951 to June 30, 1952 . . . . .        | 30   |
| 2.        | Average, Maximum and Minimum Daily Discharges in cfs for period July 1, 1951 to June 30, 1952 . . . . . | 36   |
| 3.        | Chemical Analysis of Sludge Samples from S-11 and S-14 . . . . .                                        | 57   |
| 4.        | Analytical Data on Samples taken May 26, 1952 from S-239.1, Clark Hill Reservoir . . . . .              | 60   |
| 5.        | Turbidity Values on Samples taken at S-239.1 on March 31 and May 6, 1952 . . . . .                      | 71   |
| 6.        | Clark Hill Reservoir and Savannah River Net Plankton Organisms per Liter - April 1952 . . . . .         | 78   |
| 7.        | Clark Hill Reservoir and Savannah River Net Plankton Organisms per Liter - May 1952 . . . . .           | 80   |
| 8.        | Clark Hill Reservoir and Savannah River Nannoplankton Organisms per 10 Milliliters - May 1952 . . . . . | 84   |
| 9.        | Clark Hill Reservoir and Savannah River Total Phytoplankton Organisms per Liter - May 1953 . . . . .    | 85   |
| 10.       | Organisms Found in Clark Hill Reservoir and Savannah River Plankton - April-May 1952 . . . . .          | 86   |
| 11.       | Algal-Mass Algae . . . . .                                                                              | 90   |
| 12.       | Animals Associated with Algal Masses . . . . .                                                          | 92   |
| 13.       | Quantitative Clay Bank Sample S-52.0 - Jul, 30, 1952. . . . .                                           | 99   |
| 14.       | Bottom Samples. - Summary . . . . .                                                                     | 102  |
| 15.       | Brush Box Sampling - Summary . . . . .                                                                  | 110  |
| 16.       | Basket Fishery - Commercial Catches Examined . . . . .                                                  | 126  |
| 17.       | Shad Fishery - Commercial Catches Examined . . . . .                                                    | 127  |
| 18.       | Composition of Hoop Net Catches - S-172.5 - S-140.0, December 3, 1952 - July 12, 1952 . . . . .         | 128  |

| Table No. |                                                                                                      | Page |
|-----------|------------------------------------------------------------------------------------------------------|------|
| 19.       | Comparison of Hoop Net and Commercial Catches . . . . .                                              | 129  |
| 20.       | Results of Catfish Marking Experiment -<br>May 8 - July 12, 1952 . . . . .                           | 139  |
| 21.       | Results of Rotenone Sampling . . . . .                                                               | 140  |
| 22.       | Estuarine Sampling - April 22-24, 1952 -<br>Analytical Results . . . . .                             | 153  |
| 23.       | Estuarine Collections, April 1954 . . . . .                                                          | 154  |
| 24.       | Radioactivity of Savannah River Water Samples . . . . .                                              | 166  |
| 25.       | Radioactivity of Fresh Water Fish - 1951-1952 . . . . .                                              | 167  |
| 26.       | Radioactivity of Fresh Water Minnows and Algae -<br>1951-1952 . . . . .                              | 169  |
| 27.       | Radioactivity of Fresh Water Brush Box Organisms -<br>1951-1952 . . . . .                            | 170  |
| 28.       | Radioactivity of Crabs, Oysters and Shrimp found in<br>Savannah River Estuaries April 1952 . . . . . | 171  |
| 29.       | Radioactivity of Fish found in the Savannah River<br>Estuaries April 22-24, 1952 . . . . .           | 172  |
| 30.       | The Maximum Radioactive Density of the Savannah<br>River Samples - 1951-1952 . . . . .               | 174  |

LIST OF FIGURES

| Figure No. |                                                                                  |    |
|------------|----------------------------------------------------------------------------------|----|
| 1.         | Map of Savannah River - Sampling Locations, Clark<br>Hill Dam to Mouth . . . . . | 15 |
| 2.         | Flow Investigation - Abercorn Creek, Effingham<br>County, Georgia . . . . .      | 31 |
| 3.         | Dissolved Oxygen - Mean and Minimum . . . . .                                    | 41 |
| 4.         | Biochemical Oxygen Demand, 5-day @ 20°C - Mean<br>and Maximum . . . . .          | 41 |
| 5.         | Coliform MPN per 100 ml. - Mean, Maximum and<br>Minimum . . . . .                | 41 |



| Figure No. |                                                                                                               | Page |
|------------|---------------------------------------------------------------------------------------------------------------|------|
| 6.         | Savannah River Estuary - Location of Sampling Stations . . . . .                                              | 47   |
| 7.         | Dissolved Oxygen - High Tide - March 28, 1952 . . . . .                                                       | 48   |
| 8.         | M. O. Alkalinity (ppm), ‰ Seawater, - High Tide - March 28, 1952 . . . . .                                    | 48   |
| 9.         | Dissolved Oxygen - Between Mean and High Tide - March 28, 1952 . . . . .                                      | 48   |
| 10.        | Per cent Seawater - Between Mean and High Tide - March 28, 1952 . . . . .                                     | 49   |
| 11.        | Dissolved Oxygen - Mean Tide - March 28, 1952 . . . . .                                                       | 51   |
| 12.        | Per cent Seawater - Mean Tide - March 28, 1952 . . . . .                                                      | 51   |
| 13.        | Dissolved Oxygen - Low Tide - March 28, 1952 . . . . .                                                        | 51   |
| 14.        | M. O. Alkalinity (ppm), ‰ Seawater, - Low Tide - March 28, 1952 . . . . .                                     | 51   |
| 15.        | Dissolved Oxygen, Surface Samples - High and Low Tide - June 23, 1952 . . . . .                               | 53   |
| 16.        | Dissolved Oxygen, Bottom Samples - High and Low Tide - June 23, 1952 . . . . .                                | 53   |
| 17.        | Dissolved Oxygen, Bottom and Surface Samples - High Tide - June 23, 1952 . . . . .                            | 53   |
| 18.        | Dissolved Oxygen, Bottom and Surface Samples - Low Tide - June 23, 1952 . . . . .                             | 53   |
| 19.        | Chlorides - ‰ Seawater - High and Low Tide - June 23, 1952 . . . . .                                          | 55   |
| 20.        | Clark Hill Reservoir Sample Station Location Map . . . . .                                                    | 61   |
| 21.        | Dissolved Oxygen - Depth Relation (1952) Clark Hill Reservoir - March 10, March 31, May 1 and May 3 . . . . . | 63   |
| 22.        | Dissolved Oxygen - Depth Relation (1952) Clark Hill Reservoir - May 6 . . . . .                               | 63   |
| 23.        | Dissolved Oxygen - Depth Relation (1952) Clark Hill Reservoir - May 13, May 21, May 26 . . . . .              | 63   |

## Figure No.

- 1 -

## Page

|     |                                                                                                                                    |     |
|-----|------------------------------------------------------------------------------------------------------------------------------------|-----|
| 24. | Dissolved Oxygen - Depth Relation (1952) Clark Hill Reservoir - June 13, July 28 . . . . .                                         | 63  |
| 25. | Daily Average Water Temperature of Augusta Canal, and Daily Flow of Savannah River - January 22, 1951 - January 21, 1952 . . . . . | 65  |
| 26. | Comparison of Air and Water Temperatures, Augusta Canal - February 4, 1951 - February 3, 1952 . . . . .                            | 67  |
| 27. | Distribution of Water Temperature - Augusta Water Intake - February 4, 1951 - February 3, 1952 . . . . .                           | 69  |
| 28. | Length-Frequency Distribution - Channel Catfish - August 1951 - May 1952 . . . . .                                                 | 119 |
| 29. | Length-Frequency Distribution - White Catfish - August 1951 - July 1952 . . . . .                                                  | 121 |
| 30. | Length-Frequency Distribution - Flat Bullheads - June 1951 - July 1952 . . . . .                                                   | 121 |
| 31. | Length-Frequency Distribution - American and Hickory Shad - February-April 1952 . . . . .                                          | 143 |
| 32. | Length-Frequency Distribution - Silvery Minnow - May 14 - July 3, 1952 . . . . .                                                   | 145 |
| 33. | Length-Frequency Distribution - Spottail Shiner . . . . .                                                                          | 145 |
| 34. | Length-Frequency Distribution - Pirate Perch . . . . .                                                                             | 145 |
| 35. | Length-Frequency Distribution - Bluegill - May-July 1952 . . . . .                                                                 | 147 |

- 1 -

LIST OF CONTRIBUTORS

The Savannah River Studies were performed under the general administrative direction of V. G. MacKenzie, Officer in Charge, and under the technical direction of P. D. Haney, Director, Field Activities, and the late C. C. Ruchhoft, Chief, Physics and Chemistry Section, all of the Robert A. Taft Sanitary Engineering Center (formerly the Environmental Health Center), Cincinnati, Ohio.

Personnel assigned to the field unit at Augusta, Georgia, were as follows:

- D. D. Gold, Sanitary Engineer, in Charge
- D. C. Scott, Aquatic Biologist
- J. M. Cohen, Chemist
- C. N. Shadix, Chemist (August 1951 - January 1952)
- A. L. Platz, Assistant Sanitary Engineer (July 1951 - February 1952)
- J. P. Sheehy, Assistant Sanitary Engineer (February 1952 - June 1952)
- Allan Hirsch, Assistant Sanitarian
- J. W. Coleman, Aide

Personnel contributing to the studies at the Robert A. Taft Sanitary Engineering Center, Cincinnati, Ohio, were as follows.

- C. M. Tarzwell, Chief, Aquatic Biology
- L. R. Setter, Assistant Chief, Physics and Chemistry
- A. S. Goldin, Radiochemist
- J. S. Nader, Physicist
- C. R. Hagee, Physicist
- R. J. Velten, Chemical Aide

ACKNOWLEDGMENTS

Valuable assistance has been rendered to the Savannah River Studies group by various municipal, State and Federal Agencies, and by the DuPont Company. Special acknowledgment is made to the following:

Georgia Department of Public Health

Mr. W. H. Weir, Director, Water Pollution Control

South Carolina State Board of Health

Mr. William T. Linton, Executive Director, Water Pollution Control Authority

State of Georgia Military Department

For the use of their buildings and grounds in Augusta, Georgia City of Augusta

Former Mayor, Dr. Will D. Jennings, and present Mayor,

Mr. Hugh Hamilton

E. I. DuPont de Nemours and Company

Mr. R. R. Balmer, Engineering Service Division, Savannah River Plant

Atomic Energy Commission, Savannah River Plant

Mr. G. H. Giboney, and Mr. Karl Herde, Technical and Production Division

U. S. Corps of Engineers, Savannah District

Mr. J. W. Harris, (now on active duty with the U. S. Navy),

Chief, River and Harbors Section (1951), and Mr. A. W. Carlson,

Chief, River and Harbors Section (1952).

U. S. Geological Survey

Mr. A. E. Johnson, District Engineer

U. S. Public Health Service, Communicable Disease Center

Dr. S. W. Simmons, Officer in Charge, Technical Development  
Branch

Savannah River Advisory Board Members

Georgia Department of Public Health

Mr. W. H. Weir, Director, Water Pollution Control  
Atlanta, Georgia

South Carolina State Board of Health

Mr. William T. Linton, Executive Director, Water Pollution  
Control Authority, Columbia 1, South Carolina

E. I. DuPont de Nemours and Company

Dr. J. E. Cole, Director, Manufacturing Division,  
Atomic Energy Division, Wilmington 98, Delaware

Atomic Energy Commission

Mr. Arthur E. Gorman, Sanitary Engineer, Division of  
Engineering, Washington 25, D. C.

Dr. Paul B. Pearson, Chief, Biology Branch, Division of  
Biology and Medicine, Washington 25, D. C.

U. S. Corps of Engineers

Mr. Harry Sugden, Savannah District, P. O. Box 889,  
Savannah, Georgia

U. S. Geological Survey

Mr. Albert E. Johnson, District Engineer, 207 Creason  
Building, Columbia 1, South Carolina

U. S. Public Health Service

Mr. Paul D. Haney, Director of Field Activities, Environ-  
mental Health Center, 1014 Broadway, Cincinnati 2, Ohio

Mr. C. C. Ruchhoft, Chief, Chemistry and Physics Section,  
Research and Development Branch, Environmental Health  
Center, 1014 Broadway, Cincinnati 2, Ohio

Mr. James G. Terrill, Jr., Acting Chief, Radiological  
Health Branch, Division of Engineering Resources,  
Washington 25, D. C.

Mr. L. F. Warrick, Chief, Technical Services Branch, *(Chairman)*  
Division of Water Pollution Control, Washington 25, D. C.

Mr. Lewis A. Young, Officer in Charge, Southeast Drainage  
Basin Office, Division of Water Pollution Control,  
50 Seventh St., N.E., Atlanta 5, Georgia

## CHAPTER I

### SUMMARY OF FINDINGS AND RECOMMENDATIONS

This study of the Savannah River extended from July 1951 into July of 1952. Most of the work was confined to the area just downstream (mile 230.7) from Clark Hill Dam to the river's mouth (mile 0.0). Situated in this reach of the Savannah River is the tract of land that was selected for the Savannah River plant of the Atomic Energy Commission. This tract comprises 250,000 acres in Aiken, Allendale and Barnwell Counties, South Carolina, about 25 miles downstream from Augusta, Georgia. Data were gathered for the most part before the impoundment of the Savannah River by Clark Hill Dam.

The principal objectives of the survey were to determine: (a) the water quality characteristics of the stream, (b) the background radio-activity of water and biota and (c) the ecology of the biota and its economic importance before initiation of operations at the Savannah River plant of the Atomic Energy Commission and before impoundment of water by Clark Hill Dam.

### FINDINGS AND CONCLUSIONS

The findings and conclusions reached are as follows:

1. Savannah River water is soft and low in mineral content throughout the year except for the estuarine section of the river which is mixed with sea water.
2. For almost 20 per cent of the days, or more than 60 days between February 4, 1951 to February 3, 1952, the water temperature



at the Augusta Water Intake was in the 82° to 86° F. temperature range. For less than ten days the water temperature was between 62° to 66° F. There were only nine days in the 38° to 42° F. temperature range.

3. The red muddy appearance of the Savannah River, which gives the illusion of high turbidity, is caused by colloidal red clay. The water on examination in the laboratory is not nearly as turbid as the stream appearance suggests, since the colloid affects "apparent color" rather than turbidity. The minimum mean turbidity value for all the Savannah River samples was 62 ppm and the maximum mean turbidity value was 115 ppm at the various stations. Single turbidity values ranged from a minimum of 8 ppm to 440 ppm.
4. Thirty-two water samples from seven Savannah River sampling stations and four tributaries were analyzed for the gross alpha and beta radioactivity associated with the suspended solids (insoluble) and in solution (soluble). The insoluble alpha activity varies from zero to a maximum of  $4.1 \pm 1.7$  micromicrocuries per liter of water. The soluble alpha activity of the sample showing the maximum insoluble activity was  $0.1 \pm 1.1$   $\mu\text{mc/liter}$ . Thus the total alpha activity was  $4.2 \pm 2.0$   $\mu\text{mc/liter}$  of water. Usually there was more insoluble than soluble alpha activity in the samples. However, in six out of 31 samples the soluble alpha activity exceeded the insoluble activity. The maximum soluble alpha activity was  $8.4 \pm 2.1$   $\mu\text{mc/l}$ . compared to an insoluble alpha activity of 0.4  $\mu\text{mc/l}$ .

5. The beta activity was appreciably higher in magnitude than the alpha activity. This is particularly true with respect to soluble activity. Thus the maximum soluble beta activity was  $31.8 \pm 14.8$   $\mu\text{mc}/\text{l}$ . This sample had an insoluble beta activity of only  $8.6 \pm 5.1$   $\mu\text{mc}/\text{l}$ . The maximum total beta activity occurred for this sample and was  $40.4 \pm 15.6$   $\mu\text{mc}/\text{l}$ .
6. Radioactive measurements were determined on 61 fresh water biological samples. These samples consisted of 40 individual or composite flesh and bony tissue samples, and composite samples of whole fish minnows, 7 samples of brush box organisms (largely insect larvae), and 7 samples of algae. The statistically significant results showed that the radioactivity of all samples exclusive of fish bone was  $47 \pm 28$   $\mu\text{mc}$  or less of alpha activity and  $14.3 \pm 5.6$   $\mu\text{mc}$  or less of beta activity per gram of wet weight sample. The minnows (whole fish samples) and the insect larvae, all of which samples probably contained soil, showed a significant alpha activity. The maximum alpha activity of fish flesh was  $1.5 \pm 0.9$   $\mu\text{mc}/\text{gram}$ . The radioactivity of the ash of fish bone was less than the statistical error of the determination which varied from  $\pm 2$  to  $\pm 28$   $\mu\text{mc}/\text{gram}$  for the alpha activity and  $\pm 20$  to  $\pm 260$   $\mu\text{mc}/\text{gram}$  for beta activity. The algae showed no significant amount of alpha activity but a measurable beta activity.

7. Radioactive analyses were made on samples of flesh of estuarine animals, which included: 9 composite samples of crabs, 8 of shrimps, 2 of oysters, and 32 of fishes; and the bone or shell of 17 composite samples from various organisms. The maximum alpha activity in muscular tissues was  $0.22 \pm 0.17 \mu\text{mc/g.}$  in an oyster sample. All but one composite flesh sample had a small but significant beta activity which may be attributed to  $\text{K}^{40}$ . The beta activity varied from  $0.5 \pm$  to  $0.2 \mu\text{mc/g.}$  in an oyster sample to  $4.5 \pm 1.1 \mu\text{mc/g.}$  in a fish sample.
8. Plankton populations in the Savannah River were quite low. They did not appear to be significantly influenced by the plankton which developed in the Clark Hill Reservoir or by the organic enrichment afforded by Augusta sewage. Some plankton organisms, particularly diatoms, appeared either to multiply in the river itself or to be added to the river from backwater areas. The general paucity of river plankton can probably be attributed to the scarcity of inorganic nutrients and the low transparency of the water. If sedimentation in Clark Hill Reservoir results in some clarification of the river water, the plankton populations of the river may be expected to increase.
9. Benthic algal growths and their associated organisms formed transient communities of seasonal importance. These could only maintain themselves when river levels remained constant

over periods of several weeks or when they were formed on substrates which could rise and fall with the water. Such communities of algae, herbivores and carnivores might well become of much greater importance if stable water levels and clearer waters prevail over considerable parts of the year.

10. The hard clay banks which form a large part of the river's edge supported dense populations of nymphs of burrowing mayflies (Tortopus). These insects and the organisms associated with them formed an important part of the biota of the river.
11. The sand and mud of the river bottom proper, the most abundant substrates available to benthic organisms, supported the most sparse populations. Much of this bottom material was in an unstable condition which prevented the establishment of most bottom organisms. In areas of strong current, these materials were constantly shifting, while in still waters the bottom was constantly subjected to siltation. The organic enrichment supplied by Augusta sewage provided a habitat for the development of a fauna of pollution tolerant forms immediately below the mouth of Butler Creek. General observations, at periods of low water, indicated that mussels were important constituents of the benthic fauna.
12. Wherever stable substrates, organic or inorganic in nature, were present in the river current, a diverse population of organisms became established. Since the river was abundantly provided with such substrates in the form of brush, trees,

roots, etc., such habitats were considered to be among the most important encountered. Organisms that were found in them consisted largely of detritus feeders and carnivores.

13. Commercial fishing on the Savannah River was quite limited in extent. A small basket fishery for channel catfish and a small seasonal gill net fishery for shad were established. Although it was not possible to assess the absolute value of these fisheries, they were obviously of negligible economic importance to the surrounding region. Sport fishing while indulged in by a few enthusiasts was of little importance on the river. Tributaries and the flood plain lakes were apparently more heavily fished.
14. The fish population of the Savannah River was typical for a muddy river of the southeast Atlantic Coast. The channel catfish was one of the most important constituents of the population. Two other catfish were also abundant, the white catfish and the flat bullhead. Suckers were moderately abundant, as were carp. A number of forage minnows was present. Among the spiny-rayed fish the sunfishes, largemouth bass and both black and white crappies were all present in moderate numbers. The American and hickory shads and the striped bass were the most important anadromous fishes to utilize the river. A total of 60 species of fish known to occur in fresh waters of the Savannah River are reported here.

15. With the exception of the Savannah Harbor proper, which is degraded from a combination of pollution and silt, the estuarine waters around the mouth of the river are quite typical of this stretch of coast. Shrimp and crabs are taken commercially in the immediate vicinity of the river mouth. Sport fishing for various estuarine fishes is also of considerable importance in this area. Oysters, while abundant at the river mouth, are contaminated by sanitary wastes and are not utilized.
16. Raw sewage from Augusta, Georgia, is discharged into Beaver Dam Ditch where it receives some natural treatment when the Ditch passes through Phinizy swamp to join Butler Creek, about 300 yards above the confluence of Butler Creek and Savannah River. Even in the summer months during low flow there was sufficient dilution of swamp treated Augusta sewage so that but little oxygen depletion occurred in the Savannah River below Augusta, Georgia.
17. During this period of the studies the sewered population was about 3500 persons in North Augusta, South Carolina. The Savannah River, at low flow, affords sufficient dilution so that little effect is noticed on the oxygen resources of the river.
18. The Savannah River downstream from Savannah, Georgia, is sometimes entirely depleted of its dissolved oxygen content, which results in an "oxygen block" to fish during the summer months. Domestic pollution from the city of Savannah, Georgia, and



industrial pollution from paper and pulp mills located on the Georgia side of the river above Savannah are important factors in the depletion of dissolved oxygen in the river below Savannah, Georgia. There are extensive mobile sludge banks up to 12 feet in depth in the Savannah River section, between mile 20 to mile 3. These sludge banks contain from 10 to 30 per cent volatile matter on a dry weight basis.

#### RECOMMENDATIONS

The observations and findings from these studies indicate a need for continued or new work, as follows:

1. Efforts should be made to determine the effect the Atomic Energy Commission's Savannah River Plant may have on the radioactivity of the Savannah River waters and its biota.
2. Possible increases in water temperature with the addition of quantities of heat from cooling waters released by the Savannah River plant may be either beneficial or detrimental to the total biota of the river immediately below the plant outfall. The effects of increased river water temperature on the biota in the river should be studied.
3. Impoundment of Savannah River water by Clark Hill Dam may induce appreciable changes in water quality and quantitative as well as qualitative changes in river biota. Studies should be performed to ascertain such changes, since the quality of the Savannah River water is of vital importance to industries and municipalities as a source of water supply.



4. Public concern and apprehension regarding waters which receive cooling water and/or waste discharge from atomic energy plants necessitate that a record of studies by an agency other than the Atomic Energy Plant or Operations Management should be available for informational use by State Health Departments and other agencies of government. Continuing short studies at reasonable intervals are desirable to satisfy the need for protecting public welfare and public relations.
5. More accurate as well as more simplified radiochemical analytical techniques and counting equipment are needed for determining low levels of radioactivity.
6. Studies of a research nature are needed to develop information relating to (1) - the effect of silt load in rivers upon adsorption of radioactive substances, (2) - radioactive adsorption characteristics of various types of silt commonly present in rivers, and (3) - the effect of river eddies and currents on time-distance relations of silt load deposition.

## CHAPTER II

GENERAL INFORMATION

The Savannah River studies were planned and every effort made to conduct them in such a manner as to meet the needs and responsibilities of the States involved, the Atomic Energy Commission, and the Public Health Service. Because the river is vital to the operations of the Savannah River plant of the Atomic Energy Commission, and serves as a source of water supply for the city of Augusta and as a partial source of water supply for the city of Savannah, it is important to have a full knowledge of the river and its changing characteristics.

## OBJECTIVES

The main objectives of the Public Health Service program were as follows:

1. To determine hydrological, physical, chemical, radiological and biological characteristics of the Savannah River above and below the Atomic Energy Commission's Savannah River plant, including the area above and below Clark Hill Dam.
2. To develop new and to improve existing techniques as applied to stream sanitation objectives.
3. To establish the biological conditions prevailing in the Savannah River so that future changes resulting from impoundment behind Clark Hill Dam, operation of the Atomic

Energy Commission's Savannah River plant and attendant increased population can be detected and measured.

4. To determine the relative abundance of the various species of fish in the river and the extent to which they were being utilized for sport or commercial purposes.

A comparison of the results of future investigations with those of the present one should permit the detection and evaluation of the environmental changes caused by anticipated activities in the Savannah River watershed.

#### INITIATION OF STUDIES

In July 1951, the Surgeon General of the U. S. Public Health Service formed the Savannah River Studies Advisory Board by appointing representatives from the States of Georgia and South Carolina, the E. I. duPont de Nemours and Company, the Atomic Energy Commission, the U. S. Army Corps of Engineers, the U. S. Geological Survey, and the U. S. Public Health Service to the status of Board Members. The function of the Savannah River Studies Advisory Board is to review the data obtained by the survey group and to aid in planning continuing studies on the Savannah River.

The field work of the Savannah River Studies' program was initiated on July 1, 1951, with the basing of two mobile laboratory units in the Georgia National Guard area in Augusta, Georgia. By the end of July conversion of a building in the area into a laboratory was completed and operations were transferred from the mobile units to the building.

#### SAMPLING PROGRAM


A mileage index system, listed below, was used to identify the location of sampling stations, landings and other important points on the Savannah River and its tributaries. This system involves the use of one or more letters to identify the stream, followed by a number which represents the distance in river miles from the mouth. On the Savannah River, the letter "S" designates the stream. On the tributaries the first and second letters of the name of the stream are used to designate the stream, with the exception of Upper Three Runs and Lower Three Runs for which the three letters "UTR" and "LTR" are used, respectively. The first and second letters of the tributaries are preceded with the notation, "S- confluence mile." For example, the sampling station on Steel Creek would be designated on the mileage index system as S-155.2, St.-.01. This indicates that the station is located on Steel Creek .01 mile from its confluence with the Savannah River and the confluence is 155.2 miles from the mouth of the Savannah River. If there were two Steel Creeks, one in Georgia and one in South Carolina, the stream designation would have been followed by the abbreviation of the name of the State in which it was flowing, e.g., Steel Creek in South Carolina would be designated as S-155.2, St(S.C.)-.01.

LOCATION BY RIVER MILES OF SAMPLING STATIONS,  
LANDINGS, TRIBUTARIES, ETC.

|          |                                                                    |
|----------|--------------------------------------------------------------------|
| S-246    | Sample Station in Clark Hill Reservoir, 8 miles above Dam          |
| S-239.1  | Sample Station at confluence of Little River with Savannah River   |
| S-238.1  | Clark Hill Dam                                                     |
| S-230.7  | Woodlawn Landing (Railroad Bridge), S. C.                          |
| S-218.5  | Sample Station just above Rae's Creek, Ga.                         |
| S-217.8  | Sibley Mill Gate, Ga.                                              |
| S-217.6  | King Mill Gate, Ga.                                                |
| S-217.2  | Hawks Gully Gate, Ga.                                              |
| S-216.5  | North Augusta                                                      |
| S-216.0  | Augusta                                                            |
| C-215.79 | Southern Railway Bridge                                            |
| S-215.64 | Fifth Street Bridge (USGS Alternate Gage)                          |
| S-215.6  | Fifth Street Landing, Ga.                                          |
| S-213.36 | Horse Creek, S. C.                                                 |
| S-212.0  | Sample Station 1/4 mile above Sand Bar Ferry Road Bridge           |
| S-211.73 | Sand Bar Ferry Road Bridge                                         |
| S-211.38 | C & W C Railway Bridge                                             |
| S-202.97 | New Savannah Bluff Lock and Dam                                    |
| S-202.9  | Sample Station just below New Savannah Bluff Lock and Dam          |
| S-202.65 | Butler Creek, Ga. (USGS Augusta Gage)                              |
| S-201.65 | 1 mile below Butler Creek                                          |
| S-198.95 | Thomas Creek, Ga.                                                  |
| S-198.42 | Spirit Creek, Ga.                                                  |
| S-197.0  | Sample Station                                                     |
| S-191.8  | Hollow Creek, S. C.                                                |
| S-189.4  | Silver Bluff Landing, S.C. (Good landing and access road)          |
| S-186.57 | Bridge Creek, Ga.                                                  |
| S-178.57 | McBean Creek, Ga.                                                  |
| S-176.66 | Boggy Gut Creek, Ga.                                               |
| S-176.4  | Shell Bluff Landing, Ga.                                           |
| S-172.0  | Sample Station                                                     |
| S-171.59 | Upper Three Runs, S. C.                                            |
| S-165.55 | Hancock Landing, Ga. (Good landing with boom & winch, access road) |
| S-164.26 | Fourmile Branch, S. C.                                             |
| S-161.98 | Joblers Creek, Ga.                                                 |
| S-157.3  | Brighams Landing, Ga. (Good landing and access road)               |
| S-155.2  | Steel Creek, S. C.                                                 |
| S-152.0  | Sample Station                                                     |
| S-146.4  | Entrance to Dead River (Little Hell Landing), S. C.                |
| S-145.38 | Sweetwater Creek, Ga.                                              |
| S-144.65 | Stony Bluff Landing, Ga. (Good landing and access road)            |
| S-137.32 | Lower Three Runs, S. C.                                            |
| S-131.27 | Rocky Creek, Ga.                                                   |
| S-129.25 | Highway 301 Bridge (Dorton's Ferry Bridge, USGS Gage)              |

|           |                                                                     |
|-----------|---------------------------------------------------------------------|
| S-128.23  | Brier Creek, S. C.                                                  |
| S-124.76  | Swift Creek, Ga.                                                    |
| S-113.7   | Cohens Bluff Landing, S. C.                                         |
| S-106.8   | Swift Creek, Ga.                                                    |
| S-105.5   | Brier Creek and Landing, Ga.                                        |
| S-98.93   | Buck Creek, Ga.                                                     |
| S-91.7    | Poor Robin Landing, Ga.                                             |
| S-82.53   | Black Creek, Ga.                                                    |
| S-82.19   | Blue Springs Creek, Ga.                                             |
| S-81.2    | Hudsons Ferry, Ga.                                                  |
| S-68.0    | Bostick and Stokes Landing, S. C.                                   |
| S-64.9888 | S.A.L. Railway Bridge (USGS Clys Gage)                              |
| S-52.0    | Sample Station                                                      |
| S-48.2    | Ebenezer Creek, Ga.                                                 |
| S-48.1    | Ebenezer Landing, Ga.                                               |
| S-46.33   | Lockner Creek, Ga.                                                  |
| S-44.74   | Mill Creek, Ga.                                                     |
| S-43.24   | Big Collis Creek, Ga.                                               |
| S-42.4    | Bear Creek, Ga.                                                     |
| S-36.2    | Purysburg, S. C.                                                    |
| S-29.83   | Big Collis Creek, Ga.                                               |
| S-27.36   | A.C.L. Railway Bridge                                               |
| S-22.0    | Highway Bridge Landing (Hy.17); Wild Life Indg. on Back River, S.C. |
| S-16.5    | Seaboard RR Bridge                                                  |
| S-14.5    | Foot of Bull St., Ga. (Corps of Engineers Dock & Laboratory)        |
| S-11.0    | Sample Station 1/4 mile below Fort Jackson                          |
| S- 0.5    | Old Quarantine Station, Cockspur Island                             |
| S- 0.0    | Vicinity of Fort Pulaski                                            |





SAVANNAH RIVER  
 CLARK HILL DAM TO MOUTH

FIG.1

The survey group concentrated on the following stations: S-218.5; S-202.9; S-197; S-172; S-152; S-11 and four creeks, namely, S-164.26, Mo-.01; S-155.2, St-.01; S-151.29, UTR-.01 and S-137.52, LTR-5.75. Other sampling stations were used occasionally for chemical, bacteriological and biological samples. Samples were taken at 1/3, 1/4, or 1/2 points across the river for bacteriological determinations. In the early stages of the survey, samples for chemical determinations were taken at 1/3, 1/4, or 1/2 points across the river at sample stations, but this practice was discontinued when the chemical determinations showed no variation in such sampling. Samples for chemical determinations were subsequently taken in mid-stream at sampling locations.

Routine mineral analyses were performed at the Augusta laboratory on the same sampling schedule as for samples sent to the Environmental Health Center laboratory at Cincinnati.

Sampling for radiological background and spectrographic analysis was discontinued April 10, 1952 when it was apparent that sufficient samples for background radioactivity had been obtained.

Biological sampling stations were selected in the vicinity of the chemical and bacteriological sampling stations, where practicable. However, because the location of organism concentrations is dependent upon various river features such as bars, velocity, currents, depth, etc., the selection of biological sampling stations was primarily governed by these features.

Sampling was conducted to secure a representative portion of water at each sampling station. The sample was drawn from midstream at a depth of three feet. A copper depth sampler, which held special bottles for dissolved oxygen and bacteriological samples, was used. When water samples for only bacteriological determinations at 1/4 or 1/3 points across the stream were collected, a pole-type dip sampler holding a sterile bottle was used. A Kemmerer bottle was used to obtain water samples for chemical and physical analyses in the deep waters in Clark Hill Reservoir and in the lower Savannah River.

Samples were transported in special ice chests from the sample locations to the laboratory. The ice chests were of such construction that no sample was in direct contact with the ice. The bottles containing water samples were supported in an upright position in the ice chest. The samples were brought to the laboratory as quickly as possible so that analyses could be initiated with a minimum elapsed time.

#### LABORATORY ACTIVITIES

During the early stages of the studies, chemical and bacteriological determinations were performed in two mobile trailer units. Later, laboratory facilities were constructed in a building in the National Guard area at No. 1 Milledge Road, Augusta, Georgia. The arrangements for the use of this building were made through the Georgia Department of Public Health. Laboratory operations were transferred from the trailer units to the building during September 1951. The Savannah River studies group was also granted the use of a large brick building for repair and storage facilities and two open walled sheds for drying fish nets and for gasoline storage.

The Corps of Engineers, Savannah District, furnished the Savannah River studies group the use of facilities in their laboratory located on their dock at the foot of Bull Street in Savannah. They also made available the use of their dock and boat launching equipment for sample collection activities.

Facilities suitable for a field laboratory were made available by personnel attached to the Fort Pulaski National Monument on Cockspur Island.

A total of 490 water samples from the Savannah River and tributary streams was taken for routine chemical and physical analysis on which 4,014 determinations were performed; 603 water samples were collected for bacteriological analysis and 603 determinations were performed. In addition, 32 water samples were shipped to the Environmental Health Center laboratories in Cincinnati, Ohio, for radioactive background counts and spectrographic analysis. A total of 38 water samples was taken for mineral analysis, on which a total of 582 determinations were performed. Numerous other samples were collected for dissolved oxygen and salinity studies in the lower Savannah River and for the Clark Hill Reservoir studies on which hundreds of determinations were performed.

#### SPECIAL STUDIES

Special studies that were carried out to supplement routine data collections and analyses are listed below:

1. Several dissolved oxygen and salinity profiles on the lower Savannah River (mile 23 to mile 0) were made at various tidal stages and seasons.

2. Many dissolved oxygen, temperature, iron and manganese profiles (depth) were made in Clark Hill Reservoir starting on March 10, 1952.

3. Studies in estuarine waters relating to shellfish, fish and other biological organisms were performed. Some chemical determinations were performed as a supplement to the biological studies in these waters.

4. Rotenone was used for the collection of fish not obtainable by seining or trapping.

## CHAPTER III

GENERAL DESCRIPTION

The Savannah River basin is predominantly a rural area, with about 20 per cent of the population residing in Augusta and Savannah. The principal uses of the basin water resources include the generation of hydroelectric power, shellfish production, and domestic and industrial water supplies. The Basin's watercourses receive domestic, raw and treated, sewage from 53 municipalities having a sewered population in excess of 240,000. It also receives industrial wastes from two pulp and paper plants which are estimated to have a population equivalent in excess of 200,000.

THE SAVANNAH RIVER BASIN

The drainage basin covers some 10,500 square miles of Georgia, South Carolina, and a small section of North Carolina.

The extreme headwaters of the Savannah River lie in the Blue Ridge Province of North Carolina, South Carolina, and Georgia, a heavily wooded mountainous area at altitudes as high as 5,000 feet. The Savannah River is formed by the confluence of the Tugaloo and Seneca Rivers near Hartwell, Georgia, and flows some 300 miles to discharge into the Atlantic Ocean near Savannah Beach, Georgia.

TOPOGRAPHY AND GEOLOGY (Clark Hill Dam to Atlantic Ocean)

From the foot of Clark Hill Dam to Augusta, a distance of some 18 miles, the Savannah River drops about 70 feet in elevation over a rocky bed. Its course is obstructed by two dams, Stephen's Creek Dam (29 feet in height) and the Augusta City Dam (10 feet in height). Because of the steep



gradient of the river in this stretch the backwaters of neither of these dams is very extensive. The backwater at the Stephen's Creek Dam has been so extensively silted in that a short distance above the dam it is only a few feet deep. Below Augusta at mile S-202.95, the New Savannah Bluff Lock and Dam creates a backwater which extends to mile S-219, just above Augusta. Here the last, naturally occurring, igneous rock is exposed in a series of rapids. From the foot of the New Savannah Bluff Dam to the region of marked tidal influence, at about mile S-30, the character of the river changes but little. The Sparganium marsh which is characteristic of the seacoast extends up the river to Big Collis Creek at mile S-29.83 where it ends abruptly. This may indicate the limit of regular tidal influence. However, a relatively wide range of tidal fluctuations at the Savannah River entrance sometimes causes noticeable semi-diurnal variations of water level to occur as much as fifty miles up the channel from the sea. The extent of the tidal influence depends upon the relative magnitude of the tide and the river flow. There are no tributaries entering the river in the Coastal Plain whose average flow is greater than 700 cfs. Over this 200-mile stretch the river follows a generally tortuous, meandering channel through a densely wooded 3 to 5 miles wide flood plain. The river bottom is composed largely of shifting sand; however, there are a number of places where the river swings to the edge of its flood plain and encounters the sedimentary formations of the Coastal Plain. At these points a variety of types of bottom materials are encountered, including clay, marl, gravel, sandstone and limestone. At meanders, where the river is cutting away at its own flood plain, the usual caving, vertical or undercut, clay bank

occurs on the outer edge of the meander. Banks may drop precipitously below the normal water level to the deepest part of the channel (often 10 to 12 feet at low water). From this maximum depth the river bed rises gradually to the water level on the inner side of meanders. The familiar pattern of erosion of the outer bank and silt deposition on the inner one is everywhere present. The successional nature of the sand bars is evident and they can be observed in all stages of deposition, overgrowth by willow, and conversion into the river swamp.

#### POPULATION

The Savannah River basin is predominantly a rural area having a 1940 population of 939,353, which represents an average density of 62 persons per square mile. About seventeen per cent of the total population (1940) reside in Augusta and Savannah, the basin's major cities. According to the 1950 census population figures for Savannah and Augusta are 119,669 and 71,507, respectively.

#### LAND USE

Agriculture is currently the chief economic activity of the basin. The farms of the region produce a large variety of products which include cotton, corn, tobacco, sugar cane, peanuts, vegetables, fruit, livestock, and poultry. Manufacturing activities are confined chiefly to the production of textiles, textile materials, paper, paper products, clay products and naval stores.

#### WATER USE AND DEVELOPMENT

##### Fish and Wildlife

Extensive areas of the lower Savannah drainage basin are utilized

for feeding, nesting and breeding areas by migratory wildfowl. The Savannah Bird Refuge is located on the delta lands in the lower river above Savannah, and the Tybee National Wildlife Refuge is located on the delta below Savannah.

The main stem of the Savannah River and its network of tributaries maintains both commercial and recreational fishing. However, the most important fisheries resource is the basin's shellfish industry. The waters of the minor delta channels in the vicinity of the mouth of the Savannah River are utilized as a nursery ground for young shrimp.

#### Recreation

Because of the climate and the natural and developed recreational areas, the region above Augusta is attractive to tourists and vacationists. Two of the principal storage reservoirs of the Tallulah River power system have been developed as recreational areas. The Savannah River between Augusta and Savannah has some recreational value. Below the city of Savannah the beaches on Tybee Island provide attractive sea bathing facilities.

#### Reservoirs and Diversion Projects

##### Clark Hill

The Clark Hill Project, a multi-purpose reservoir, has recently been completed except for the power house which is now under construction. The top five feet of the reservoir, elevation 330 to 335, will be used as storage for flood control. Below elevation 330 the reservoir will be used primarily to regulate flow for power generation. It is estimated by the Corps of Engineers that \$104,000 annual flood damage in the valley below

the dam will be averted. Once in operation at power pool level, a minimum of 5,500 cfs regulated flow at Butler Creek gaging station below Augusta will be discharged. It is expected that the year-round natural channel depth between Augusta and Savannah will be increased to 7.3 feet by regulated flow.

Clark Hill Reservoir at top flood control pool (335 feet) will cover an area of 78,000 acres and will extend from the dam to about 37 miles up the Savannah River, with arms extending 29 miles up Little River, Georgia, and 17 miles up Little River, South Carolina. The shore line will be about 1,200 miles long and the reservoir will contain 2,900,000 acre-feet.

The average head for power will be about 130 feet with a minimum of 111 feet to a maximum of 151 feet. The ultimate power capacity of Clark Hill Dam will be 260,000 kilowatts.

#### Proposed Dams and Reservoirs

In addition to Clark Hill Dam and other existing works on the Savannah River and its tributaries, the Corps of Engineers has selected 10 future dam sites. None of these sites are downstream from Augusta. Hartwell Dam is authorized but no construction funds have been appropriated. Its site is on the Savannah River about eight miles below the junction of the Seneca and Tugaloo Rivers. Other selected sites are as follows: Newry-Old Pickens Site on Little River; War Woman, Sand Bottom, Rogues Ford and Camp Creek Sites on the Tugaloo River; Middleton Shoals and Goat Island Sites on the Savannah River between the Tugaloo and Broad Rivers; Tallow Hill and Anthony Shoals Sites on the Broad River.

### Diversion of Savannah River Water

The city of Augusta Diversion Dam on the Savannah River is located a short distance downstream from the Stevens Creek Dam. The flow of Savannah River water to the Augusta Power Canal is regulated to the canal by gates at the head of the canal. The city of Augusta uses the canal as its source of water supply. Large textile mills use the canal as a source of hydroelectric power. Water used for such power generation is returned to the Savannah River via Sibley Mill gates, King Mill gates and Hawks Gully gates. The remaining water in the canal flows into Beaver Dam Ditch which carries most of Augusta's sewage and passes through Phinizy Swamp to join Butler Creek about 300 yards upstream from the confluence of Butler Creek with the Savannah River.

### Navigation

The New Savannah Bluff Lock and Dam was completed April 1, 1937. It controls the depth of the Savannah River for 12 miles below Augusta. During the year 1949 the depth of the Savannah River was maintained at 4.0 feet or more for 95% of the time.

The operation of Clark Hill Reservoir should provide a year around depth of 7.2 feet in the Savannah River channel below Augusta.

Congress has authorized, but has not appropriated, funds for a 9 foot channel 90 feet wide from Augusta to Savannah Harbor. It is possible to establish a channel depth of 12 feet through the construction of two lock and dam systems below Augusta.

### Water Supplies

Augusta, Georgia, uses Savannah River water as its total water



source while Savannah, Georgia, depends only partly on Savannah River water, with deep wells supplying most of its domestic needs.

Savannah's filtration plant, located near Port Wentworth, receives surface water from an intake on Abercorn Creek (Figure 2). The water in Abercorn Creek is largely Savannah River water with some swamp drainage. Most of the water from this plant is used for industrial purposes, though some 8 MGD is used for domestic purposes. For the first eight months of 1952 the water treated daily in this plant averaged 19.79 MGD.

Savannah obtains about 14 MGD of water from deep wells. There are indications that the artesian supply for the city of Savannah is being depleted. <sup>\*</sup>/ The charging area for these wells is in the area around Augusta.

Augusta is the only other city in the Savannah River Basin that uses Savannah River water. Its water supply intake is located in the Augusta Power Canal. Augusta's water filtration plant has seven 3 MGD filters. The maximum raw water pumping capacity is 19.75 MGD.

The other cities and towns in the basin use tributary streams or ground water as their source of water supply.

#### Sources of Pollution

##### Domestic Sewage

The estimated population connected to sewers in the Savannah River Basin is 306,000. Of this amount 150,000, or nearly half is accounted for by the two largest cities in the basin, Savannah and Augusta. Aiken,

<sup>\*</sup>/ U. S. Geol. Survey Water-Supply Paper 987, p. 135, 1945.



South Carolina, discharges treated (secondary) wastes to Horse Creek and thence to the Savannah River. The pollution load to the Savannah River of these treated wastes is negligible.


North Augusta, South Carolina, discharges its raw sewage to the Savannah River at about mile S-216. Its present sewered population amounts to only 3,500, which is small compared to the dilution afforded by the river. However, when present sewer estimations are completed by July 1, 1954, its sewered population will be 12,000.

The pollution load from Augusta enters the Savannah River at two different points. The sewage from about 15% of the population, which amounts roughly to 10,000 P.E. (BOD), is discharged to the river near mile S-215. The remainder is discharged to Beaver Dam Ditch which flows through Phinizy Swamp and enters Butler Creek about 300 yards from the junction of Butler Creek with the Savannah River. Phinizy Swamp acts as a treatment plant to the Augusta sewage. It is estimated that at least 40 per cent of the raw sewage is oxidized and the pollution load to the Savannah River at mile S-202.65 is about 37,000 P.E. (BOD).

About 30,000 of Savannah's population (1950 census) of 119,689 receive secondary treatment. The treated sewage effluent is not discharged to the Savannah River, but to the Vernon River in the Intra-Coastal Waterway. The sewage from the remainder of the population is discharged raw to the Savannah River. It is estimated that the population equivalent of the domestic wastes from Savannah and of the industrial wastes from plants above Savannah that are discharged to the Savannah River is at least 300,000.

### Industrial Wastes

Estimates in population equivalent figures of the amount of industrial pollution to the Savannah River are not available so no figure is included for this type of pollution. Some textile mills and finishing plants discharge wastes to tributaries of the Savannah River but their pollution effect on the river is negligible. For example, though the dissolved oxygen in Horse Creek (S.C.) which joins the Savannah River at mile S-213.4, is depleted by textile wastes during warm weather, its effect on the Savannah River is small because of the low flow in Horse Creek. There are no important industrial waste sources in Augusta, Georgia. At Savannah the two main sources of industrial pollution to the Savannah River are the Union Bag and Paper Company and the Southern Paperboard Corporation.



## CHAPTER IV

STREAM FLOWS

Information on stream flows and characteristics are available from many sources. Data used in this report were obtained principally from the U. S. Geological Survey.

## DISCHARGE DATA

Savannah River at Augusta, Georgia

From U.S.G.S. data the average discharge at Augusta, Georgia, for the years 1898-1906, 1927-1931, 1938-48 is 10,860 second-feet. The drainage area at Augusta, including that of Butler Creek, is 7,508 square miles. The maximum discharge of official record was 350,000 second-feet on October 3, 1929. The maximum gage height was 46.3 feet September 27, 1929 at the auxiliary gage on Fifth Street Bridge on U. S. Highway 1 at Augusta. The minimum daily discharge was 1,040 second-feet on October 2, 1927.

The monthly average daily discharges at Augusta from July 1, 1951 to June 30, 1952 are presented in Table 1 and the average maximum and minimum daily discharges for that period are shown in Table 2.

Savannah River at Buatons Ferry Bridge

The U.S.G.S. records available at this gaging station cover the period October 1939 to September 1948. The maximum discharge during this period occurred on August 18, 1940 and amounted to 141,000 second-feet (gage height, 27.0 feet). A minimum daily flow of 2,240 second-feet occurred on October 18, 25, 1941. The average discharge for the water

year, 1947-48, was 15,000 second-feet. The drainage area is 8,650 square miles.

The monthly daily average discharges at Burtons Ferry Bridge on July 1, 1951 through June 30, 1952 are presented in Table 1. The average, maximum and minimum daily discharges during that period are shown in Table 2.

#### Savannah River Near Clyo

The U.S.G.S. gage is located on the Seaboard Air Line Railroad bridge at S-65. The drainage area amounts to 9,850 square miles.

For an 11 year period (October 1937-September 1948) the average discharge was 11,530 second-feet. During the same period the maximum discharge of 128,000 second-feet occurred on August 21, 22, 1940. The minimum daily discharge of 2,830 second-feet occurred October 19, 25-28, 1941.

The monthly average daily discharges at S-65 (near Clyo) from July 1, 1951 through June 30, 1952 are presented in Table 1. The average maximum and minimum daily discharges for that period are shown in Table 2.

#### Augusta Canal near Augusta, Georgia

The average discharge (U.S.G.S. data) for 17 years (1931-48) was 2,636 second-feet. The maximum daily discharge occurred on January 2, 28, 1948 and was 4,380 second-feet. There was no flow in the canal from April 8 to May 10, 1936.

#### Brier Creek at Millhaven, Georgia

The drainage area is 656 square miles. For an 11 year period (April 1937-September 1948) the average discharge for that period was 673

second-feet. A maximum discharge of 25,400 second-feet occurred on August 16, 1940. A minimum discharge of 110 second-feet was recorded on June 30, July 2, 1945.



TABLE 1 - DAILY AVERAGE SAVANNAH RIVER DISCHARGE,  
IN CFS, FOR THE PERIOD  
JULY 1, 1951 TO JUNE 30, 1952  
BASED ON PROVISIONAL DATA FROM THE U.S.G.S.

| Month        | Augusta<br>(S-202.65) | Burtons Ferry Bridge<br>(S-129.25) | Near Clyo<br>(S-65) |
|--------------|-----------------------|------------------------------------|---------------------|
| July 1951    | 4,570                 | 5,230                              | 5,740               |
| August       | 3,470                 | 4,150                              | 4,790               |
| September    | 3,390                 | 3,860                              | 4,210               |
| October      | 2,730                 | 3,600                              | 4,210               |
| November     | 4,200                 | 5,100                              | 5,870               |
| December     | 8,660                 | 8,630                              | 8,740               |
| January 1952 | 8,650                 | 9,920                              | 11,590              |
| February     | 5,840                 | 7,310                              | 8,910               |
| March        | 29,090                | 28,640                             | 27,930              |
| April        | 21,810                | 26,650                             | 31,150              |
| May          | 6,780                 | 8,170                              | 10,040              |
| June 1952    | 4,340                 | 4,680                              | 5,680               |

TABLE 2 - AVERAGE, MAXIMUM AND MINIMUM DAILY DISCHARGES  
IN CFS, FOR THE PERIOD  
JULY 1, 1951 TO JUNE 30, 1952

|         | Augusta<br>(S-202.65) | Burtons Ferry Bridge<br>(S-129.25) | Near Clyo<br>(S-65) |
|---------|-----------------------|------------------------------------|---------------------|
| Average | 8,640                 | 9,670                              | 10,740              |
| Maximum | 38,600                | 37,100                             | 41,300              |
| Minimum | 1,710                 | 2,120                              | 2,860               |



## CHAPTER V

CHEMICAL, BACTERIOLOGICAL AND PHYSICAL STUDIES

The routine laboratory tests included dissolved oxygen, biochemical oxygen demand, oxygen consumed, pH, turbidity, alkalinity, color and chloride. The routine bacteriological test performed was the coliform MPN (most probable number) per 100 ml. The above data are tabulated in Appendix "A."

The mineral analysis of Savannah River water samples include the following determinations: suspended solids (total and ash), silica, sulfate, magnesium, sodium and potassium, total dissolved solids, iron, fluoride, nitrate, manganese, orthophosphate, total phosphate, calcium, bicarbonate, chloride, iron and aluminum oxides, and aluminum.

Chemical and bacteriological and physical determinations were for the most part performed in accordance with Standard Methods for the Examination of Water and Sewage, Ninth Edition, 1946, of the American Public Health Association. Minor modifications of the standard procedures and new methods used were developed at the Robert A. Taft Sanitary Engineering Center, Cincinnati, Ohio. These were described in the "Interim Report on Savannah River Studies, July 1951 - March 1952." Iron and manganese determinations are presented in Appendix "F."

Especially pertinent chemical, bacteriological, and physical data are discussed below as such relate to water quality of the section of the Savannah River that was surveyed. Also, special studies that relate specifically to the objectives of this survey are included here.

## WATER QUALITY

### Dissolved Oxygen

In general, the mean data curve shows that the Savannah River from above Augusta, Georgia, to Savannah, Georgia, is good. Below Savannah the dissolved oxygen conditions deteriorate rapidly, as shown in Figure 3.

The minimum dissolved oxygen values indicate the most critical conditions encountered. Such conditions usually occur during warm summer months at times of low flows. Figure 3 shows that the Savannah River was in good condition from above Augusta to Savannah but just below Savannah at mile 8-11 there were times when the Savannah River was completely depleted of oxygen. Such a depletion of oxygen causes what is sometimes termed an "oxygen block" in the river. If an "oxygen block" occurs when fish are migrating downstream to the ocean or from the ocean upstream, large numbers of fish may die in their efforts to continue their journey through the "oxygen block". Fish kills have been reported from the lower river in very recent years but their nature or extent could not be ascertained.

### Biochemical Oxygen Demand

In general, the mean and maximum BOD values, as shown in Figure 4, are quite low. There is an increase in BOD below Savannah, as is to be expected below a point of pollution. It is of interest to note that Augusta and North Augusta do not appreciably increase the mean BOD values. The maximum values are slightly over 2 ppm and indicate there is some pollution but not enough to significantly deplete the oxygen resources.

Individual BOD values are presented in Appendix "A".

#### Coliform MPN

Coliform organisms in large numbers are indicative of fecal pollution. However, these may also be found in large numbers in surface runoff from cities and agricultural areas.

A summary of the median, maximum and minimum values for the coliform MPN per 100 ml. test data are presented as a bar diagram in Figure 5. Samples for the coliform test were usually taken at third or quarter points across the stream at each sample station. Upon summarization there was not a sufficient difference at points across the stream to warrant plotting on a stream width as well as stream length basis. Often there was an appreciable difference in the individual samples across the stream.

Usually the median coliform values, as shown in Figure 5, are in excess of 40,000 MPN per 100 ml. for 50 miles of river below Augusta.

The median coliform values of the Savannah River above Augusta are usually less than 1,000 MPN per 100 ml. The Savannah River above Augusta is a suitable source of surface water for Augusta, from the bacteriological standpoint.

#### Minerals

The Savannah River water is mostly in contact with relatively insoluble silicate minerals and is therefore soft and very low in mineral content except for the estuarine portion which is mixed with sea water.

The results from the mineral determinations are presented in tabular form in Appendix "B".

Excluding samples from Station S-11 which are partially sea water, only one analysis out of 23 on the main stem of the river showed a total dissolved solids content greater than 80 ppm. Analyses of tributary waters sometimes exceeded 120 ppm total dissolved solids.

Excluding samples from S-11, of 33 analyses the only mineral constituent exceeding 10 ppm was silica except one result for sulfate and one for chloride. Silica seldom exceeded 25 ppm and was generally less than 20 ppm. The Savannah River contained less than 0.2 ppm fluoride. The manganese content was seldom over 0.03 ppm and frequently less than 0.01 ppm before water was impounded in Clark Hill Reservoir. After water impoundment in Clark Hill Reservoir, a manganese content of 0.6 ppm was found in the river water just below Clark Hill Dam (Page 9 of Appendix "C").

The Savannah River and the tributaries through the Savannah River Project Plant area are low in nitrate and total phosphate. Nitrate values seldom exceed 2 ppm and were usually less than 1 ppm. The results for total phosphate were less than 0.7 ppm and few exceeded 0.5 ppm.

Excluding the samples from S-11, the results from magnesium determinations of main stem and tributary waters were less than 2 ppm. Results for calcium were less than 6 ppm in the main stem. Determinations on water samples from Steel Creek, South Carolina, showed slightly more calcium but not more than 9 ppm. The waters in Lower Three Runs, South Carolina, contained as much as 17 ppm calcium in one of two determinations. Sodium and potassium together were generally less than 10 ppm.

Bicarbonate is the principal acid radical in Savannah River water.




FIG. 3




FIG. 4




FIG. 5



The bicarbonate content of the Savannah River and tributary waters is variable. It ranged from 3.5 ppm to 22 ppm.

### Turbidity

In summarizing the turbidity data at each sampling station on the Savannah River it was found that the mean value varied considerably from station to station. The minimum mean value was 62 ppm and the maximum mean value was 115. Single turbidity values ranged from a minimum of 8 ppm to a maximum of 440 ppm.

When one looks at the Savannah River from its shore, the river has a red muddy appearance and gives the illusion of high turbidity. This red muddy appearance of the stream is caused by colloidal red clay. The water on examination in the laboratory is not nearly as turbid as the stream appearance suggests since the colloid affects "apparent color" rather than turbidity.

### Color

There are two types of "color" in water: "true color" defined as the color due to dissolved substances only, and "apparent color" defined as the color contributed by the suspended material as well as by substances in solution.

The color determinations performed in the Savannah River Studies laboratory were for "true color."

Mean values of color were calculated from data for each station. These mean values ranged from 18 to 132 ppm. Single values from all the stations ranged from 10 to 240 ppm. Most of the "true color" in the Savannah River water is caused by organic material from decomposed plant

residues, since large swampy areas border both sides of the stretch of the Savannah River between Augusta and the Atlantic Ocean. The absence of swamps above Augusta accounts for the lower color values in this reach of river.

The "apparent color" of the water is high because of the red clay colloid which is difficult to remove. The colloid could not be removed from the water by filtration through filter paper but it was removed satisfactorily by a centrifuge.

#### SPECIAL STUDIES

##### Dissolved Oxygen and Salinity Studies in the Savannah River Estuary

On March 28, 1952, when the estimated flow in the river at Savannah was 37,300 cfs, dissolved oxygen and salinity were determined at 2 mile intervals from mile 23 to mile 2 during high tide, between mean and high tide, during mean and low tide. Alkalinity was determined on samples taken at high and low tides. On bottom samples taken 3 feet from the bottom at high tide the critical point for the dissolved oxygen profile was at mile 5 as shown in Figure 7. The D.O. value at this point was 4.7 ppm. For a sample taken 3 feet from the surface at the same time, the curve was still descending at mile 2. The D.O. value at this point was 5 ppm. Samples for chloride and alkalinity were taken at the same time and at the same points as for dissolved oxygen. At high tide the surface samples (3 feet from surface) and bottom samples (3 feet from bottom) have alkalinity and salinity values that parallel from mile 23 to mile 8 as shown in Figure 8. The bottom samples increased very rapidly in alkalinity and salinity from mile 5 to mile 2, while the increase for the surface samples was small.

Between mean and high tide on March 28, 1952 the critical point of the oxygen sag curves for bottom and surface samples shown in Figure 9 were at mile 6 and mile 5, respectively. Compared with the oxygen sag curves at high tide, the critical points had moved upstream 3 miles.

As shown in Figure 10, the salinity in terms of per cent sea water increases rapidly from mile 11 downstream as determined from samples taken between mean and high tide.

During mean tide on March 28, 1952 the critical point of the oxygen sag curve for the bottom samples shown in Figure 11 was at mile 11. The dissolved oxygen value at this point was 3 ppm (estimated Savannah River flow 37,300 cfs.) The oxygen sag curve determined from surface samples was still descending at mile 2 so no critical point can be designated in the range, 3-23 to 3-2. The salinity of the bottom samples increased rapidly from mile 6 downstream, as shown in Figure 12, and amounted to 55 per cent sea water at mile 2. The salinity of the surface samples increased more gradually from mile 6 downstream and amounted to 9 per cent sea water at mile 2.

As shown in Figure 13, the dissolved oxygen values for both bottom and surface water decreased downstream from mile 20 at low tide. At mile 2 the dissolved oxygen values were 6.3 and 5.2 ppm for surface and bottom samples respectively.

Referring to Figure 14, alkalinity values for surface and bottom samples were approximately the same for the entire section. In terms of per cent sea water the salinity increase was moderate starting from less than 1 per cent at mile 6 and increasing to 15 per cent at mile 2 for

bottom samples (Figure 14.) At low tide on March 28, 1952 the salinity of surface samples increased from less than 1 per cent at mile 5 to about 2 per cent sea water at mile 2.


On June 23, 1952 when the estimated river flow was 5,400 cfs at Savannah another dissolved oxygen and salinity study was made covering the Savannah River section, S-23 to S-0.





During high tide the critical point of the oxygen sag curve for surface samples occurred at mile 17 as shown in Figure 15, while the critical point at low tide occurred at mile 3. The critical oxygen values at high and low tides for surface samples were 1.4 ppm and 2.6 ppm, respectively. For bottom samples, Figure 16, critical oxygen values of 1.0 ppm at high tide and 1.5 ppm at low tide occurred at mile 17 and 14, respectively. Comparison of the oxygen sag curves for surface and bottom samples at high tide, as shown by Figure 17, showed little difference between them. The oxygen sag curves for surface and bottom samples at low tide, Figure 18, showed similarity from mile 23 to 18 and from mile 8 to 3. There is quite a difference in the stretch, mile 18 to 8, with the critical value of the bottom sample amounting to 1.5 ppm at mile 14 while that for surface samples is 2.6 ppm at mile 3.

As shown in Figure 19 there is a marked difference in salinity between surface and bottom samples from mile 20 downstream. There is also a marked difference in salinity at low and high tides below mile 20. The data discussed above are tabulated in Appendix "D."

#### Study of Sludge Bank in Lower Savannah River Section, S-14 to S-3

During sampling operations in the Lower Savannah River section it









FIGS. 9 & 10

FIGS. 7 & 8









FIGS 11 & 12



FIGS 13 & 14



FIGS. 15 & 16



FIGS. 17 & 18




FIG. 19



was noticed that a sludge bank existed in the main channel. The sludge bank varied in depth from 6 to 11 feet from approximately mile 14 to mile 8 or mile 8 to mile 2 depending on the flow in the river and the time. The sludge bank actually extends upstream to mile 20 but it is usually less than 6 feet thick above mile 14.

The Corps of Engineers, Savannah District, obtained samples of the sludge at several depths in the sludge bank at mile 11 and mile 14.

The main purpose of the sludge study was to determine the amount of organic matter present at various sludge depths so that the effect on the dissolved oxygen in the stream could be appraised. The results of the chemical analyses are presented in Table 3.

TABLE 3 - CHEMICAL ANALYSES OF SLUDGE SAMPLES FROM S-11 AND S-14

| Sample Station                       | S-11   | S-11    | S-11    | S-14   | S-14   |
|--------------------------------------|--------|---------|---------|--------|--------|
| Depth from Surface of Sav. River     | 33'    | 36'     | 37'     | 32'    | 35'    |
| Lab. No.                             | 573    | 574     | 575     | 576    | 577    |
| Specific Gravity                     | 1.007  | 1.087   | 1.108   | 1.042  | 1.081  |
| % Solids                             | 2.044  | 13.79   | 17.91   | 7.557  | 13.05  |
| Total Susp.                          |        |         |         |        |        |
| Solids - ppm <sup>a</sup> /          | 16,450 | 120,000 | 137,000 | 56,530 | 90,240 |
| Total Susp. Ash - ppm <sup>a</sup> / | 13,810 | 102,370 | 116,510 | 47,980 | 55,010 |
| Total Susp.                          |        |         |         |        |        |
| Volatile - ppm <sup>a</sup> /        | 2,640  | 17,630  | 20,490  | 8,550  | 43,200 |
| TSS- Dry Basis %                     | 80.48  | 37.02   | 76.49   | 74.80  | 75.28  |
| TSA- Dry Basis %                     | 67.56  | 74.23   | 55.05   | 63.49  | 42.12  |
| TSV- Dry Basis %                     | 12.92  | 12.78   | 11.44   | 11.31  | 33.10  |
| Chlorides ppm Cl <sub>2</sub> /      | 2,330  | 5,060   | 4,900   | 4,800  | 6,120  |
| Chlorides Dry Basis %                | 13.64  | 3.67    | 2.74    | 6.35   | 4.59   |
| Oxygen Consumed ppm <sup>a</sup> /   | 1,810  | 11,600  | 15,500  | 6,520  | 11,200 |
| Oxygen Consumed Dry Basis %          | 8.85   | 8.3     | 9.20    | 3.62   | 8.57   |

<sup>a</sup> Values corrected for specific gravity

From the results presented in Table 3 it can be seen that the oxygen consumed values on sludge samples increased greatly the deeper the sample was in the sludge bank, however, on a dry basis there was little difference with increase in depth. The reason for this is obvious. The sludge at increasing depths decreases in water content. On a dry basis the volatile matter in the sludge at mile 11 amounted to about 12 per cent for all three depths while at mile 14 an increase of 3 feet in depth in the sludge bank increased the volatile solids content from a value of 11 per cent to one of 33 per cent.

The results of the chemical analysis of the sludge samples indicate that the sludge bank at mile 14 and mile 11 is probably a mixture of Savannah River silt, solids from city of Savannah raw sewage and fiber from pulp and paper industries upstream.

#### Clark Hill Reservoir Studies

The studies relating to Clark Hill Reservoir were originally intended only to determine the build-up of soluble iron and manganese in the lower depths of the reservoir. However, it was decided to determine the temperature and dissolved oxygen profiles, because of their biological significance and as indicators of currents and stratification in the reservoir; and turbidity because of the information it could give regarding the settling characteristic of the suspended load entering the reservoir.

Shortly after January 20, 1952 the pool elevation of Clark Hill Reservoir passed the 270 ft. level and on March 5, 1952 the pool elevation was about 300 ft. Sampling activities were initiated on March 10, 1952. A map of Clark Hill Reservoir at full pool level on which the

sampling locations are located by arrows is shown in Figure 20. A summary of individual analytical results for Clark Hill Reservoir appears in Appendix "C."

Water samples taken on March 10, 1952 at a point one mile above Clark Hill Dam where the total water depth was 98 ft. showed increasing manganese content as the sample depth increased. The values of soluble manganese content at depths of 23 ft. and 73 ft. were 0.002 and 0.012 ppm, respectively, and the soluble iron content was 0.54 and 0.55 ppm, respectively. On May 6, 1952, at the same distance above the dam and where the total water depth was 114 ft., values for manganese content at depths of 64 ft. and 111 ft. were 0.0013 and 0.27 ppm, respectively. The iron content of these same samples taken at 64 ft. and 111 ft. depths was 0.048 and 0.68 ppm, respectively.

Other samples taken at the previously cited station and at other points in the reservoir, as well as below the dam, confirm the conclusion that the build-up in soluble iron and manganese near the bottom is rapid.

A water sample taken from the Savannah River about 1/4 mile downstream from Clark Hill Dam on May 26, 1952 contained 0.60 ppm of manganese and 0.04 ppm of iron. The temperature of the water sample was 27°C, the dissolved oxygen content was 9.2 ppm and turbidity was 8 ppm. Since the solubility of oxygen in water at a temperature of 27°C is 8.1 ppm, the water sample was supersaturated with oxygen. The aeration of the water as it passes over the spillways of the dam will oxidize some of the soluble iron and manganese converting them to insoluble compounds. Such aeration action is desirable.



Water samples were taken at several depths from Clark Hill Reservoir at a point one mile above Clark Hill Dam (S-239.1). The total water depth at this point was 115 feet. Analytical data on the water samples taken at this station are presented in Table 4.

TABLE 4 - ANALYTICAL DATA ON SAMPLES TAKEN MAY 26, 1952 FROM S-239.1


Clark Hill Reservoir

| Depth<br>ft. | Temp.<br>°C. | Manganese<br>ppm | Iron<br>ppm | Diss. Oxygen<br>ppm | Turbidity<br>ppm |
|--------------|--------------|------------------|-------------|---------------------|------------------|
| 3            | 25           | -                | -           | 7.7                 | 1                |
| 13           | 24           | -                | -           | 7.4                 | 1                |
| 38           | 22           | -                | -           | 4.9                 | 2                |
| 63           | 21           | 0.033            | 0.08        | 4.0                 | 4                |
| 88           | 20           | 0.24             | 0.032       | 2.0                 | 4                |
| 113          | 19           | 1.12             | 0.12        | 0                   | 15               |

The determinations of iron and manganese were performed on clarified samples. Out of curiosity, determinations were performed on non-clarified water samples. While there was no change in the manganese value, the value for iron was considerably greater (over 1.0 ppm) in the non-clarified sample.

The relationship of dissolved oxygen to depth covering a four month period at Stations S-239.1 is shown in Figures 21, 22, 23, and 24. Stratification is quite obvious beginning on May 1 through July 28 with the exception of May 21.


Turbidity values of samples taken at Station S-239.1, one mile above Clark Hill Dam, on March 31 and May 6, 1952, are shown for comparison in Table 5.






CLARK HILL RESERVOIR  
LOCATION MAP

FIG 20

Preceding page blank




- 63 -




D.O. (p.p.m.)

DISSOLVED OXYGEN-DEPTH RELATION (1952)


CLARK HILL RESERVOIR



- 65 -



52 G1F  
SAVANNAH RIVER STUDIES  
JAN. 22, 1951 TO JAN. 21, 1952



# COMPARISON OF AIR & WATER TEMPERATURES

AUGUSTA CANAL  
FEBRUARY 4, 1951 TO FEBRUARY 3, 1952

FIG 26




FIG. 27



TABLE 5 - TURBIDITY VALUES ON SAMPLES TAKEN AT S-239.1  
ON MARCH 31 AND MAY 6, 1952

| March 31, 1952       |                  | May 6, 1952          |                  |
|----------------------|------------------|----------------------|------------------|
| Depth<br>Ft.         | Turbidity<br>ppm | Depth<br>Ft.         | Turbidity<br>ppm |
| 5                    | 140              | 3                    | 1                |
| 25                   | 135              | 14                   | 2                |
| 50                   | 105              | 39                   | 3                |
| 75                   | 175              | 54                   | 8                |
| 100                  | 98               | 89                   | 4                |
| 110 (Total<br>Depth) | -                | 114 (Total<br>Depth) | -                |

Table 5 shows that in 5 weeks time the suspended load has almost settled completely.

On June 13, 1952 temperature determinations were made at various depths in Clark Hill Reservoir at S-239.1 with an electrical resistance thermometer (Foxboro Thermometer.) The temperature decreased gradually by about 0.2 to 0.3 degrees Centigrade per each 5 foot increase in depth from 0 to 105 ft. except for a decrease of 4.5°C. for the 5 ft. interval from the 20 to 25 ft. depth. The temperature at the surface was 29.0°C. and at a depth of 105 ft. it was 16.4°C.

On July 28, 1952 the temperature profile was again determined for 10 ft. intervals in depth from 0 to 110 ft. at S-239.1. No large change in temperature decrease occurred for any 10 ft. interval in this range. The water temperature at the surface was 31.7°C. and that at a depth of 110 ft. was 20.0°C.

#### SAVANNAH RIVER WATER TEMPERATURE IN AUGUSTA CANAL AT AUGUSTA, GEORGIA

Various temperature groupings are used in summarizing the temperature data obtained from the Augusta Water Department, Augusta, Georgia.

The air and water temperature as obtained from the Water Department consisted of 6 readings each, taken at 4 hour intervals during a 24 hour period. The daily temperatures used in the histogram and plots of air and water temperatures were an average of 6-4 hour readings.

Distribution in groups of 4 degrees Fahrenheit of the mean daily water temperatures is shown in the histogram in Figure 27. For almost 20 per cent of the days or more than 60 days of the year's period, February 4, 1951 to February 3, 1952, the water temperature at the Augusta Water Intake was in the 82° to 86° F. group. For less than 10 days the water temperature was in the 62° F. to 66° F. group. Of interest is the 38° to 42° F. temperature group in which there were only 9 days.

Savannah River water temperatures as might be expected of a relatively shallow stream, follow rather closely a running average of the mean daily air temperatures of the preceding seven days. Plots of the mean daily water temperature, running average of the preceding seven days mean air temperatures and the running average of the preceding seven days mean water temperature are shown in Figure 26. The water temperatures taken in the Augusta Canal by the Water Department should be practically the same as that in the Savannah River at Augusta since the point in the Augusta Canal at which the temperatures were read are only a short distance from where the Savannah River water enters the Augusta Canal.

Plots of the daily flow of the Savannah River at Augusta, Georgia, and the daily-average water temperature of the Augusta Canal are shown in Figure 25 for comparison purposes. Though it appeared that there is some negative correlation between flow and water temperature, flow does not

have nearly the effect of air temperature. The effect on water temperature by flow is transmitted through depth changes. An increase in depth diminishes the effect of air temperature. A decrease in depth causing increased turbulence in streams with rough bottoms will increase the effect of air temperature.

## CHAPTER VI

BIOLOGICAL STUDIES

In fulfilling the objectives set forth in the introduction, investigations were undertaken to obtain information on the biota present in various habitats in the Savannah River. A comparison of the results of future biological studies with the background information presented here should aid in detecting and evaluating changes that might occur in the river. Limitations of time and personnel prevented these studies from being all-inclusive. With the exception of the work on plankton, most attention was given macroscopic forms which were most readily collected. Those groups of organisms which appeared to be most important in the biology of the river or of most interest to man were given special attention. Nearly all of the biological work was confined to the strictly fresh-water portion of the river. The observations made on the estuarine portion of the river are treated separately at the end of the biological section.

FRESH-WATER ENVIRONMENTAL ASSOCIATIONS

The groups of organisms associated with the various habitats represented in the fresh-water sections of the Savannah River encompassed by this study may be classified and briefly defined as follows:

1. Plankton - organisms floating or drifting in the water which are not dependent upon a fixed substrate for their existence and organisms which may be normally attached but which have

been swept from their substrate and mixed with the true plankton. The plankton consists largely of the free-floating algae and of a variety of small invertebrates.

2. Benthos - organisms which are more or less dependent upon a solid substrate which is usually fixed but which may, on occasion, be shifting, drifting or floating. This group consists mainly of the insects, oligochaeta, mollusks and certain other invertebrates, and of attached algae. Many benthic organisms may be further subclassified on the basis of the type of substrate which they occupy. These organisms also differ in their manner of association with the substrate; some attach to it, others burrow into it, and others simply move about its surface.

A number of distinct types of habitats for benthonic organisms were present in the river. Separate associations of organisms inhabited different types of substrates; among which were:

- |                             |                               |
|-----------------------------|-------------------------------|
| a. masses of attached algae | e. fixed organic debris (i.e. |
| b. compact clay             | brush, trees, roots, etc.)    |
| c. sand                     | f. rock and gravel            |
| d. mud                      |                               |

3. Nekton - swimming organisms which are able to move about freely and which can maintain themselves against the current. The most important organisms in this group are the fish. Some large, free-swimming crustacea also belong in this assemblage.

Each of these groups of organisms is dealt with separately in the following section on presentation of results. Details of methodology are

also included here since these varied from group to group.

### Plankton

Preliminary collections of plankton taken from the Savannah River in 1951 showed that the plankton population of the river was quite low. Concentrates of river water made in 1951 contained so much silt and so few organisms that quantitative analysis of them were not considered feasible. No extensive sampling for plankton was conducted prior to April 1952. At that time, plankton net tows made in Clark Hill Reservoir showed a sizeable zooplankton population. A sampling program was initiated to determine the fate of the Clark Hill Reservoir plankton populations in the Savannah River downstream from the reservoir.

During the period April 11-14, 1952 quantitative net plankton samples were taken at various depths from two stations (S-239.1 and S-238.6), in the reservoir and from the Savannah River at two stations (S-237.8 and S-219.2), below the reservoir. (Table 6). During the period May 1-5, 1952 quantitative net plankton and nanoplankton <sup>1/</sup> samples were taken at various depths from one station (S-239.1) in the reservoir, and from ten stations in the river below the reservoir, S-237.8, S-219.2, S-219, S-218.4, S-203.1, S-202.9, S-201.65, S-147.6, S-146.4, and S-145.5. (Table 8).

The net plankton samples collected in April 1952 showed a fairly abundant zooplankton population in Clark Hill Reservoir, consisting

<sup>1/</sup> Nanoplankton consists of those organisms small enough to pass through a standard plankton net with #20 silk bolting cloth.



largely of microcrustacea and rotifers. (Table 6). The greatest concentrations of these organisms were in the surface waters, both in the main body of the reservoir and in the arm of the reservoir that was sampled. The samples taken from the Savannah River below the reservoir at this time showed the zooplankton in the river to be much less abundant than that in the reservoir. The water then being discharged to the river was drawn largely from the lower levels of the reservoir, where the plankton was relatively low.

In contrast to the fairly high zooplankton in the reservoir, the net phytoplankton was extremely low; the total numbers of these organisms were never as much as 1,000 organisms per liter. Net plankton and nanoplankton samples were collected simultaneously, in order that the total plankton could be evaluated.

The series of net plankton samples taken in May showed an apparent decrease from April in the total numbers of zooplankton present in Clark Hill Reservoir, but a sizeable population still remained. (Table 7). The zooplankton population in the river downstream from the reservoir remained low, the zooplankton varying from 18 to 55 per liter at different stations.

The net phytoplankton populations in the reservoir remained small in May. In only one instance was the density as great as 1,000 organisms per liter. This paucity of net phytoplankton extended downstream from the reservoir with the only significant variation being noted in the oxbow at mile 146.4, where the net phytoplankton count was more than seven times the number of net phytoplankton organisms found anywhere else in the

TABLE 6 -- CLARK HILL RESERVOIR AND SAVANNAH RIVER NET PLANKTON ORGANISMS PER LITER  
APRIL 1952

|                         |                      |     |     |     |                |      |         |    |     |     |    |
|-------------------------|----------------------|-----|-----|-----|----------------|------|---------|----|-----|-----|----|
| Date :                  | 4/11/52              |     |     |     | 4/14/52        |      |         |    |     |     |    |
| Location :              | CLARK HILL RESERVOIR |     |     |     | SAVANNAH RIVER |      |         |    |     |     |    |
| River Mile:             | S-239.1              |     |     |     | S-239.8*       |      |         |    |     |     |    |
| Depth :                 | Surface              | 3'  | 25' | 50' | 75'            | 100' | Surface | 3' | 25' | 40' | :  |
| MYXOPHYCEAE             |                      |     |     |     |                |      |         |    |     |     |    |
| Blue-Green unicells     |                      |     | 2   | 3   |                |      |         |    |     |     |    |
| TOTAL                   |                      |     | 2   | 3   |                |      |         |    |     |     |    |
| BACILLARIOPHYCEAE       |                      |     |     |     |                |      |         |    |     |     |    |
| Melosira                |                      |     |     |     |                | 3    |         |    |     |     | 13 |
| Others                  |                      |     |     |     |                | 3    |         |    |     |     | 2  |
| TOTAL                   |                      |     |     |     |                |      |         |    |     |     | 15 |
| CHRYSOPHYCEAE           |                      |     |     |     |                |      |         |    |     |     |    |
| Mallomonas              |                      |     | 2   | 6   | 6              | 31   |         | 3  |     |     |    |
| TOTAL                   |                      |     | 2   | 6   | 6              | 31   |         | 3  |     |     |    |
| CHLOROPHYCEAE           |                      |     |     |     |                |      |         |    |     |     |    |
| Eudorina                |                      | 77  | 17  | 9   | 6              |      | 6       | 9  | 2   |     | 4  |
| Others                  |                      | 41  | 17  | 12  | 18             |      | 18      | 7  | 7   |     | 4  |
| TOTAL                   |                      | 118 | 34  | 21  | 24             |      | 24      | 9  | 9   |     |    |
| EUCIOPHYCEAE            |                      |     |     |     |                |      |         |    |     |     |    |
| Trachelomonas           |                      | 22  | 17  | 7   | 3              | 6    |         |    | 19  |     | 4  |
| Others                  |                      |     |     |     |                | 12   |         |    |     |     |    |
| TOTAL                   |                      | 22  | 17  | 7   | 3              | 18   |         |    | 19  |     | 4  |
| DINOPHYCEAE             |                      |     |     |     |                |      |         |    |     |     |    |
| Glenodinium             |                      | 6   | 19  | 15  | 5              |      |         |    | 4   |     |    |
| Others                  |                      | 18  |     |     |                |      |         |    |     |     |    |
| TOTAL                   |                      | 24  | 19  | 15  | 5              |      |         |    | 4   |     |    |
| TOTAL NET PHYTOPLANKTON |                      | 164 | 74  | 72  | 36             | 76   |         | 6  | 12  | 32  | 25 |

TABLE 2 (Continued)

|                   |     |     |    |     |     |     |     |     |    |    |
|-------------------|-----|-----|----|-----|-----|-----|-----|-----|----|----|
| <b>PROTOZOA</b>   |     |     |    |     |     |     |     |     |    |    |
| <b>Actinopoda</b> |     |     |    |     |     |     |     |     |    |    |
| Diffugia          | 25  | 7   | 17 | 22  | 2   | 2   | 13  | 4   | 11 | 4  |
| Ciliata           | 22  | 37  | 22 | 22  | 30  | 30  | 4   | 2   |    | 2  |
| TOTAL             | 47  | 45  | 25 | 44  | 32  | 32  | 19  | 6   | 11 | 10 |
| <b>NOCTURIA</b>   |     |     |    |     |     |     |     |     |    |    |
| Keratella         | 205 | 102 | 23 | 115 | 149 | 149 | 149 | 149 | 11 | 4  |
| Polyarthra        | 55  | 53  | 4  | 3   | 5   | 5   | 34  | 2   |    | 2  |
| Other             | 55  | 3   | 2  | 34  | 5   | 5   | 223 | 6   |    | 4  |
| TOTAL             | 345 | 158 | 32 | 124 | 155 | 155 | 223 | 19  | 11 | 10 |
| <b>NEMATODA</b>   |     |     |    |     |     |     |     |     |    |    |
| TOTAL             |     |     |    | 3   | 3   |     |     |     |    |    |
| <b>CRUSTACEA</b>  |     |     |    |     |     |     |     |     |    |    |
| Bosmina           | 12  | 9   |    |     | 6   | 6   | 9   | 2   |    | 2  |
| Daphnia           | 12  |     |    |     | 3   | 3   | 3   |     |    | 4  |
| Diaptomus         | 18  | 59  | 2  | 3   | 59  | 59  | 40  | 2   | 22 | 4  |
| Diaptomus         | 100 | 158 | 2  |     |     |     | 3   | 6   | 26 | 5  |
| Diaptomus         | 19  | 3   |    |     |     |     | 112 | 9   |    | 12 |
| Diaptomus         | 298 | 254 | 13 | 12  | 205 | 322 | 167 | 10  |    |    |
| Diaptomus         | 445 | 403 | 17 | 27  | 477 | 477 | 390 | 16  | 37 | 54 |
| TOTAL             | 821 | 641 | 75 | 198 | 877 | 877 | 64  |     |    |    |

\*Small Arm of the Reservoir. Estimated to be at S-238.8

TABLE 7 -- CLARK HILL RESERVOIR AND SAVANNAH RIVER NET PLANKTON ORGANISMS PER LITER  
MAY 1952

| Date :                   |     | 5/1/52                                                             |     | 5/2/52                                                           |      | 5/5/52                                |      |
|--------------------------|-----|--------------------------------------------------------------------|-----|------------------------------------------------------------------|------|---------------------------------------|------|
| Location :               |     | CLARK HILL RESERVOIR                                               |     | SAVANNAH RIVER                                                   |      |                                       |      |
| River Mi.:               |     | S-239.1                                                            |     | S-237.8:219.2:219.2:218.4:203.1:202.9:201.65:S-147.0:146.0:145.5 |      |                                       |      |
| Depth :                  |     | Surface 3' 25' 50' 70' 100' :Surface 3' 3' 3' 3' 3' 3' 3' 3' 3' 3' |     | Surface 3' 3' 3' 3' 3' 3' 3' 3' 3' 3'                            |      | Surface 3' 3' 3' 3' 3' 3' 3' 3' 3' 3' |      |
| <b>MYXOPHYCEAE</b>       |     |                                                                    |     |                                                                  |      |                                       |      |
| Chroococcus              | 14  |                                                                    |     |                                                                  |      | 1                                     | 6    |
| Others                   |     |                                                                    |     |                                                                  |      | 1                                     | 6    |
| TOTAL                    | 14  |                                                                    |     |                                                                  |      |                                       |      |
| <b>BACILLARIOPHYCEAE</b> |     |                                                                    |     |                                                                  |      |                                       |      |
| Melosira                 | 18  | 3                                                                  | 4   | 6                                                                | 2    | 3                                     | 17   |
| Others                   | 18  | 6                                                                  | 4   | 82                                                               | 5    | 10                                    | 41   |
| TOTAL                    | 18  | 9                                                                  | 8   | 99                                                               | 8    | 16                                    | 58   |
| <b>CHRYSTOPHYCEAE</b>    |     |                                                                    |     |                                                                  |      |                                       |      |
| Dinobryon                | 19  |                                                                    |     | 54                                                               |      | 4                                     | 15   |
| Mallomonas               | 218 |                                                                    |     |                                                                  |      | 1                                     |      |
| Synura                   | 9   |                                                                    |     | 54                                                               |      | 5                                     | 15   |
| TOTAL                    | 246 |                                                                    |     |                                                                  |      |                                       |      |
| <b>CHLOROPHYCEAE</b>     |     |                                                                    |     |                                                                  |      |                                       |      |
| Eudorina                 | 5   | 5                                                                  | 8   | 2                                                                | 2    | 5                                     | 6    |
| Others                   | 5   | 5                                                                  | 8   | 8                                                                | 4    | 5                                     | 6    |
| TOTAL                    | 10  | 10                                                                 | 16  | 10                                                               | 6    |                                       |      |
| <b>EUGLENOPHYCEAE</b>    |     |                                                                    |     |                                                                  |      |                                       |      |
| Trachelomonas            | 5   | 37                                                                 | 124 | 4                                                                |      | 1                                     | 56   |
| Others                   | 5   | 5                                                                  | 4   | 4                                                                |      | 1                                     | 7    |
| TOTAL                    | 10  | 37                                                                 | 124 | 8                                                                |      | 2                                     | 63   |
| <b>DINOPHYCEAE</b>       |     |                                                                    |     |                                                                  |      |                                       |      |
| Glenodinium              |     |                                                                    | 8   | 983                                                              | 13   | 1                                     | 84   |
| Others                   |     |                                                                    | 8   | 6                                                                | 1    |                                       |      |
| TOTAL                    |     |                                                                    | 8   | 989                                                              | 13   | 1                                     | 84   |
| TOTAL NET PHYTOPLANKTON  | 15  | 346                                                                | 9   | 24                                                               | 1119 | 70                                    | 1258 |
|                          |     |                                                                    |     |                                                                  |      | 37                                    | 160  |

[illegible][illegible]

\* Oz-Bot. 60 5-115.4.

Reproduced from  
best available copy.

stretch of river sampled. This count was still only 1,258 organisms per liter, or slightly more than one organisms per milliliter. This increase in net phytoplankton was not reflected in the sample from the river below the ox-bow (mile 145.5); the net phytoplankton there was only 160 organisms per liter.

Nannoplankton was also extremely rare in the river. (Table 8). Nannophytoplankton for each station was added to the net phytoplankton to attain the total numbers of phytoplankton. The total phytoplankton (Table 9) of the river showed no significant changes over the stretch of river studied, varying from 3,200 to 6,200 organisms per liter at different stations.

The plankton populations in Clark Hill Reservoir did not appear to influence the plankton populations in the river to any considerable extent, during April and May 1952. The sampling errors involved in dealing with such small numbers of organisms would make it impossible to detect any such influence unless it occurred to a marked degree. Some of the organisms comprising the river plankton appeared to be contributed by Clark Hill Reservoir, while others apparently entered the river from other sources such as sloughs, ox-bows, and tributaries draining the flood plain. Diatoms, in particular, appeared to be multiplying in the river, or entering it from some source other than Clark Hill Reservoir. Eleven genera of diatoms were identified in the river plankton, and only five of these were found in Clark Hill Reservoir. (Table 10). The total numbers of diatoms were consistently higher in the nannoplankton counts from the river than they were in the counts from the reservoir. This was also generally true of the diatoms in the net plankton. The limited



data from mile 146.4 indicated that some of this slight increase in diatoms may have originated in ox-bows on the river.

The only major pollution entering the stretch of river sampled for plankton was the sewage from Augusta which is discharged to the river at two points. The sewage from about 15% of the population entered the river at mile 215, and the remainder of the sewage entered the river through Butler Creek at mile 202.65. Plankton samples taken downstream from these points at miles 203.1 and 201.65, respectively, did not show any detectable quantitative changes in the total phytoplankton. Such increases would ordinarily be expected, due to the fertilizing effects of organic pollution; however, the low transparency of the river water was probably a limiting factor in these areas.

It is expected that Clark Hill Reservoir will continue to contribute some organisms to the plankton of the river, but it is difficult to determine to what extent future increases in the plankton of the reservoir will be reflected in the plankton of the river downstream. Any lowering of turbidity in the river, brought about by settling of suspended silt in Clark Hill Reservoir would undoubtedly improve conditions for the survival and reproduction of plankton populations in the river.

### Benthos

#### Benthic Algal Growths and Associated Organisms

The low transparency of the water and the extreme variation of water levels prevalent at the time of the present studies apparently limited the growth of attached algae in the river. Masses of attached algae were observed, however, during the few periods of stable water

|            |                          |   |     |   |                                                                            |   |  |   |        |
|------------|--------------------------|---|-----|---|----------------------------------------------------------------------------|---|--|---|--------|
| Date :     | 5/1/52                   | : |     | : | 5/2/52                                                                     | : |  | : | 5/5/52 |
| Location : | CLARK HILL RESERVOIR     | : |     | : | SAVANNAH RIVER                                                             | : |  | : |        |
| River Mi.: | S-239.1                  | : |     | : | S-237.8; 219.2; 219.2; 218.4; 203.1; 202.9; 201.65; S-147.0; 146.4*; 145.5 | : |  | : |        |
| Depth:     | Surface 3'    25'    50' | : | 75' | : | Surface: 3'    3'    3'    3'    3'    3'    3'    3'    3'                | : |  | : |        |

## MYXOHYCEAE

|              |   |   |   |   |   |
|--------------|---|---|---|---|---|
| Chirococcus  | 5 | 2 | 5 | 2 | 5 |
| Others       | 5 | 2 | 5 | 3 | 5 |
| <u>TOTAL</u> | 5 | 2 | 5 | 5 | 5 |

## BACILLARIOPHYCEAE

|                  | 5  | 2  | 14 | 2  | 9 | 14 | 2 | 11 | 5 | 24 | 45 |
|------------------|----|----|----|----|---|----|---|----|---|----|----|
| <u>Gyrosigma</u> |    |    |    |    |   |    |   |    |   |    |    |
| <u>Melosira</u>  |    |    |    |    |   |    |   |    |   |    |    |
| <u>Navicula</u>  |    | 2  |    |    |   |    |   |    |   |    |    |
| <u>Nitzschia</u> |    |    |    |    |   |    |   |    |   |    |    |
| <u>Synedra</u>   | 5  |    | 2  |    |   |    |   |    |   |    |    |
| <u>Others</u>    |    |    | 14 | 2  | 9 | 14 | 2 | 11 | 5 | 24 | 45 |
| <u>TOTAL</u>     | 14 | 25 | 2  | 16 | 9 | 2  | 2 | 2  | 2 | 2  | 2  |

CHLOROPHYCEAE

|                     |   |   |   |   |   |   |   |
|---------------------|---|---|---|---|---|---|---|
| Chlamydomonas group | 5 | 7 | 5 | 9 | 2 | 6 | 2 |
| Others              | 2 | 2 | 2 | 2 | 2 | 1 | 2 |
| TOTAL               | 7 | 7 | 7 | 9 | 2 | 7 | 2 |

## EUGLENOPHYCEAE

|               |    |    |   |   |   |   |   |   |
|---------------|----|----|---|---|---|---|---|---|
| Euglena       | 7  | 9  | 2 | 5 | 5 | 2 | 5 | 5 |
| Trachelomonas | 5  | 2  |   |   |   | 2 |   | 2 |
| TOTAL         | 12 | 11 | 2 | 5 | 5 | 2 | 5 | 7 |

ДИ. ПУЧКА:

|             |    |    |   |   |   |
|-------------|----|----|---|---|---|
| Glenodinium | 34 | 13 | 2 | 5 | 2 |
| Others      |    |    |   |   |   |
| TOTAL       | 34 | 18 | 2 | 5 | 2 |
| TOTAL       |    |    |   |   |   |

|                 |    |                    |    |    |    |    |    |                            |
|-----------------|----|--------------------|----|----|----|----|----|----------------------------|
|                 | 67 | 56                 | 11 | 39 | 18 | 11 | 39 | 38                         |
| EALHOPIPLANKTON | *  | Cz-Bow at 5-146 h. |    |    |    |    |    | Wannoplankton sample lost. |

\*

TABLE No. 9

## CLARK HILL RESERVOIR AND SAVANNAH RIVER

## TOTAL PHYTOPLANKTON ORGANISMS PER LITER

| <u>DATE</u> | <u>LOCATION</u>            | <u>RIVER MILE</u> | <u>DEPTH</u> | <u>TOTAL<br/>NET<br/>PHYTO-<br/>PLANKTON</u> | <u>TOTAL<br/>NANNO-<br/>PHYTO-<br/>PLANKTON</u> | <u>TOTAL<br/>PHYTO-<br/>PLANKTON</u> |
|-------------|----------------------------|-------------------|--------------|----------------------------------------------|-------------------------------------------------|--------------------------------------|
| 5/1/52      | Clark<br>Hill<br>Reservoir | S-239.1           | Surface      | 0                                            | 6700                                            | 6700                                 |
|             |                            |                   | 3'           | 400                                          | 5600                                            | 6000                                 |
|             |                            |                   | 25'          | 0                                            | 1100                                            | 1100                                 |
|             |                            |                   | 50'          | 0                                            | 3000                                            | 3000                                 |
|             |                            |                   | 75'          | 1100                                         | 1800                                            | 2900                                 |
| 5/2/52      | Savannah<br>River          | S-237.8           | 100'         | 100                                          | 1100                                            | 1200                                 |
|             |                            |                   |              |                                              |                                                 |                                      |
|             |                            | S-237.8           | Surface      | 0                                            | 3900                                            | 3900                                 |
|             |                            | S-219.2           | 3'           | 0                                            | 3800                                            | 3800                                 |
|             |                            | S-219             | 3'           | 200                                          | 5200                                            | 5400                                 |
|             |                            | S-218.4           | 3'           | 100                                          | 4500                                            | 4600                                 |
|             |                            | S-203.1           | 3'           | 100                                          | 3100                                            | 3200                                 |
|             |                            | S-202.9           | 3'           | 0                                            | 6200                                            | 6200                                 |
| 5/5/52      |                            | S-201.65          | 3'           | 0                                            | 6000                                            | 6000                                 |
|             |                            | S-147.0           | 3'           | 100                                          | 6000                                            | 6100                                 |
|             |                            | S-146.4*          | 3'           | 1300                                         | --                                              | --                                   |
|             |                            | S-145.5           | 3'           | 200                                          | 4800                                            | 5000                                 |

\* Ox-Bow at S-146.4. Nannoplankton sample lost.

TABLE NO. 10

ORGANISMS FOUND IN CLARK HILL RESERVOIR AND SAVANNAH RIVER PLANKTON  
APRIL-MAY 1952

| ORGANISMS                 | SAVANNAH RIVER |     | CLARK HILL RESERVOIR |     |
|---------------------------|----------------|-----|----------------------|-----|
|                           | APRIL          | MAY | APRIL                | MAY |
| <u>MYXOPHYCEAE</u>        |                |     |                      |     |
| <u>Chroococcus</u>        |                | x   |                      | x   |
| <u>Oscillatoria</u> group |                | x   |                      |     |
| <u>Phormidium</u>         |                | x   |                      |     |
| Undetermined unicells     |                | x   |                      |     |
| <u>BACILLARIOPHYCEAE</u>  |                |     |                      |     |
| <u>Asterionella</u>       |                | x   |                      |     |
| <u>Ceratoneis</u>         |                | x   |                      |     |
| <u>Cyclotella</u>         |                | x   |                      | x   |
| <u>Cymbella</u>           |                | x   |                      |     |
| <u>Fragilaria</u>         |                | x   |                      |     |
| <u>Gomphonema</u>         | x              | x   |                      | x   |
| <u>Gyrosigma</u>          |                | x   |                      |     |
| <u>Melosira</u>           | x              | x   |                      | x   |
| <u>Navicula</u>           |                | x   |                      | x   |
| <u>Nitzschia</u>          |                | x   |                      |     |
| <u>Synedra</u>            |                | x   |                      | x   |
| Undetermined Pennales     |                | x   | x                    | x   |
| <u>CHRYSTOPHYCEAE</u>     |                |     |                      |     |
| <u>Dinobryon</u>          |                | x   |                      |     |
| <u>Malomonas</u>          |                | x   | x                    | x   |
| <u>Synura</u>             |                |     |                      | x   |
| <u>CHLOROPHYCEAE</u>      |                |     |                      |     |
| <u>Ankistrodesmus</u>     |                | x   |                      |     |
| <u>Cerasterias</u>        |                | x   | x                    | x   |
| <u>Chlamydomonas</u>      |                | x   |                      |     |
| <u>Closterium</u>         |                | x   |                      |     |
| <u>Coelastrum</u>         |                | x   | x                    |     |
| <u>Cosmarium</u>          |                | x   |                      |     |
| <u>Desmidium</u>          |                | x   |                      |     |
| <u>Eudorina</u>           |                |     | x                    |     |
| <u>Gloeocystis</u>        |                |     | x                    |     |
| <u>Micractinium</u>       |                |     | x                    |     |
| <u>Oedogonium</u>         |                | x   |                      |     |
| <u>Oocystis</u>           |                |     | x                    |     |
| <u>Pandorina</u>          |                |     | x                    |     |
| <u>Pediastrum</u>         |                |     | x                    |     |
| <u>Pleodorina</u>         |                | x   | x                    |     |

TABLE NO.10 (continued)  
 ORGANISMS FOUND IN CLARK HILL RESERVOIR AND SAVANNAH RIVER PLANTATION  
 APRIL-MAY 1952

| ORGANISMS                          | SAVANNAH RIVER |     | CLARK HILL RESERVOIR |     |
|------------------------------------|----------------|-----|----------------------|-----|
|                                    | APRIL          | MAY | APRIL                | MAY |
| CHLOROPHYCEAE (continued)          |                |     |                      |     |
| <u>Scenedesmus</u>                 |                |     |                      | X   |
| <u>Stigeoclonium</u>               |                | X   |                      |     |
| <u>Tetraedron</u>                  | X              |     |                      |     |
| <u>Volvox</u>                      |                | X   | X                    | X   |
| <u>Volvocales cysts</u>            |                | X   |                      | X   |
| EUGLENOPHYCEAE                     |                |     |                      |     |
| <u>Euglena</u>                     |                | X   |                      | X   |
| <u>Trachelomonas</u>               |                | X   | X                    | X   |
| <u>Undetermined Euglenophyceae</u> | X              | X   | X                    | X   |
| DINOPHYCEAE                        |                |     |                      |     |
| <u>Ceratium</u>                    |                |     | X                    |     |
| <u>Glenodinium</u>                 |                | X   | X                    | X   |
| <u>Peridinium</u>                  |                | X   | X                    | X   |
| PROTOZOA                           |                |     |                      |     |
| <u>Diffugia</u>                    |                | X   | X                    | X   |
| <u>Actinopoda</u>                  |                |     | X                    |     |
| <u>Undetermined Ciliata</u>        | X              | X   | X                    | X   |
| NEMATODA                           |                |     |                      |     |
|                                    |                | X   | X                    |     |
| ROTATORIA                          |                |     |                      |     |
| <u>Conochilus</u>                  |                | X   | X                    | X   |
| <u>Keratella</u>                   | X              | X   | X                    | X   |
| <u>Monostyla</u>                   |                | X   | X                    |     |
| <u>Notholca</u>                    |                | X   | X                    |     |
| <u>Polyarthra</u>                  | X              | X   | X                    | X   |
| <u>Undetermined Rotatoria</u>      | X              | X   | X                    | X   |
| <u>Rotifer eggs</u>                |                | X   |                      |     |
| CRUSTACEA                          |                |     |                      |     |
| <u>Bosmina</u>                     |                | X   | X                    | X   |
| <u>Chydorus</u>                    |                |     | X                    | X   |
| <u>Daphnia</u>                     | X              | X   | X                    | X   |
| <u>Cyclops</u>                     | X              | X   | X                    | X   |
| <u>Diaptomus</u>                   |                |     | X                    | X   |
| <u>Nauplii</u>                     | X              | X   | X                    | X   |
| INSECTA                            |                |     |                      |     |
| <u>Tendipedidae</u>                |                | X   |                      |     |

levels. Even on these occasions the vertical zone wherein effective photosynthesis could occur was quite narrow. Substrates such as floating debris, willow branches, and roots, which could rise and fall with the water level keeping the algae near the surface, supported algal growths for longer periods than fixed substrates. Algal growths on fixed substrates were often stranded by receding water or kept below the photosynthetic zone by rising water.

Periods of algal abundance which persisted for intervals of several weeks were observed during the winter, spring and summer of 1952. They were attributed to extended periods of relatively stable water levels. The filamentous green alga, Stigeoclonium lubricum, was the most abundant species observed during the winter and spring of 1952. It was often found growing in long streamers attached to willow branches in swift water. The red alga, Auchmoneia, and the filamentous green algae, Heliothrix and Vaucheria, were frequently found in similar situations. Other genera of algae sometimes found in abundance were the diatoms, Fragilaria, Terpsinoe, and Malacra; the green algae, Stauroneis, Microspora and Tetraspora; and the blue-green alga, Cylindrocapsa. Some of these species preferred quieter waters and were found in flooded swamp areas and back waters rather than in the river proper. Spirogyra became extremely abundant during the summer, when it was often found in quiet areas sheltered by sand bars. Phormidium was observed throughout the year growing in closely adherent mats on submerged sticks and logs and on shallow sand bars. Seasonal changes in the composition of the algae were probably caused by increased water temperatures which created unfavorable conditions



for the species predominating earlier in the year and favored warm water species.

Filamentous algae growing in fast water were generally found in almost pure stands, while the algae growing in quieter water often had many other organisms growing in association with them. For example, protozoans, rotifers, nematodes and oligochaeta were often found abundantly living in association with the algae in stagnant areas. Masses of Stigeoclonium, Vaucheria, and Judoniella growing attached to willow branches in fast water at S-165.4 collected on January 1, 1952 contained sizeable populations of insects. This fauna was largely composed of the immature stages of mayflies (Ephemerella sp.), midges (Spaniotoma sp.) and caddis flies (Leptocella sp.). Other immature insects present were stoneflies (Hydroperla harti), caddis flies (Macronemum carolina, Hydroptila sp. and undetermined Psychomyiidae), midges (Polypedilum sp. and Tanytarsus dissimilis), snipe flies (Atherix variegatus) and beetles of the family Elmidae.

Twelve samples of algal masses were collected from the river at S-171.62; S-170.5; S-169.0; S-165.4; S-155.3; S-152.0, and S-146.4 and from Upper Three Runs on April 18, 1954 for critical examination of the invertebrate fauna associated with them. The water level was high at that time, and an area for some distance upstream from the mouth of Upper Three Runs was inundated by Savannah River water. Samples of algal masses were taken at the mouth of this tributary, S-171.59, UTR-.1, and two miles upstream from the river backwater at S-171.59, UTR-200. A single algal mass was collected from the river on May 12, 1952 and another one on July 23, 1952 at S-155.3 and S-169.0. These algal masses were brought into the

TABLE 11 - SAVANNAH RIVER STUDIES  
ALGAL-MASS ALGAE

8

|                  |                           |                                                                                                          |                                                                                                          |                                                                                                          |                                                                                                          |                                                                                                          |                                                                                                          |
|------------------|---------------------------|----------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------|
| Location(River): | S-171.62                  | S-171.59                                                                                                 | S-171.59                                                                                                 | S-171.59                                                                                                 | S-171.59                                                                                                 | S-171.59                                                                                                 | S-171.59                                                                                                 |
| Mile)            |                           | Ut-2.0                                                                                                   | Ut-0.1                                                                                                   | Ut-0.1                                                                                                   | Ut-2.0                                                                                                   | Ut-0.1                                                                                                   | Ut-0.1                                                                                                   |
| Date (1952)      | 4/18/52                   | 4/18/52                                                                                                  | 4/18/52                                                                                                  | 4/18/52                                                                                                  | 4/18/52                                                                                                  | 4/18/52                                                                                                  | 4/18/52                                                                                                  |
| Predominant      | Stigeoclonium: Melosira   | Vaucheria                                                                                                | Melosira sp, Melosira                                                                                    | Melosira                                                                                                 | Melosira                                                                                                 | Stigeoclonium: Stigeoclonium                                                                             | Stigeoclonium: Stigeoclonium                                                                             |
| Algae            | lubricum                  | polysarum, Stigeoclonium: italicum, Stigeoclonium: lubricum                                              | lubricum                                                                                                 | lubricum                                                                                                 | Oedogonium sp, Fragilaria sp                                                                             | lubricum                                                                                                 | nium                                                                                                     |
|                  | sp.                       |                                                                                                          |                                                                                                          |                                                                                                          | Fragilaria sp                                                                                            |                                                                                                          | lubricum                                                                                                 |
|                  |                           |                                                                                                          |                                                                                                          |                                                                                                          |                                                                                                          |                                                                                                          | Spirogyra                                                                                                |
|                  |                           |                                                                                                          |                                                                                                          |                                                                                                          |                                                                                                          |                                                                                                          | sp,                                                                                                      |
|                  |                           |                                                                                                          |                                                                                                          |                                                                                                          |                                                                                                          |                                                                                                          | Vaucheria                                                                                                |
|                  |                           |                                                                                                          |                                                                                                          |                                                                                                          |                                                                                                          |                                                                                                          | sp.                                                                                                      |
| Other            | Lyngbya sp.               | Asterionella: Navicula sp, Vaucheria sp, Navicula sp, Zygnema sp, Fragilaria                             | Navicula sp, Ulothrix sp, Spirogyra sp, Eunotia sp, Tetraspora sp, sp,                                   | Navicula sp, Ulothrix sp, Spirogyra sp, Eunotia sp, Tetraspora sp, sp,                                   | Navicula sp, Ulothrix sp, Spirogyra sp, Eunotia sp, Tetraspora sp, sp,                                   | Navicula sp, Ulothrix sp, Spirogyra sp, Eunotia sp, Tetraspora sp, sp,                                   | Navicula sp, Ulothrix sp, Spirogyra sp, Eunotia sp, Tetraspora sp, sp,                                   |
| Algae            | Frustulia                 | Spirogyra sp, Fragilaria sp, Synedra sp, Ankistrodes-                                                    | Spirogyra sp, Fragilaria sp, Synedra sp, Ankistrodes-                                                    | Spirogyra sp, Fragilaria sp, Synedra sp, Ankistrodes-                                                    | Spirogyra sp, Fragilaria sp, Synedra sp, Ankistrodes-                                                    | Spirogyra sp, Fragilaria sp, Synedra sp, Ankistrodes-                                                    | Spirogyra sp, Fragilaria sp, Synedra sp, Ankistrodes-                                                    |
| present          | vulgaris,                 | Fragilaria sp, Oedogonium sp, Mus sp, Desmidiun sp, Mougeotia sp, Ulothrix sp, Anabaena sp, Oscillatoria | Fragilaria sp, Oedogonium sp, Mus sp, Desmidiun sp, Mougeotia sp, Ulothrix sp, Anabaena sp, Oscillatoria | Fragilaria sp, Oedogonium sp, Mus sp, Desmidiun sp, Mougeotia sp, Ulothrix sp, Anabaena sp, Oscillatoria | Fragilaria sp, Oedogonium sp, Mus sp, Desmidiun sp, Mougeotia sp, Ulothrix sp, Anabaena sp, Oscillatoria | Fragilaria sp, Oedogonium sp, Mus sp, Desmidiun sp, Mougeotia sp, Ulothrix sp, Anabaena sp, Oscillatoria | Fragilaria sp, Oedogonium sp, Mus sp, Desmidiun sp, Mougeotia sp, Ulothrix sp, Anabaena sp, Oscillatoria |
|                  | Tabellaria                | Oedogonium                                                                                               | Oedogonium                                                                                               | Oedogonium                                                                                               | Oedogonium                                                                                               | Oedogonium                                                                                               | Oedogonium                                                                                               |
|                  | flocculosa, sp,           | Navicula sp, Melosira sp, Lyngbya sp.                                                                    | Navicula sp, Melosira sp, Lyngbya sp.                                                                    | Navicula sp, Melosira sp, Lyngbya sp.                                                                    | Navicula sp, Melosira sp, Lyngbya sp.                                                                    | Navicula sp, Melosira sp, Lyngbya sp.                                                                    | Navicula sp, Melosira sp, Lyngbya sp.                                                                    |
|                  | Navicula sp, Melosira sp, |                                                                                                          |                                                                                                          |                                                                                                          |                                                                                                          |                                                                                                          |                                                                                                          |
|                  | Eunotia sp, Lyngbya sp.   |                                                                                                          |                                                                                                          |                                                                                                          |                                                                                                          |                                                                                                          |                                                                                                          |
|                  | Closterium                |                                                                                                          |                                                                                                          |                                                                                                          |                                                                                                          |                                                                                                          |                                                                                                          |
|                  | sp,                       |                                                                                                          |                                                                                                          |                                                                                                          |                                                                                                          |                                                                                                          |                                                                                                          |
|                  | Oedogonium                |                                                                                                          |                                                                                                          |                                                                                                          |                                                                                                          |                                                                                                          |                                                                                                          |
|                  | sp,                       |                                                                                                          |                                                                                                          |                                                                                                          |                                                                                                          |                                                                                                          |                                                                                                          |
|                  | Tetraspora                |                                                                                                          |                                                                                                          |                                                                                                          |                                                                                                          |                                                                                                          |                                                                                                          |
|                  | sp,                       |                                                                                                          |                                                                                                          |                                                                                                          |                                                                                                          |                                                                                                          |                                                                                                          |
|                  | Anabaena sp,              |                                                                                                          |                                                                                                          |                                                                                                          |                                                                                                          |                                                                                                          |                                                                                                          |
|                  | Oscillatoria              |                                                                                                          |                                                                                                          |                                                                                                          |                                                                                                          |                                                                                                          |                                                                                                          |
|                  | sp,                       |                                                                                                          |                                                                                                          |                                                                                                          |                                                                                                          |                                                                                                          |                                                                                                          |
|                  | Undet.                    |                                                                                                          |                                                                                                          |                                                                                                          |                                                                                                          |                                                                                                          |                                                                                                          |
|                  | Euglenophyta:             |                                                                                                          |                                                                                                          |                                                                                                          |                                                                                                          |                                                                                                          |                                                                                                          |
| Weight of Algal: |                           |                                                                                                          |                                                                                                          |                                                                                                          |                                                                                                          |                                                                                                          |                                                                                                          |
| Mass(grams)      | 13.2                      | 15.8                                                                                                     | 22.9                                                                                                     | 2.7                                                                                                      | 24.5                                                                                                     | 19.                                                                                                      |                                                                                                          |

TABLE II - (Continued)

|                   |            |              |               |                |            |                |              |
|-------------------|------------|--------------|---------------|----------------|------------|----------------|--------------|
| Location (River): | S-170.5    | S-170.2      | S-169.0       | S-165.4        | S-155.3    | S-152.0        | S-146.4      |
| Mile)             |            |              |               |                |            |                |              |
| Date              | 4/18/52    | 4/18/52      | 7/23/52       | 4/18/52        | 5/12/52    | 4/18/52        | 4/18/52      |
| Predominant       | Vauleria   | Stigeoclo-   | Spirogyra sp. | Stigeoclonium  | Oedogonium | Stigeoclonium  | Oedogonium   |
| Algae             | polysperma | nium         |               | lubricum       |            | lubricum, nium |              |
|                   |            | lubricum     |               |                |            | Melosira sp.   | sp.          |
|                   |            |              |               |                |            |                | Fragilaria   |
|                   |            |              |               |                |            |                | laria sp.    |
| Other             |            | Oedogonium   | Schizomeru    | Oedogonium sp. |            | Fragilaria sp. | Stigeoclo-   |
| Algae             |            | sp.          | sp.           | Melosira sp.   |            | Spirogyra      | nium         |
| present           |            | Comphonema   |               |                |            | tenuissima     | lubricum     |
|                   |            | sp.          |               |                |            | Eunotia sp.    | Fragilaria   |
|                   |            | Lanobya sp.  |               |                |            | sp.            |              |
|                   |            | Tetraspora   |               |                |            |                | Eunotia sp.  |
|                   |            | Vauleria sp. |               |                |            |                | Melosira sp. |
|                   |            | Melosira sp. |               |                |            |                | Oscilla-     |
|                   |            | Tatellora    |               |                |            |                | toria sp.    |
|                   |            | sp.          |               |                |            |                |              |
| Height of algal   |            |              |               |                |            |                |              |
| thrust (cm)       | 46.7       | 41.3         | 89.2          |                | 32.7       | 14.8           | 5.0          |

TABLE 12 - SAVANNAH RIVER STUDIES  
ANIMALS ASSOCIATED WITH ALGAL MASSES

|                 |                                 |
|-----------------|---------------------------------|
| Location (River | :S-171.62:171.59:171.59:171.59: |
| Mile)           | : :Ut-2.0:Ut-0.1:Ut-0.1:        |
| Date (1952)     | :4/18/52 :4/18 :4/18 4/18 :     |

## ANNELIDA:OLIGOCHAETA

|                                              |   |    |    |
|----------------------------------------------|---|----|----|
| <u>Nais communis</u> , <u>Nais josinae</u> , |   |    |    |
| Undet. Nais                                  |   | 83 |    |
| <u>Naidium</u> sp.                           |   |    |    |
| <u>Pristina aequiseti</u>                    |   | 46 |    |
| <u>Stylaria fossularis</u> ,                 |   |    |    |
| <u>Stylaria proboscidea</u> ,                |   |    | 12 |
| Undet. Naididae                              | 3 | 4  | 51 |
| Undetermined                                 |   |    |    |

## ARTHROPODA:CRUSTACEA

|                                                    |    |   |  |
|----------------------------------------------------|----|---|--|
| Copepoda: <u>Cyclops</u> sp., <u>Diaptomus</u> sp. |    |   |  |
| Ostracoda:Undetermined                             |    |   |  |
| Cladocera: <u>Daphnia</u> sp.                      |    |   |  |
| <u>Ceriodaphnia</u> sp.                            |    |   |  |
| <u>Eurycerus lamellatus</u> ,                      |    |   |  |
| <u>Croperus harpae</u> , <u>A. angustatus</u> ,    |    |   |  |
| <u>Pleuroxus denticulatus</u>                      | 71 | 1 |  |
| Amphipoda: <u>Crangonyx gracilis</u>               |    |   |  |

## ARTHROPODA:ARACHNIDA

|                          |  |   |
|--------------------------|--|---|
| Hydracarina:Undetermined |  | 1 |
|--------------------------|--|---|

## ARTHROPODA:INSECTA

|                                            |  |   |  |
|--------------------------------------------|--|---|--|
| Flecoptera: <u>Psephenus placida</u>       |  |   |  |
| <u>Isoperla bilineata</u>                  |  | 1 |  |
| Ephemeroptera: <u>Stenonema</u> sp.        |  |   |  |
| <u>Caenis</u> sp.                          |  |   |  |
| <u>Ephemerella</u> spp. <u>invaria</u> gr. |  |   |  |
| <u>Leptophlebia</u> sp.                    |  |   |  |
| Undet. Baetinae                            |  | 3 |  |
| Odonata: <u>Neurocordulia</u> sp.          |  |   |  |
| Neuroptera: <u>Corydalis cornutus</u>      |  |   |  |
| Trichoptera: <u>Hydropsyche simulans</u> , |  |   |  |
| Undet. <u>Hydropsyche</u>                  |  |   |  |
| <u>Macronemum carolina</u>                 |  | 2 |  |
| <u>Leptocella candida</u>                  |  |   |  |
| <u>Triaenodes</u> nr. <u>injusta</u>       |  |   |  |

(continued)

|                               |                 |        |        |        |        |        |
|-------------------------------|-----------------|--------|--------|--------|--------|--------|
| 18-171.59:171.59:171.59:170.5 | :170.5          | :169.0 | :167.4 | :155.3 | :152.0 | :146.4 |
| Ut-2.0                        | :Ut-0.1:Ut-0.1: | :      | :      | :      | :      | :      |
| 4/18                          | :4/18           | :4/18  | :4/18  | :4/18  | :5/12  | :4/18  |

(continued)

TABLE 12 (Continued) - SAVANNAH RIVER STUDIES  
ANIMALS ASSOCIATED WITH ALGAL MASSES

|                 |                                  |
|-----------------|----------------------------------|
| Location (River | : -171.62:171.59: 171.59:171.59. |
| Mile            | : :Ut-2.0: Ut-0.1:Ut-0.1.        |
| Date (1952)     | :4/18/52 :4/18 : 4/18 :4/18 :    |

ARTHROPODA:INSECTA (Continued)

|                                  |     |    |        |      |
|----------------------------------|-----|----|--------|------|
| Coleoptera:Stenelmis sp.         |     |    |        | 4A** |
| Macronychus glabratus            |     |    |        | 5A   |
| Undet. Hydroporinae              |     |    |        |      |
| Gyrinus sp.                      |     |    |        |      |
| Tropisternus sp.                 |     |    |        |      |
| Diptera:Agrotis variegata        |     |    |        |      |
| Ablatesmyia carnea               |     |    |        |      |
| Ablatesmyia monilis              |     |    |        |      |
| Undet. Ablatesmyia               |     |    |        |      |
| Ablatesmyia flavifrons           |     |    |        |      |
| or melanops                      |     |    |        |      |
| Ablatesmyia pilosella            |     |    |        |      |
| Procladius sp.                   |     | 1  |        |      |
| Coryphoptera scutellata(?)C.spp. | 1   |    | 1      |      |
| Thienemannella sp.               | 1   |    | 10     | 3    |
| Spanictora sens.lat.spp.         | 212 |    | 242,6* | 472  |
| Psectrocladius sp.               |     |    |        |      |
| Larytus sens nr. dispilms        |     | 2  |        |      |
| Larytus sens.lat.spp.            |     |    |        |      |
| Micropectra sp.                  |     |    |        |      |
| Microdiplos pedellus             |     |    |        |      |
| Polygaster illinoensis           | 3   |    |        |      |
| Undet. Polygaster                |     | 41 | 13     | 3    |
| Endochironomus nigricans         |     |    |        | 3    |
| Limnochironomus modestus         |     |    |        |      |
| Undet. Chironominae              |     |    |        |      |
| Polpomyia spp.                   |     |    |        |      |
| Stilobezzia sp.                  |     |    |        |      |
| Undet. Ceratopogonidae           |     |    |        |      |

MOLLUSCA:GASTROPODA

Undet.Flanorbidae

|                 |     |    |     |     |
|-----------------|-----|----|-----|-----|
| TOTAL ORGANISMS | 217 | 43 | 412 | 567 |
|-----------------|-----|----|-----|-----|

\* = Pupa

\*\* = Adult



2 OF 3

PB

229 203




TABLE 12 (Continued) - SAVANNAH RIVER STUDIES  
ANIMALS ASSOCIATED WITH ALGAL MASSES

S-171.59:171.59:171.59:170.5 :170.5 :169.0 :165.4 :155.3 :152.0 :140.4 :  
:Ut-2.0 :Ut-0.1:Ut-0.1:Ut-0.1 : : : : : : :  
:4/18 :4/18 :4/18 :4/18 :4/18 :7/23 :4/18 :5/12 :4/18 :4/18 :

|    |     |     |        |        |         |        |        |     |      |
|----|-----|-----|--------|--------|---------|--------|--------|-----|------|
|    |     |     | 1A**   | 2      | 8       |        |        |     |      |
|    |     |     | 1      |        |         |        |        |     |      |
|    |     |     |        |        | 1       |        |        |     |      |
|    |     |     | 1      | 1      | 1       |        |        |     |      |
|    |     |     |        | 101P** | 10      |        |        |     |      |
|    |     |     | 1      |        |         | 1      |        |     |      |
| 2  |     |     | 1      |        |         |        |        |     |      |
|    |     |     |        | 1      |         |        |        |     | 5    |
|    | 109 | 3   | 1      | 41     | 1       | 110,3P | 7      |     | 1    |
|    |     |     | 2      |        | 17      |        |        |     | 5    |
| 3  | 706 | 170 | 203,5P | 772    | 129,20P | 94     | 415,4P | 435 | 90   |
|    |     |     |        |        | 11      |        |        |     |      |
|    |     |     |        |        | 35,1P   |        | 2      |     | 1    |
|    |     |     |        |        |         |        |        |     | 1    |
|    |     |     |        |        |         |        |        |     | 1    |
|    |     |     |        | 2      |         |        |        | 111 |      |
| 2  | 2   | 1   | 14     |        | 6       | 7      | 242,1P | 15  |      |
|    |     |     |        | 1      |         |        |        | 3   | 4    |
|    |     |     |        |        | 29,1P   |        |        |     |      |
|    |     |     | 11     |        |         |        |        | 31  |      |
|    |     |     |        |        |         |        |        |     | 4    |
|    |     | 1   |        |        |         |        |        |     |      |
| 1  |     |     |        |        |         |        |        |     |      |
| 11 | 817 | 175 | 244    | 822    | 210     | 214    | 847    | 605 | 2191 |

\* P = Pupa

\*\* A = Adult

laboratory where the animals were separated from them. The algae were then identified, blotted dry, and weighed. The animals were preserved and later identified. The data from these determinations are recorded in Tables 11 and 12. The algal masses collected from the river and from the mouth of Upper Three Runs were all heavily populated with animal life. The sample from Dead River at S-146.4 had large numbers of microcrustacea present, but in all other cases chironomid larvae were the most abundant organisms. Spaniotoma was by far the most common chironomid genus present in all these algal masses. While Spaniotoma spp. were found widely distributed in brush box samples and bottom samples, they are principally algal feeders and were never found elsewhere in the high numbers that they attained in algal masses.

The samples collected from the area above the backwater (station) on Upper Three Runs showed much lower animal populations than the other samples in this series. Most of the algae in these particular samples were in a partially decayed condition, possibly because the growth of foliage on the trees had shaded many areas of the stream previously exposed to sunlight. The lower numbers of insects may be attributed to the fact that the decaying algae formed an unsuitable habitat for algal feeders such as Spaniotoma.

Any clarification of river water and stabilization of water levels which should result from the operation of Clark Hill Reservoir would increase the opportunities for the development of benthic algae by widening the photosynthetic zone, and would reduce the periodic stranding or submersion of the algae. These conditions would also result in increased numbers of algal-feeding invertebrates.

#### Organisms Associated with Compact Clay

On the outer side of every meander of the river and at many places between meanders, the banks of the river are quite steep and composed of compact clay. Wherever this clay which underlies the soil of the flood

plain, or wherever the resistant clays and marls of the Coastal Plain sediments are exposed to the current of the river, these substrates are

literally honey-combed with the burrows of mayfly nymphs. The mayfly

nymphs and the organisms found in their abandoned burrows form a characteristic assemblage of organisms that is to be found abundantly throughout the Coastal Plain portion of the river. Bottoms of clay, present where steep clay banks had crumbled into the river, had the same burrowing mayfly fauna as was found on the clay banks themselves.

Since the substrate occupied by these organisms is always rather hard, nearly always vertical, and is usually exposed to swift water, it is exceedingly difficult to sample adequately. A number of qualitative collections were made from these clay banks at miles S-171.7, S-165.5, and S-154.6 during June and July 1952. On July 30, 1952 a quantitative sample was taken from a clay bank at S-52.0. A block of clay was removed from the bank and its volume was measured by displacement. It was then broken up and washed through a series of screens. The organisms were picked out, identified, counted, and their volume determined by displacement.

Nymphs of the burrowing mayfly, Tortopus, were the most abundant organisms encountered in the qualitative collections. A variety of other organisms was also present. It is believed that the quantitative collection from S-52.0 recorded in Table 13 is fairly representative of compact clay substrates throughout the Coastal Plain portion of the river. In this collection Tortopus nymphs comprised 54% of the total number of organisms present and 94% of the total volume. Many of the Tortopus

nymphs appeared to be about to emerge as adults (i.e. in their last nymphal instar). Thus, the population of this species of organism might be assumed to be near its maximum volume.

The strategic location of these mayfly burrows at just the points where the current of the river impinges upon a firm substrate, undoubtedly aids the erosive action of the river against banks that would otherwise be very resistant to such attack. One peculiar feature of Tortopna is its apparent migratory nature. On many banks, abandoned mayfly burrows were found in abundance five or six feet above low water levels while both occupied and unoccupied burrows were at the same time found at least six feet below the water surface. Since the uppermost of these burrows could only have been occupied for a relatively short part of the nymphal life of a mayfly, it seems obvious that some individuals must move about in this honey-comb, building new burrows or occupying old ones above the normal water levels while the river is high, and retreating to succeeding lower ones as the river drops. Considering the area of exposed surface represented, the numbers and volume of organisms taken from the quantitative sample were high. The organisms found in association with Tortopna do not usually burrow in hard clay themselves and the burrowing activities of the mayflies provided them with a suitable habitat. This one quantitative sample indicates that large populations of organisms are present in this habitat. This assumption is substantiated by the qualitative sampling of clay banks elsewhere in the river which appeared to parallel this one very closely. Since clay banks are present at many places on the



TABLE 13 - QUANTITATIVE CLAY BANK SAMPLE 3-52.0  
SAVANNAH RIVER STUDIES  
JULY 30, 1952

Volume of clay substrate - 8,000 cc., Area of Exposed surface 1,600 cm<sup>2</sup>

|                        |                                   | Number | Volume<br>(cc.) |
|------------------------|-----------------------------------|--------|-----------------|
| ORGANISMS              |                                   |        |                 |
| Nematoda:              | Undetermined                      | 107    | 0.05            |
| Arthropoda, Insecta */ |                                   |        |                 |
| Plecoptera:            | <u>Acroneuria</u> sp.             | 4      | 0.05            |
| Ephemeroptera:         | <u>Tortopus</u> sp. last instar   | 57     | 4.80            |
|                        | <u>Tortopus</u> sp. other instars | 457    | 22.40           |
|                        | <u>Stenonema</u> spp.             | 18     | 0.02            |
|                        | <u>Isonychia</u> spp.             | 99     | 0.24            |
|                        | <u>Tricorythodes</u> sp.          | 4      | 0.00            |
|                        | <u>Caenis</u> spp.                | 5      | 0.01            |
|                        | Baetinae, undetermined            | 1      | 0.00            |
| Odonata:               | Cordulinae, undetermined          | 1      | 0.00            |
|                        | <u>Argia</u> sp.                  | 1      | 0.00            |
| Neuroptera:            | <u>Corydalus cornutus</u>         | 5      | 0.80            |
| Trichoptera:           | <u>Cheumatopsyche</u> sp.         | 5      | 0.00            |
|                        | <u>Macronema carolina</u>         | 15     | 0.15            |
|                        | <u>Macronema carolina</u> , pupa  | 1      | 0.04            |
|                        | <u>Potamyia flava</u>             | 1      | 0.00            |
|                        | <u>Stenelmis decorata</u> , adult | 117    | 0.17            |
| Coleoptera:            | <u>Stenelmis</u> sp., larva       | 26     | 0.05            |
|                        | <u>Palpomyia</u>                  | 1      | 0.00            |
| Diptera:               | <u>Pentaneura</u> (monilis group) | 4      | 0.00            |
|                        | Chironomidae, undetermined        | 2      | 0.00            |
| Mollusca               |                                   |        |                 |
| Gastropoda:            | <u>Pleurocera</u> sp.             | 15     | 0.10            |
| Pelecypoda:            | <u>Sphaerium</u> sp.              | 1      | 0.00            |
| TOTAL                  |                                   | 947    | 23.88           |

\* Unless otherwise indicated, all Arthropoda were represented by immature forms.

river, this community of organisms, dominated by Tortopus, appears to be an important part of the Savannah River fauna.

#### Organisms Associated with Sand and Mud

The most extensive and widespread substrates available to benthic organisms in the Savannah River are the sands and silts of the river bottom proper. Much of this material is constantly undergoing erosion and deposition. Such instability greatly limits the potentiality of supporting any sizeable community of organisms.

Areas of sand and mud bottoms were often contiguous, or intermingled, and the same technique was necessarily used for sampling both; thus, these two types of substrate are herein considered together. Bottom samples were taken by Petersen dredge at miles S-223 (Stephens Creek Impoundment), S-203.3 (New Savannah Bluff Impoundment), S-201.55, S-171.7, S-165.5, S-146.4 (Dead River), S-141.0, and S-51.0.

Except in the case of the impoundments, which had fairly uniform bottoms, samples were taken in transects across the river. The spacing of the samples across each transect depended upon the rate of flow of the river and the nature of the bottom materials. Areas that appeared to be uniform with regard to bottom materials, depth, and current speed, were sampled but once on any given transect. The numbers of organisms collected from mud and sand substrates were generally small as compared with those taken from other substrates. The numbers of individuals and species present depended upon the nature of the bottom sampled. The areas of shifting sand, which formed the majority of the bottom materials in areas of swift current, were practically devoid of macroscopic



organisms. Bottoms of sand, mud, and organic debris such as those present in impoundments and backwaters, and other areas where deposition of silt was taking place, supported larger numbers and a greater variety of macroscopic organisms. Transects across the river at stations where the several different bottom types were present showed a fauna varying from one side of the stream to the other. (Table 14).

Midge larvae were the most ubiquitous organisms in the bottom fauna. Chironomids or Ceratopogonids were present in almost every bottom sample taken. The greatest numbers of chironomids and oligochaeta worms were taken from a sample on the Georgia side of the river at mile S-201.55, apparently reflecting the organic enrichment of the river by the sewage from Butler Creek. Several genera of burrowing mayflies other than Tortopus were collected, and the emergence of one of these genera was observed in July 1952. Large numbers of Hexagenia bilineata were seen swarming on willows above areas of soft mud bottom at miles S-146.4, S-141.0, and S-130.0.

Fresh-water mussels were moderately abundant as was evidenced by the numbers of dead shells along the shores. This important group of organisms was inadequately sampled by dredge. The low river levels prevailing during the fall of 1951 and the summer of 1952 made it possible to collect a number of living mussels by hand. In 1952 it was noted that large numbers of these animals were killed by being stranded in drying pools behind sand bars. Mussels have yet to be identified.

The populations of organisms living in sand and mud habitats were generally low, particularly in the areas of shifting sand. However, since these areas afford the most extensive habitats available to benthic

TABLE NO. 14 - SAVANNAH RIVER STUDIES

## SUMMARY

## Bottom Samples \*

|                        |         |         |          |       |   |      |
|------------------------|---------|---------|----------|-------|---|------|
| Sampling Point         | S-223.0 | S-203.3 | S-201.55 |       |   |      |
| Date                   | 8/23/51 | 9/25/51 | 8/1/52   |       |   |      |
| Depth (Feet)           | 6       | 8       | 3        | 2     | 9 | 9    |
| Bottom Materials **    | MS,OD   | MS      | MS,OD    | SG,OD | S | S,OD |
| Number of Dredge Grabs | 2       | 4       | 3        | 3     | 3 | 3    |

## NEMATODA

Undetermined

## ANNELIDA: OLIGOCHAETA

Pelosclex sp.

Undetermined

22

3

## ANNELIDA: HIRUDINEA

Helobdella

2

## ARTHROPODA: INSECTA

Ephemeroptera: Ephemera sp.Hexagenia sp.

6

Tortopus sp.Pentagenia sp.Stenonema spp.Isonychia spp.Ephemerella sp.Caenis sp.Odonata: Gomphus sp.Argia sp.Neuroptera: Sialis sp.

1

Corydalis cornutusTrichoptera: Chimarra aterimaMacronemum carolina

6

3

Undet. Hydroptilidae

8P

Leptocella sp.

1P

Coleoptera: Macronychus glabratusStenelmis sp.

29

1

\*

All organisms represent larval  
forms except where indicated

A = Adult

P = Pupa

\*\*

M = mud

S = sand

C = clay

G = gravel

OD = organic debris

P = Phormidium

(continued)

TABLE NO. 14 - SAVANNAH RIVER STUDIES (Continued)

SUMMARY

Bottom Samples \*

| S-171.7 |       |      |        | S-165.5 |    |   |   | S-146.4  |   |   |       | S-141.0 |   |    |       | S-51.0  |      |      |      |
|---------|-------|------|--------|---------|----|---|---|----------|---|---|-------|---------|---|----|-------|---------|------|------|------|
| 7/23/52 |       |      |        | 7/21/52 |    |   |   | 11/12/51 |   |   |       | 7/10/52 |   |    |       | 7/31/52 |      |      |      |
| 12      | 8     | 8    | 3      | 10      | 10 | 3 | 5 | 4        | 3 | 3 | 6     | 11      | 8 | 12 | --    | 2       | 6    | 8    | 7    |
| **C     | SC,OD | S,OD | S,C,OD | S       | S  | S | M | M        | M | M | MS,OD | SP      | S | S  | MS,OD | M,OD    | S,OD | S,OD | C,OD |
| 3       | 3     | 3    | 3      | 3       | 3  | 3 | 2 | 2        | 2 | 2 | 3     | 3       | 3 | 3  | 3     | 3       | 3    | 3    | 3    |

1

1

7

9

2

1

22

3

11

1

2

149

7

1

9

1

1

1

1

1

1

1

1

1

1

10

1P

13

1

3,6A 6

2A

19,1A

1A

2  
1A

\*/ All organisms represent larval forms except where indicated  
A = Adult      P = Pupa

\*\*/ M = mud  
S = sand  
C = clay  
G = gravel  
OD = organic debris  
P = phormidium

(continued)

TABLE 14 - SAVANNAH RIVER STUDIES (continued)

## SUMMARY

## Bottom Samples \*

| Sampling Point         | S-223.0 | S-203.3 | S-201.55 |        |   |       |
|------------------------|---------|---------|----------|--------|---|-------|
| Date:                  | 8/23/51 | 9/25/51 | 8/1/52   |        |   |       |
| Depth (Feet)           | 6       | 8       | 3        | 2      | 9 | 9     |
| Bottom Materials **    | MS, OD  | MS      | MS, OD   | SG, OD | S | S, OD |
| Number of Dredge Grabs | 2       | 4       | 3        | 3      | 3 | 3     |

## ARTHEROPODA: INSECTA (continued)

## Diptera: Chaoborus sp.

Ablabesmyia monilis

1P

1

Ablabesmyia flavifronsAblabesmyia pilosella

1

1

Procladius sp.Coelotanytus nr. concinusSpaniotoma sens. lat. spp.

3

Tanytarsus sens. lat. spp.Stenochironomus exquisitus

1

Stenochironomus flavicingula

1

1

Polypedilum sp.Endochironomus tendens

1

Cryptochironomus stylifer

9

1

Cryptochironomus sp.

2

Limnochironomus modestus

19

30

Parachironomus tenuicaudatus

84

Limnochironomus sp.

8

Chironomus sp.

Undet. Chironomidae

1

Undet. Ceratopogonidae

## MOLLUSCA: GASTROPODA

Physa sp.Ferrissia sp.

## MOLLUSCA: PELECYPODA

Pisidium sp.

1

Sphaerium sp.

7

Musculium sp.

4

unionidae

1

## TOTAL ORGANISMS

10 39 124 39 0 48

\* All organisms represent larval forms except where indicated  
A = Adult P = Pupa

\*\* M = mud  
S = sand  
C = clay  
G = gravel  
OD = organic debris  
P = phormidium

TABLE NO. 14 - SAVANNAH RIVER STUDIES (Continued)

## SUMMARY - Bottom Samples \*

| S-171.7 |       |      |   | S-165.5 |    |   |   | S-146.4  |   |   |       | S-141.0 |   |    |       | S-51.0  |      |      |      |
|---------|-------|------|---|---------|----|---|---|----------|---|---|-------|---------|---|----|-------|---------|------|------|------|
| 7/23/52 |       |      |   | 7/21/52 |    |   |   | 11/12/51 |   |   |       | 7/10/52 |   |    |       | 7/31/52 |      |      |      |
| 12      | 8     | 8    | 3 | 10      | 10 | 3 | 5 | 4        | 3 | 3 | 6     | 11      | 8 | 12 | --    | 2       | 6    | 8    | 7    |
| **C     | SG,OD | S,OD | S | C,OD    | S  | S | M | M        | M | M | MS,OD | SO      | S | S  | MS,OD | M,OD    | S,OD | S,OD | C,OD |
| 3       | 3     | 3    | 3 | 3       | 3  | 3 | 2 | 2        | 2 | 2 | 3     | 3       | 3 | 3  | 3     | 3       | 3    | 3    | 3    |

4 2 4  
 1 2 2  
 1 5  
 16 12 15 1  
 2 2 1  
 1 1 8 1  
 3 4 1  
 8 1 1 1  
 1 1 1  
 2 1 1  
 1 3 1,1P  
 8 3  
 6 2 7 1 1 9  
 1 1  
 1  
 1  
 1  
 173 22 11 8 33 0 3 25 19 26 7 23 28 9 0 1 38 1 30 11

\* All organisms represent larval  
 forms except where indicated  
 A = Adult P = Pupa

\*\* M = mud  
 S = sand  
 C = clay  
 G = gravel  
 OD = organic debris  
 P = phormidium



invertebrates, they must be considered somewhat more important in the productive economy of the river than the low densities of the organisms inhabiting them indicate.

#### Organisms Associated with Brush, Trees, Rocks, Gravel, etc.

Piles of tangled debris, formed by roots, fallen trees, brush and materials swept into the river from its flood plain and tributary streams, were abundant in the river. It was observed early in the study that these substrates often supported large and diverse populations of invertebrates. Natural outcroppings of sandstone and chert, and banks riprapped with granite also provide stable substrates for invertebrates. Such areas were not common.

Devices called "brush boxes" were constructed in order to obtain representative samples of the organisms occupying such substrates. (Appendix G). Table 15 presents summarized data of such collections; complete data are shown on Table 1, Appendix "G".

The use of brush boxes to sample benthic organisms has several advantages over random collection. Brush boxes may be set in almost any situation from which a sample is desired. Thus, the investigator is not dependent upon the chance occurrence of suitable substrates at a location of particular interest and samples may be easily obtained from situations varying with regard to depth, current, bottom materials, proximity to strategic tributaries, and other considerations. The groups of organisms collected from boxes placed at different stations may be compared on a relative basis since the materials placed in the boxes offer similar substrates for colonization. Thus the species composition of the population



of any one insect group, for example, the diptera, may be compared from one box to another. Brush box sampling is apt to be less selective than some other methods of collecting since a given amount of the substrate material from a definable environmental situation is examined in detail regardless of the paucity or richness of the fauna.

The limitations encountered in the use of brush boxes were primarily due to the loss and silting-in of boxes. If the wires holding the boxes were fixed to the bank below the water surface, where they could not be seen and tampered with by curious persons, a rise in the river level made them difficult to find and recover. Boxes placed in strong current risked being carried away completely, as they became clogged with drifting debris, while those placed in very sheltered situations occasionally became deeply covered by silt.

In order to establish a standard time interval of convenient length during which a representative fauna could accumulate, a series of four boxes was set side by side at mile S-219 on July 23, 1951, and raised after intervals of 7, 14, 21, and 28 days. A comparison of the columns under S-219 in Table 15 reveals that most of the groups of organisms encountered were fairly well represented after 21 days, at least at this season of the year. Accordingly, 21 days was chosen as the standard time for leaving brush boxes in place. Vagaries of weather and river levels did not allow the raising of every box exactly on schedule, but 20 of the boxes placed were allowed to remain near enough to the standard length of time to make them comparable on a time basis. Other boxes were allowed to remain in the river for periods up to 165 days.

These long term boxes were used to determine whether or not organisms would appear which were different from those found in the 21-day boxes. Data from comparable boxes are available from mile S-201.55 (one box left for 21 days, the other for 165 days) and from mile S-165.4 (one box left for 21 days, the other for 139 days). In both these instances the long-term member of the pair contained fewer organisms of the type which cling closely to the substrate (Plecoptera, Ephemeroptera and Trichoptera) than did their 21-day counterparts. This difference can be attributed to the accumulation of slime and silt on the available surfaces. In both these cases the boxes were set where the current was rather weak and such accumulations could take place over long periods of time. Both long-term boxes contained a few burrowing Odonata while the 21-day boxes contained none. The boxes set at mile S-201.55 were in an area heavily enriched by sewage from Augusta, they thus contained appreciable numbers of such forms as oligochaeta, gastropoda and sphaeriid clams. All these groups showed a marked increase in numbers in the 165-day box set at this locality over those in the 21-day box. The presence of many Chironomus riparius in the 165-day box at mile S-201.55 reflects both the pollution and the physical environment at this station. The dipteran populations in each of these pairs of boxes are remarkably similar both quantitatively and qualitatively. Spongille fragilis was the only organism found in the long-term boxes which was not found in any of the short-term boxes. It occurred in detectable quantities in only two boxes that had been left in the river for 165 and 139 days thus indicating that this form requires a fairly long period for its development.

Since most of the brush boxes were set below the photosynthetic zone, herbivorous animals dependent upon algal growth were relatively scarce. A comparison of the dipteran fauna collected from brush boxes with those collected from algal masses substantiates this assumption. Spaniotoma, a herbivore and the most abundant dipteran in the algal masses, was present in most of the brush boxes, but it was never very common, nor did it ever form a predominant part of the dipteran fauna.

The highest total numbers of organisms were taken from boxes set in the polluted waters below Horse Creek and Butler Creek, both in moderate current, and from boxes set in swift water at miles S-171.7 and S-171.15, although the species compositions from these areas varied considerably. Brush boxes placed in moderate current at the two latter stations did not become as heavily populated as those in swift water at the same place or those in comparable current at polluted stations.

General observations on the distribution of the various groups of organisms in brush box collections follow.

Porifera - Spongilla was found in only two cases, both in boxes that had been in the river over 100 days. It was abundant in the box at S-201.55 below the pollution from Butler Creek.

Turbellaria - Dugesia tigrina was found at widespread locations throughout the river, but it was very abundant only in the boxes from the polluted areas below Horse Creek (S-213.3) and Butler Creek.

Rotatoria - Although many kinds of rotifers were present in the river, the only one readily collected because of its large size in brush box sampling, was the sessile, colonial form, Floscularia, sp. It was present at a

TABLE NO. 15

## SAVANNAH RIVER STUDIES

July 1951-July 1952

SUMMARY - Brush Box Sampling

Numbers of Organisms

| Location (River Mi.) | S-223.15 | S-219.0 |        |         |         |          |         | S-215.5 |
|----------------------|----------|---------|--------|---------|---------|----------|---------|---------|
| Date Removed         | 8/24/51  | 7/30/51 | 8/6/51 | 8/13/51 | 8/20/51 | 10/16/51 | 8/27/51 |         |
| No. of days in place | 21       | 7       | 14     | 21      | 28      | 21       | 21      |         |

## PORIFERA

Total

## TURBELLARIA

Total

1

## ROTATORIA

Total

Pr.\*

Pr.

Pr.

Pr.

## BRYOZOA

Total

Pr.

## ANNELIDA: OLIGOCHAETA

Total

8

2

9

## ANNELIDA: HIRUDINEA

Total

3

1

## ARTHROPODA: CRUSTACEA

Copepoda

Ostracoda

Isopoda

Amphipoda

Decapoda

145

6

10

43

16

1

Total CRUSTACEA

145

6

10

44

16

## ARTHROPODA: ARACHNIDA

Total

13

## ARTHROPODA: INSECTA

Plecoptera

Ephemeroptera

Odonata

Neuroptera

Trichoptera

Hemiptera

Coleoptera

Diptera

28

46

50

51

81

58

11

34

30

24

26

56

64

11

1

8

12

21

16

25

15

6

7

12

27

11

40

30

109

30

214

216

292

161

125

Total INSECTA

187

130

336

320

494

328

153

## MOLLUSCA: GASTROPODA

Total

8

3

1

2

20

14

43

## MOLLUSCA: PELECYPODA

Total

1

## TOTAL ORGANISMS

365

133

337

332

536

386

61

\* Pr. = Present

(continued)

TABLE NO. 15 (Continued) SAVANNAH RIVER STUDIES July 1951-July 1952  
SUMMARY - Brush Box Sampling - Numbers of Organisms

| 8-215.5 |          | 8-213.3 | 8-203.3 | 8-201.55 |         | 8-172.0 |
|---------|----------|---------|---------|----------|---------|---------|
| 9/19/51 | 10/16/51 | 9/19/51 | 8/27/51 | 8/28/51  | 7/27/52 | 9/4/51  |
| 21      | 21       | 21      | 21      | 22       | 165     | 21      |

Pr.

|      |     |      |     |      |      |     |
|------|-----|------|-----|------|------|-----|
| 9    | 1   | 238  |     | 12   | 174  |     |
| Pr.* | Pr. | Pr.  | Pr. | Pr.  | Pr.  |     |
|      |     |      |     |      | Pr.  | Pr. |
|      |     | 62   | 3   | 14   | 399  |     |
| 3    |     | 31   |     | 1    | 16   |     |
| 79   | 1   | 20   | 3   |      |      |     |
| 79   | 1   | 20   | 3   |      |      |     |
| 3    |     | 5    |     | 4    | 3    |     |
| 7    | 1   | 1    | 13  | 71   | 2    | 39  |
| 24   | 26  | 66   | 28  | 72   | 30   | 42  |
| 28   | 6   | 147  | 4   | 182  | 27   | 1   |
| 3    | 2   | 6    |     | 5    | 69   | 40  |
| 553  | 33  | 1125 | 318 | 692  | 640  | 50  |
| 615  | 68  | 1345 | 363 | 1022 | 768  | 203 |
| 159  | 281 | 122  | 1   | 11   | 73   |     |
| 1    | 1   | 11   |     | 9    | 74   |     |
| 870  | 353 | 1835 | 371 | 1074 | 1510 | 204 |

\*/ Pr. = Present

(continued)



TABLE NO. 15 (continued)

## SAVANNAH RIVER STUDIES

July 1951-July 1952

SUMMARY - Brush Box Sampling

Numbers of Organisms

| Location (River Mi.) | S-171.7 | S-171.6  | S-171.52 |          | S-171.15 | S-165.4 |         |
|----------------------|---------|----------|----------|----------|----------|---------|---------|
| Date Removed         | 10/3/51 | 12/20/51 | 10/4/51  | 10/31/51 | 9/4/51   | 9/4/51  | 7/17/52 |
| No. of days in place | 21      | 37       | 22       | 26       | 21       | 21      | 139     |

Pr.

## PORIFERA

Total

## TURBELLARIA

Total

1

1

4

## ROTATORIA

Total

Pr.\*

Pr.

## BRYOZOA

Total

Pr.

Pr.

## ANNELIDA: OLIGOCHAETA

Total

1

1

60

## ANNELIDA: HIRUDINEA

Total

1

## ARTHROPODA: CRUSTACEA

Copepoda

Ostracoda

Isopoda

Amphipoda

Decapoda

1

Total CRUSTACEA

1

1

1

## ARTHROPODA: ARACHNIDA

Total

1

5

1

## ARTHROPODA: INSECTA

Plecoptera

Ephemeroptera

Odonata

Neuroptera

Trichoptera

Hemiptera

Coleoptera

Diptera

87

476

35

11

3413

107

720

40

47

2

54

21

105

37

384

88

2

142

516

154

49

254

16

2

630

164

372

18

527

134

13

4113

105

547

17

69

29

15

2643

50

117

3

4

5

61

27

71

## MOLLUSCA: GASTROPODA

Total

1

1

1

3

## MOLLUSCA: PELECYPODA

Total

1

TOTAL ORGANISMS

4851

270

1330

1491

5465

2943

237

\*/ Pr. = Present



TABLE NO. 15 (Continued) SAVANNAH RIVER STUDIES July 1951-July 1952  
SUMMARY - Brush Box Sampling - Numbers of Organisms

| 8-160.75 | 8-155.25 | 8-155.2<br>8-155.2 | 8-155.17 | 8-146.4  | 8-144.7  | 8-136.6 | 8-51.0  |
|----------|----------|--------------------|----------|----------|----------|---------|---------|
| 11/2/51  | 10/1/51  | 9/26/51            | 9/6/51   | 10/11/51 | 10/11/51 | 2/6/52  | 7/31/52 |
| 32       | 24       | 20                 | 22       | 21       | 21       | 53      | 160     |

Pr.\*

Pr.

|      |      |     |      |     |      |     |      |
|------|------|-----|------|-----|------|-----|------|
| 1    | 8    | 3   |      |     |      |     |      |
| 6    | 1    | 4   | 2    |     |      | 5   |      |
|      | 2    |     | 1    |     | 2    |     |      |
|      | 13   |     | 6    |     |      |     |      |
|      |      |     | 1    |     |      |     |      |
| 2    |      |     |      |     | 1    |     |      |
| 2    | 14   |     | 7    |     | 1    |     |      |
| 2    | 2    | 6   | 33   |     | 8    |     |      |
| 45   | 7    |     | 15   | 1   | 75   | 33  | 7    |
| 317  | 168  | 1   | 81   | 7   | 514  | 150 | 154  |
| 31   | 34   | 24  | 38   | 39  | 2    | 2   | 37   |
| 2    |      |     | 7    |     | 1    | 1   | 2    |
| 3437 | 522  | 17  | 1047 | 8   | 2252 | 86  | 857  |
| 104  | 12   | 30  | 109  | 19  | 151  | 69  | 242  |
| 1021 | 440  | 120 | 324  | 73  | 2362 | 34  | 81   |
| 5132 | 1183 | 192 | 1621 | 147 | 5357 | 375 | 1380 |
| 17   |      | 1   | 1    | 2   | 9    |     |      |
| 9    |      |     | 1    |     |      |     |      |
| 5171 | 1186 | 227 | 1625 | 193 | 5366 | 386 | 1386 |

\* / Pr. = Present

number of stations but was most abundant at polluted water stations.

Bryozoa - Plumatella repens was found at a number of stations, but it appeared to be most abundant at polluted water stations.

Annelida - Oligochaeta and leeches were both found at a number of stations but were most abundant at polluted water stations.

Crustacea - The numbers of small crustaceans, such as copepods and ostracods, recorded in the accompanying tables are not reliable indications of their relative abundance at various localities. These small organisms were difficult to detect in a sample, and the greatest numbers were taken when the samples were sorted in living condition and the animals could be seen swimming about in the sorting tray. The large free-swimming river shrimps, Macrobrachium ohione and Palaemonetes paludosus, occurred occasionally in the brush boxes.

Arachnida - The small water mites presented the same limitations to estimation of their relative abundance in various localities as did the smaller crustacea.

Insecta - The insects were the most diverse and abundant group obtained in brush box sampling.

The stoneflies (Plecoptera) were found in abundance wherever brush boxes were set in fast water at unpolluted stations.

Mayflies (Ephemeroptera) were present in all of the samples. Burrowing mayflies occurred where boxes were set in soft bottoms or when they accumulated silt deposits. The most abundant genera of mayflies were Stenonema, Oaenis, and Tricorythodes. Only one nymph (Hexagenia) was taken from the polluted area below Horse Creek, and one from Steel Creek.

Dragonflies and damselflies (Odonata) were taken in all the boxes except the one from Upper Three Runs. Burrowing dragonflies were present in boxes where accumulating silt provided suitable habitats for them. Climbing nymphs such as those of Argia, Macromia, and Neurocordulia were the most frequently encountered forms.

Larvae of the dobson fly, Corydalis cornutus, (Neuroptera), a carnivorous species, was common in areas of fast water where large numbers of other insects were present.

The caddisflies (Trichoptera) were the most abundant organisms in many of the brush boxes. Macronemum carolina, Hydropsyche spp. and Neuraclipsis sp. were extremely abundant in areas with swift current where leaves and detritus accumulated on the box.

Only one water strider (Hemiptera) was taken in brush box sampling, but since these insects live on the surface of the water, its occurrence in this sample was probably accidental.

Three genera of elmids beetles (Coleoptera) which were common in almost all the samples were Ancyronyx, Macronychus, and Stenelmia. Berosus was found only in polluted areas. Other beetles were scattered in distribution.

The bulk of the dipteran populations in the brush boxes was composed of midge larvae (Chironomidae). Ablabesmyia monilia, Limnochironomus modestus and Chironomus riparius were abundant in polluted areas. Endochironomus nigricans was abundant in polluted water, but not in swift current. Polypadilum spp. were abundant in swift water, as were Tanytarsus spp., but the latter forms were absent from polluted areas.

The snipe-fly, Atherix variegata, and black-fly, Simulium sp. were both found in Upper Three Runs and in swift-water areas below that point. Ceratopogonids were found in scattered localities, but never in large numbers.

Mollusca: Both pulmonate and gilled snails were most abundant in polluted areas. Fingernail clams, Sphaeriidae, occurred when bottom deposits accumulated in the brush boxes. They were also most abundant in polluted areas.

#### Nekton

The four principal objectives of the investigation of fish populations in the river were to determine: (a) the species composition of the population; (b) the extent of utilization by man; (c) some important aspects of the biology of the principal species (particularly catfish); and (d) fish movement.

Three methods were used to obtain information applicable to the above objectives; (a) the observation of commercial and sport fishing, (b) the use of seines, and nets, for sampling, and (c) the use of rotenone to obtain samples.

Each of these methods yielded different information in the study of fish populations and will be discussed separately.

#### COMMERCIAL AND SPORT FISHING

##### Commercial Fishing

Two types of commercial fishing were observed in the fresh-water areas of the Savannah River; a basket fishery, principally for channel catfish, and a gill net fishery for American shad and other anadromous

herrings. Contacts with the fishermen engaged in these activities were entirely fortuitous due to the scattered and unsystematic nature of the commercial fishing on the river. Observations of sport fishing were limited to chance interviews with sport fishermen encountered during the course of the study.

#### Basket Fishing for Catfish

The basket fishery for catfish is the most intensive commercial fishery in the fresh-water areas of the river. Three fishermen were observed operating regularly in the area from Augusta (S-216.0) to Porter's Landing (S-74.3), and others were observed operating in the area around Ebenezer Landing (S-48.1) and Lockner Creek (S-46.33). None of them was entirely dependent upon commercial fishing for a livelihood. The wire baskets used in this fishery are highly selective for catfish. The principal and most valuable species taken was the southern channel catfish (Ictalurus lacustris punctatus). The two other catfish appearing regularly in catches were the white catfish (Ictalurus catus) and the flat bullhead (Ameiurus platyrhynchus). These were of little commercial value. A number of other species occurred occasionally in the catch.

Selected fish from several catches were examined in July and August 1951, and twelve total catches were examined from August 1951 to May 1952. In examination of these catches all fish were identified and measured. The sex of some mature individuals was determined and pectoral spines were removed from the catfish. <sup>2/</sup>

---

<sup>2/</sup> These spines are being retained at the Department of Biology, University of Georgia, for age and growth rate studies.



Table 16 shows a summary of these catches. Channel catfish comprised 84.3% of the catch, white catfish 10.6%, flat bullheads 3.6%, and other species 1.5%. The length-frequency distributions of catfish from commercial catches are shown in Figures 28, 29, and 30.

The size of the catches was extremely variable. The same fisherman reported catches of as much as 700 and as little as 50 pounds per week. The fishery was somewhat seasonal; fishermen reported fishing to be best in the spring and the fall and poor during the summer months. The commercial value of these catches to the fisherman averaged about 40 cents per pound live weight. Most of the fish were sold to eating establishments located near the river.

#### Shad Fishing

The American shad and two or three other anadromous species of herring ascend the river to spawn in late winter and spring and are subjected to a limited local fishery. In 1952 the operation of this fishery was observed directly by interviewing fishermen and accompanying them on the rounds of their nets whenever possible. The large meshed (5 - 6" stretched) gill nets used in this fishery are either placed at stationary sets in eddy waters, or are drifted down straight stretches of river channel. Fishing for the two commercially valuable species, American shad (Alosa sapidissima) and hickory shad (Pomolobus mediocris), began early in January on the lower river (around mile S-48), and about a month later at mile S-146. Fishing continued into the month of April when longnose gar (Lepisosteus osseus) and bowfin (Amia calva) became so numerous in the catches as to interfere with the operation of the nets and thus






FIG. 28




FIG. 29




FIG. 30

render fishing unprofitable. After mid-April, a large percentage of the caught shad were spent and worthless. The last spent shad observed in the river was seen floating at mile S-155.3 on June 16.

Seven gill net catches, in which 100 American shad and 19 hickory shad were present, were examined. The length, weight, and sex were recorded for each fish. A scale sample was removed from each fish for age determination. <sup>\*</sup>/

Table 17 summarizes the catches examined. Considering the shad only, the American shad comprised 34% of the catch and the hickory shad 16%. These figures are not considered to represent the relative abundance of the two species in the river since the large meshed nets of the fishery are obviously selective for the larger species. This is evidenced by the fact that many more females than males of both species were caught, the females averaging considerably larger than the males. The length-frequency distributions of the shad from these catches are shown on Figure 31.

At least two other species of anadromous clupeid fishes ascend the Savannah River to spawn. The glut herring (Pomolobus aestivalis), a small but abundant species, is of no commercial value but is often taken in small-meshed gill nets to be used as cut bait. Fishermen report that another clupeid (probably the alewife, Pomolobus pseudo-harengus) preceded the shad run, but specimens of this fish were not obtained so its identity remains in doubt. The striped bass (Morone saxatilis)

\* Scale samples have been transmitted to the United States Fish and Wildlife Service for use in their study of shad in the southeastern States.

saxtilis) is occasionally caught in the shad nets, as is the sturgeon (probably Acipenser oxyrhynchus).

Fishermen operating in the lower river rated shad fishing as fairly good in 1952 when compared with that of the previous few years. Many of the fish caught in this area are sold in Savannah, Springfield and neighboring towns. The American shad were sold for from \$1.25 to \$2.00 each, while the hickory shad usually brought about \$.75 each. This type of fishing seemed to be engaged in as much for sport as for profit by some of the fishermen interviewed.

#### Sport Fishing

Observations of sport fishing on the Savannah River indicate that hook-and-line fishing may be classified as follows:

1. Bank line, trot line and cane pole fishing for channel catfish.
2. Cane pole fishing and casting for largemouth black bass, crappie, and the various species of pan-sized sunfish.
3. Casting for striped bass.

The value and importance of sport fishing on any large stream is difficult to assess. Very few sport fishermen were encountered on the river proper and sport fishing seemed to be much more intensive on some of the tributary streams and flood plain lakes than on the river itself. While sport fishing on the river itself was apparently of negligible importance at the time these studies were made, it might well increase considerably if the operation of Clark Hill Dam causes the river to become clearer. At the same time increasing human population in the

Augusta area will no doubt result in greater use of the river for sport fishing.

#### SEINE AND NET SAMPLING

##### Hoop Netting

Hoop nets with a diameter of five feet and mesh size of approximately one inch were used to collect fishes for radiological background determinations and to obtain additional information on the fish populations of the river. The nets were baited with cottonseed-meal cake and usually set in fast water. This type of sampling was conducted at intervals from December 1951 into July 1952, over the stretch of river from S-172.5 to S-140.1. A total of 117 net hauls were made during this period. The results of this sampling are shown on Table 18.

Hoop nets operate in a manner similar to that of the wire baskets used by commercial fishermen and they were set in much the same situations; however, they appeared to be somewhat less selective for catfish. Catfish comprised 85.6% of the commercial catches examined and 77.8% of the hoop net catches. Considering only the three common species of catfish, these differences in the composition of the catch are shown in Table 19.

TABLE 16

SAVANNAH RIVER STUDIES  
BASKET FISHERY  
Commercial Catches Examined

| Date              | Location<br>River Miles | Numbers of Fish    |                  |                  | Others**/<br>No. | Total<br>No. | Total<br>Weight<br>Pounds |
|-------------------|-------------------------|--------------------|------------------|------------------|------------------|--------------|---------------------------|
|                   |                         | Channel<br>Catfish | White<br>Catfish | Flat<br>Bullhead |                  |              |                           |
| 8/31/51           | S-91.7 - S-74.3         | 309                | 20               | 13               |                  | 340          | 167                       |
| 9/7/51            | " "                     | 198                | 15               | 3                |                  | 216          | 100                       |
| 9/14/51           | " "                     | 117                | 3                | 6                | 13               | 139          | 78                        |
| 9/28/51*          | Vicinity S-122          | 133                | 58               | 21               |                  | 212          | 150                       |
| 10/5/51           | S-152.2 -S-146.4        | 271                | 65               | 8                |                  | 344          | 284                       |
| 10/13/51          | " " "                   | 62                 | 12               | 3                |                  | 77           | -                         |
| 10/23/51          | " "                     | 42                 | 2                | 4                |                  | 48           | -                         |
| 10/26/51          | " "                     | 124                | 9                |                  | 5                | 138          | -                         |
| 10/29/51          | " "                     | 235                | 11               |                  | 3                | 249          | -                         |
| 11/22/51          | " "                     | 91                 | 20               | 4                | 1                | 116          | 141                       |
| 11/30/51          | " "                     | 31                 | 7                | 8                | 7                | 53           | -                         |
| 5/12/52           | " "                     | 152                |                  | 7                | 2                | 161          | -                         |
| TOTAL NO.         |                         | 1,765              | 222              | 75               | 31               | 2,093        |                           |
| % of Total Catch  |                         | 84.3               | 10.6             | 3.6              | 1.5              |              |                           |
| Average Length mm |                         | 294                | 340              | 226              |                  |              |                           |

\*/ Figures represent approximately 75% of total catch

\*\*/ American Eel, Red Horse Suckers, Carp, Brown Bullhead, White Crappie, Redbelly Sunfish, Striped Bass.



TABLE 17  
SAVANNAH RIVER STUDIES  
SHAD FISHERY  
Commercial Catches Examined

| Date                   | Location<br>River Miles | Numbers of Fish      |                              |                          |                            | Total<br>No. |
|------------------------|-------------------------|----------------------|------------------------------|--------------------------|----------------------------|--------------|
|                        |                         | American Shad        |                              | Hickory Shad             |                            |              |
|                        |                         | <u>Alosa</u><br>male | <u>sapidissima</u><br>female | <u>Pomolobus</u><br>male | <u>mediocris</u><br>female |              |
| 2/22/52                | S-44.4 - S-48.1         | 4                    | 9                            |                          | 7                          | 20           |
| 3/7/52                 | S-44.4 - S-46.4         | 12                   | 14                           | 1                        |                            | 27           |
| 3/7/52                 | S-36.5                  | 3                    | 7                            |                          | 9                          | 19           |
| 3/12/52                | S-46.4                  | 1                    | 10                           |                          |                            | 11           |
| 3/12/52                | S-36.5                  | 13                   | 7                            |                          | 1                          | 21           |
| 3/21/52                | S-46.4                  | 1                    | 13                           |                          |                            | 14           |
| 4/10/52                | S-146.4                 |                      | 6                            | 1                        |                            | 7            |
| <hr/>                  |                         |                      |                              |                          |                            |              |
| TOTAL NO.              |                         | 34                   | 66                           | 2                        | 17                         | 119          |
| Average Total L. in mm |                         | 480                  | 540                          | 370                      | 446                        |              |
| Average Weight in gm   |                         | 1,272                | 2,010                        | --                       | 1,034                      |              |

TABLE 18

SAVANNAH RIVER STUDIES  
COMPOSITION OF HOOP NET CATCHES  
S-172.5 -- S-140.1  
December 3, 1951 - July 12, 1952

| <u>Species</u>                          | <u>Number</u> |
|-----------------------------------------|---------------|
| <u>Lepisosteus osseus osseus</u>        | 18            |
| <u>Alosa sapidissima</u>                | 1             |
| <u>Carpionodes</u> n sp.                | 9             |
| <u>Moxostoma</u> spp.                   | 34            |
| <u>Minytrema melanops</u>               | 22            |
| <u>Cyprinus carpio</u>                  | 6             |
| <u>Ictalurus lacustris punctatus</u>    | 1,160         |
| <u>Ictalurus catus</u>                  | 209           |
| <u>Ameiurus platycephalus</u>           | 124           |
| <u>Perca flavescens</u>                 | 1             |
| <u>Pomoxis annularis</u>                | 5             |
| <u>Pomoxis nigro-maculatus</u>          | 10            |
| <u>Lepomis auritus</u>                  | 1             |
| <u>Lepomis macrochirus purpurascens</u> | 1             |
| <u>Roccus saxatilis</u>                 | 6             |
| <hr/>                                   |               |
| TOTAL                                   | 1,507         |
| <hr/>                                   |               |
| Number of Net Hauls                     | 117           |
| <hr/>                                   |               |

TABLE 19

|                 | Commercial Fishing |                          |                        | Hoop Netting   |                          |                        |
|-----------------|--------------------|--------------------------|------------------------|----------------|--------------------------|------------------------|
|                 | % by<br>Number     | Average<br>Lengths<br>mm | Modal<br>Lengths<br>mm | % by<br>Number | Average<br>Lengths<br>mm | Modal<br>Lengths<br>mm |
| Channel Catfish | 85.6               | 294                      | 210                    | 77.8           | 270                      | 260                    |
| White Catfish   | 10.8               | 340                      | 340                    | 14.0           | 272                      | 250                    |
| Flat Bullhead   | 3.7                | 226                      | 200                    | 8.3            | 219                      | 200                    |

Length-frequency distributions of these fish are shown in Figures 28, 29, and 30. The differences between percentage composition, average lengths, and modal lengths of fish from commercial and hoop net catches are probably due to differences in the size selectivity of the gear and seasons of greatest fishing effort. In many cases very small fish, game fish, and fish of undesirable species, were discarded from commercial catches before they could be examined. This fact alone could account for most of the observed differences in the catch.

Two striking abnormalities were noticed among the catfish examined. A few white catfish in both the hoop net and commercial catches were observed to be in an extremely emaciated condition. A number of these fish were autopsied but no obvious cause for their condition was discovered. Occasionally channel catfish were caught in which extensive areas of the body or head were covered with bloody mucus. Although these areas appeared to be tumorous or ulcerous, close inspection revealed that the skin beneath them was bloody but unbroken. Neither of these abnormalities (extreme emaciation and mucus accumulation) occurred in large

numbers of fish and both remain unexplained.

#### Gill Netting

Opportunities for setting small-meshed gill nets were limited. The accumulation of drifting debris on nets set in current would cause them to be swept away a few hours after they were set. Gill netting was, therefore, limited to the backwater at Little Hell Landing (S-146.4). Experimental gill nets set in the winter and spring of 1952 took small numbers of American shad, redhorse suckers, longnose gar, black bass, and other fish, but they yielded little information on the composition of fish populations.

#### Seining

Swift water and the presence of underwater obstacles such as stumps and snags made seining difficult and seine collections were generally limited to shallow areas around sand bars with small seines (10' - 25' long). Seine hauls attempted with a 100' seine and a large crew proved unprofitable because of the difficulties imposed by the current and snags. Even small seines could not be operated effectively by less than three men. Thus, a systematic program of seine sampling was not attempted.

The seining that was done yielded additional information only on the occurrence of minnows and other small fish. The billfish (Scomberomorus marinus) was the only fish that was taken by no method other than seining. The most common fish taken in the seine collections were silvery minnows (Hybognathus nuchalis regius) and the spottail shiner (Notropis hudsonius saludarius).

#### ROTENONE SAMPLING

Five fish collections were made in 1952 using rotenone in the form of a 5% emulsifiable solution; two were from sloughs, and three from the river proper. The quantity of rotenone used in sloughs was determined from a rough calculation of the total volume of water in these areas. The poison was then distributed by means of a bucket pump and hose, and with buckets. Vertical mixing was accomplished by running outboard motors over the areas where the poison had been distributed. It was more difficult to make satisfactory collections from the river proper. In each case five gallons of rotenone were distributed in the form of a vertical wall of high concentration which drifted downstream over the area to be poisoned. Vertical distribution was effected in deep water (3 feet or more) by pumping the mixture through a hose which was raised and lowered as the boat moved across the stream. Fluorescein dye was mixed with the rotenone so that the movement and distribution of the poison could be traced visually. Invariably the wall of poison drifted downstream diagonally due to the more rapid flow of water toward the center of the river. The size of the area which was effectively sampled could not be estimated accurately in most instances because of the variability with which the eddy currents dispersed the rotenone. All affected fish that could be reached were collected. The area that was poisoned effectively varied for different species of fish depending upon their sensitivity to rotenone. Thus, a larger area was poisoned to obtain sensitive species, such as the gizzard shad, and a lesser area for more hardy species, such as the flat bullhead. A greater percentage

of the fish poisoned in still water were retrieved than in flowing water, since the current undoubtedly removed many of the dying fish from these areas before they could be collected.

One rotenone collection was made at each of the following stations:

1 - 5/14/52, S-133.9: A backwater behind Ring Jav Point on the South Carolina side, about one acre in extent, which was broadly open to the river downstream. There were flooded areas of trees and brush along the edges of the backwater and at the upper end. The average depth was approximately 3-1/2 feet and the maximum depth about 8 feet. There was no current.

2 - 5/14/52, S-133.8: A long narrow slough on the South Carolina side less than 1/2 acre in extent which was overgrown with cypress and had a very narrow, shallow channel opening to the river. The average depth was approximately 2-1/2 feet and the maximum depth about 5 feet. There was no current.

3 - 6/30/52, S-148.8: Catfish Hole, where the river made a sharp bend toward the Georgia side and a moderate current flowed against the bank. There was a submerged sand bar at the upper end of this bend between the hole and the main channel of the river. The average depth of the hole was about 15 feet. The current of the river proper at this point was swift. The maximum depth was about 28 feet. The rotenone was placed between the sand bar and the Georgia shore. The current formed a large eddy at the lower end of the bend which sent a back-current through the hole, providing good recirculation of the poisoned water. An area of about 3/4 of an acre was sampled.



4 - 7/1/52, S-171.6 - 171.1: A cut-off on the Georgia side of the river across from the mouth of Upper Three Runs. The mouth of the cut-off was narrow with a fast current which became moderate as the channel widened to about 100 feet. The average depth was about 3 feet and the maximum depth about 6 feet. The rotenone was applied at the upper end of the cut-off and was effective throughout its entire length and for a short distance downstream in the river. The area thus poisoned was approximately 6 acres.

5 - 7/3/52 S-130.0: Three rows of pilings extended about 125 feet into the river from the Georgia side. A sandy bottom sloped gradually from the shore to a maximum depth of 10 feet at the outer edge of the pilings. The current increased from zero along the shore to rapid at the ends of the pilings. The rotenone was placed at the upper end of the middle pilings and was effective over an area of about 1-1/2 acres.

Poisoning with rotenone is one of the least selective methods of sampling fish populations. Such sampling presented the most reliable information on the composition of the fish populations, as it revealed the presence of species not collected by other methods, and yielded additional information on many other species. The results of rotenone sampling are shown in Table 21.

Samples 1 and 2, taken in sloughs, showed differences in species composition from those taken in areas more representative of the river proper. Sample 5 was obtained at a station which had both sluggish and rapid water areas, and it contained species characteristic of each of these habitats. The sloughs contained more longnose gar (Lepisosteus

osseus), carp (Cyprinus carpio), chain pickerel (Esox niger), and redbfin pickerel (Esox americanus) than did the river. White catfish (Ictalurus catus) and flat bullheads (Ameiurus platycephalus) were absent in these still water areas, and channel catfish (Ictalurus lacustris punctatus) were scarce. The largest numbers of these three catfish were taken in samples 3 and 4 from areas of flowing water. The tadpole madtom (Schilbeodes mollis) was found in the still water areas, while another madtom (Schilbeodes leptacanthus) was taken at the fast water stations. Barratt's darter (Hololepis barratti), which prefers quiet water and bottom materials rich in organic matter, was found in the collections from sloughs, while the Johnny darter (Boleosoma nigrum) was found at the three flowing water stations. The American eel (Anguilla rostrata) was common in the sloughs and at S-130.0, where some still water was present. There were also differences in the species composition of the minnow populations in the two types of habitat. Golden shiners (Notemigonus chrysolaucus borealis) preferred the still water areas, as did the shiner (Notropis maculatus). The spottail shiner (Notropis hudsonius saldanus) was scarce in the sloughs and abundant at the other stations.

#### AGE AND GROWTH DETERMINATIONS

Length-frequency distributions were constructed for those species of which adequate numbers of individuals were obtained. From these distributions some conclusions regarding the age-composition of the population can be drawn. Fish from samples 1 and 2 taken in May showed similar length-frequency distributions and could be grouped together. The same

was true for the fish taken in late June and early July. There were noticeable length differences between the fish taken in the spring (Samples 1 and 2) and those taken in the early summer (Samples 3, 4, and 5), and except in the case of bluegills, the length-frequency distributions of fish taken at these two periods were kept separate.

The histograms illustrating the length-frequency distribution of the commercial catches and the hoop net catches cannot be compared directly with that of the rotenone samples because of the size-selectivity of these operations and because these histograms represent fish taken throughout the year rather than at one season. However, these figures do indicate that the bulk of the commercial catfish fishery is composed of fish two years old or older.

Channel catfish - A sufficient number of channel catfish was taken in Samples 3, 4, and 5 to construct a length-frequency distribution histogram. (Figure 28). A small but well-defined group consisting of five young-of-the-year fish with an average length of 26 mm. was present. Because of the very small size of the individuals, this age-group was very inadequately sampled. The remaining fish form a distinctly bimodal curve with modes at 130 and 200 mm., strongly suggesting that these represent one- and two-year-old fish respectively. The scattered distribution of the larger fish offers no clue to the sizes attained by the older age groups. This age distribution remains to be confirmed by examination of sections of the pectoral spines of these fish.

Flat bullhead - The length-frequency distribution histogram (Figure 30) of the flat bullheads from Samples 3, 4, and 5 shows the presence of

a sizeable group of young-of-the-year fish with an average length of 34 mm. The year-classes of older fish are not clearly delineated, but the mode at 120 mm. obviously represents one-year-old fish. This reveals that the commercial catch of this species is composed largely of fish at least two years old.

**Silvery Minnow** - Sufficient numbers of this species were taken in all rotenone samples to determine the growth rate from May until July. (Figure 32). The young-of-the-year fish taken in May ranged in size from 20 to 33 mm. By the end of June and beginning of July these fish had reached an average length of 39 mm. The year-old fish collected in May had an average length of 71 mm. The one fish 110 mm. long was probably two years old. By the time the second series of samples had been taken, the lengths of the year old fish had increased to an average length of 80 mm.

**Spottail shiner** - The distribution of these shiners from Samples 3, 4, and 5 (Figure 33) shows two definite year-classes and the possibility of a third. The young-of-the-year fish averaged 35 mm. in length and the year-old or older fish averaged 86 mm. Of these, the small numbers of larger fish may represent two-year-olds. \*/

**Pirate perch** - Few adults of this species were taken in the rotenone collections. The growth rate of the young-of-the-year fish that were collected from May to July was determined. (Figure 34). In

---

\*/ Only about one-half of the juvenile fish of the silvery minnow and the spottail shiner collected were measured, thus, the actual proportions of young-of-the-year fish to year-old fish were about twice as great as they appear in the histograms.

Samples 1 and 2 this year-class averaged 27 mm. If 50 mm. is considered as the upper limit of this year-class in Samples 3, 4, and 5, its average size in July was 37 mm., an average length increase of 10 mm. during this period.

Bluegill - No conclusions could be drawn regarding the age-composition or growth rate of the bluegill from the information collected in rotenone sampling, because of the extended spawning period of this species which lasts throughout the summer. The length-frequency distribution histogram of the species (Figure 35), however, does show the general size-composition of the population.

#### FISH MOVEMENT STUDIES

As discussed in the section dealing with radiological studies on fishes, it is the individual fish and the amount of radioactivity that it can acquire that is of the greatest interest from the standpoint of environmental health. If a radiological health hazard should develop in the Savannah River, it would be necessary to have information on the extent of movement of individual fish from areas where high activity might be acquired into other areas. Therefore, a marking experiment was begun on May 8, 1952, and continued through July 12, 1952, to obtain information on the extent of movement of catfish in the vicinity of the Savannah River Plant of the Atomic Energy Commission.

Fish were taken in hoop nets from three areas and were marked by clipping off their pelvic fins. Fish from the upper station (S-155.35 - S-155.15) were marked by removing the right pelvic fin; those from the middle station (S-149.7 - S-149.3) by removing both pelvic fins, and those



from the lower station (S-146.3 - S-146.1) by removing the left pelvic fin. Thus, it could be determined at which area a recaptured fish had been marked. The results of this marking experiment are shown in Table 20. In addition to recaptures taken in the course of the marking experiment, one channel catfish with a double pelvic clip was taken in a rotenone sample at Catfish Hole (S-148.8), less than one mile from the point where it was first captured.

Channel catfish were found to move both up and downstream as far as 5.6 miles. A relatively small number of white catfish were marked, and only three recaptured. They were found upstream from the point of marking. Flat bullheads gave the highest proportion of recaptures of fish marked. These were all taken at the stations where they were first captured, indicating that this species is more sedentary than the other two species of catfish.

The relatively small numbers of recaptures taken during the course of fish movement studies and from the rotenone collections made within the study area indicated that a large population of catfish was present. The study, although conducted for a limited period of time and within a relatively short stretch of river, showed that channel and white catfish did move distances of several miles.\*

#### ANNOTATED LIST OF FRESH WATER FISHES

The following is an annotated list of fishes known to occur in

---

\* A more detailed study of catfish movements in the river is being continued by Mr. Robert Humphries, a graduate student at the University of Georgia, on an A.E.C. grant.



TABLE 20 - SAVANNAH RIVER STUDIES  
RESULTS OF CATFISH MARKING EXPERIMENT  
May 8 - July 12, 1952

| Location<br>(River<br>Miles)                      | Channel Catfish      |                      |                                | White Catfish                  |                                |                                | Flat Bullhead                  |                                |                                |
|---------------------------------------------------|----------------------|----------------------|--------------------------------|--------------------------------|--------------------------------|--------------------------------|--------------------------------|--------------------------------|--------------------------------|
|                                                   | No. catfish<br>hauls | Total No. Recaptures | Total No. Recaptures<br>marked | Total No. Recaptures<br>marked | Total No. Recaptures<br>marked | Total No. Recaptures<br>marked | Total No. Recaptures<br>marked | Total No. Recaptures<br>marked | Total No. Recaptures<br>marked |
| 8-155.35-<br>3-155.15<br>Right pelvic<br>clip (R) | 32                   | 224                  | 213                            | 11                             | 15                             | 15                             | 12                             | 11                             | 1                              |
| 5-145.3-<br>5-145.3<br>Double pelvic<br>clip (D)  | 34                   | 321                  | 299                            | 3                              | 12                             | 1                              | 53                             | 41                             | 12                             |
| 5-140.3-<br>5-145.1<br>Left pelvic<br>clip (L)    | 20                   | 425                  | 375                            | 2                              | 45                             | 104                            | 42                             | 37                             | 5                              |
| Total catches<br>from 8-155.6<br>to 5-145.3       | 15                   | 559                  | 527                            | 5                              | 2                              | 8                              | 16                             |                                |                                |
| TOTAL                                             | 113                  | 1531                 | 1387                           | 17                             | 16                             | 210                            | 123                            | 89                             | 1 12 5                         |

\* Including recaptures

TABLE 21  
SAVANNAH RIVER STUDIES  
Results of Rotenone Sampling

| Sample No.                       | 1       | 2                   | 3       | 4             | 5       |
|----------------------------------|---------|---------------------|---------|---------------|---------|
| Sampling Point                   | S-133.9 | S-133.8             | S-148.8 | S-171.6-171.1 | S-180.0 |
| Date                             | 5/14/52 | 5/14/52             | 6/30/52 | 7/1/52        | 7/3/52  |
| <i>Lepisosteus osseus osseus</i> | 7       | 8                   | 1       |               | 2       |
| <i>Amia calva</i>                | 2       | 1                   |         | 1             | 1       |
| <i>Pomolobusapidissima</i> )     |         |                     |         |               | 1       |
| <i>Pomolobusmodiolus</i> )       | 2       |                     |         |               |         |
| <i>Dorosomacapedianum</i>        | 2       |                     |         | 3             | 3       |
| <i>Anguilla rostrata</i>         | 14      | 26                  |         |               | 16      |
| <i>Garrpiodes n. sp.</i>         |         |                     |         | 7             |         |
| <i>Moxostoma crassilabre</i> )   |         | )                   | )       | 8             |         |
| <i>Moxostoma collapsum</i> )     | 1       | )                   | 1 )     |               |         |
| <i>Moxostoma papillorum</i> )    |         | )                   |         | 1             |         |
| <i>Minytrema melanura</i>        | 6       |                     |         | 3             | 2       |
| <i>Erimyzon sucetta sucetta</i>  | 1       |                     |         |               | 1       |
| <i>Cyprinus carpio</i>           | 13      |                     |         | 1             |         |
| <i>Etheognathus nuchalis</i>     |         |                     |         |               |         |
| <i>regina</i>                    | 24      | / fry 409 / fry 149 |         | 415           | 384     |
| <i>Notropis callisema</i>        | 94      |                     |         |               | 1       |
| <i>Notropis leiodi</i>           |         |                     | 31      | 32            | 40      |
| <i>Notropis heterodon</i>        |         |                     |         |               |         |
| <i>salicinus</i>                 | 6       |                     | 84      | 161           | 205     |
| <i>Notropis maculatus</i>        | 44      | 54                  |         |               | 4       |
| <i>Notropis petersoni</i>        | 10      | 12                  |         | 10            | 17      |
| <i>Notemigonus chrysolaemus</i>  |         |                     |         |               |         |
| <i>basci</i>                     | 12      | 1                   |         |               | 1       |
| <i>Opsopoeodus emiliae</i>       |         |                     |         |               |         |
| <i>emiliae</i>                   | 141     | 10                  |         | 8             | 2       |
| <i>Ictalurus nebulosus</i>       |         |                     |         |               |         |
| <i>punctatus</i>                 | 12      |                     | 103     | 190           | 21      |
| <i>Ictalurus catus</i>           |         |                     | 4       | 4             | 2       |
| <i>Ameiurus platycephalus</i>    |         |                     | 36      | 70            | 17      |
| <i>Ameiurus nebulosus</i>        |         |                     |         |               |         |
| <i>marmoratus</i>                | 4       |                     |         |               | 7       |
| <i>Schilbeoides leptacanthus</i> |         |                     | 1       | 8             |         |
| <i>Schilbeoides mollis</i>       | 2       | 1                   |         |               |         |


TABLE 21 (continued)  
SAVANNAH RIVER STUDIES  
Results of Rotenone Sampling

| Sample No.                        | 1       | 2       | 3       | 4             | 5       |
|-----------------------------------|---------|---------|---------|---------------|---------|
| Sampling Point                    | S-133.9 | S-133.8 | S-148.8 | S-171.6-171.1 | S-130.0 |
| Date                              | 5/14/52 | 5/14/52 | 6/30/52 | 7/1/52        | 7/3/52  |
| <i>Chologaster cornutus</i>       |         | 3       |         |               |         |
| <i>Naos americanus</i>            | 7       | 10      | 1       |               |         |
| <i>Naos niger</i>                 | 8       | 6       |         |               |         |
| <i>Gambusia affinis holbrooki</i> | 3       |         | 9       |               |         |
| <i>Aphredoderus sayanus</i>       | 10      | 42      | 65      | 42            | 27      |
| <i>Triacetes maculatus</i>        |         | 2       |         |               |         |
| <i>Labidesthes sicculus</i>       | 7       | 3       |         | 1             | 1       |
| <i>Boleosoma nigrum</i>           |         |         | 1       | 1             | 1       |
| <i>Hololepis bairdii</i>          | 1       | 6       |         |               |         |
| <i>Micropterus salmoides</i>      | 7       | 2       | 3       | 8             | 16      |
| <i>Etheocanthus gloriosus</i>     | 2       | 3       |         |               |         |
| <i>Centrarchus macropterus</i>    | 5       |         |         | 1             |         |
| <i>Chaenobrytus coronarius</i>    | 28      | 20      | 9       | 1             | 57      |
| <i>Lepomis auritus</i>            | 17      | 3       | 1       | 3             | 27      |
| <i>Lepomis macrochirus</i>        |         |         |         |               |         |
| <i>purpurascens</i>               | 76      | 35      | 48      | 30            | 95      |
| <i>Lepomis marginatus</i>         | 4       | 3       | 2       | 1             | 8       |
| <i>Lepomis microlophus</i>        | 1       | 2       | 4       | 7             |         |
| <i>Pomoxis annularis</i>          | 31      | 2       | 3       | 6             | 32      |
| <i>Pomoxis nigro-maculatus</i>    | 35      | 1       | 1       | 15            | 9       |
| <i>Elassoma zonatum</i>           |         | 5       |         |               |         |


the fresh waters of the Savannah River:

1. Atlantic Sturgeon - Acipenser oxyrinchus. Apparently a few sturgeon still ascend the river to spawn. Fishermen reported catching small individuals in gill nets at the mouth of the Lochner Creek (S-46.4) in 1952. A specimen 6' long and weighing 95 lbs. is reported to have been taken at this locality in 1951.
2. Longnose Gar - Lepisosteus osseus osseus. This species of gar appears to be abundant and ubiquitous in the river. Although it is often caught in gill nets, it is rarely used as food and is generally considered a nuisance.
3. Florida Gar - Lepisosteus platyrhincus. A single dried specimen obtained from a fisherman at Lochner Creek has been identified as this species. It is apparently very scarce in the river.
4. Bowfin - Amia calva. This species is abundant in quiet backwaters and frequently taken in gill nets. It is seldom used for food.
5. American shad - Alosa sapidissima. The American shad ascends the river in fair numbers to spawn from January through April. Although spawning was not observed, it probably occurs in the river at the mouths of tributary streams. This commercially valuable fish is the object of a limited local fishery in the lower river.
6. Hickory Shad - Pomolobus heteroclitus. The habits of the hickory shad are similar to those of the American shad and it is regularly caught with the former species. Because of its smaller size it is taken less readily in gill nets than the more valuable American shad.
7. Clut Herring - Pomolobus aestivalis. This species ascends

HICKORY SHAD  
*Pomolobus mediocris*



AMERICAN SHAD  
*Alosa sapidissima*




TOTAL LENGTH IN MILLIMETERS

LENGTH—FREQUENCY DISTRIBUTION

From commercial shad catches examined Feb.-April, 1952

FIG. 31



Rotenone collections - May 14, June 30, July 18, 3, 1952

FIG. 32




FIG. 33




FIG. 34






FIG. 35

the river in considerable numbers in March and April. Being even smaller than the hickory shad, however, it may be taken only in smaller-meshed nets. When caught it is used only as cut bait and has no commercial value.

8. Alewife - Pomolobus pseudoharengus. Fishermen have reported a clupeid fish different from the three preceding species which ascends the river before the shad run begins. It is assumed that this is the alewife, although no specimens were collected.

9. Gizzard Shad - Dorosoma cepedianum. This shad appears to be abundant throughout the river. It has no economic value.

10. American Eel - Anguilla rostrata. The eel is a fairly abundant fish. It is sometimes taken in catfish baskets and on set lines. It is used as food and as cut bait for catfish lines.

11. Carpsucker - Carpiodes n. sp. This apparently new species of carpsucker was taken occasionally by a gill net fisherman at Little Hell Landing (S-145.4). Seven adult specimens were collected in a rotenone sample at S-171.6. Young fish, presumably of this species, were seined from sandbars in this section of the river.

12. Redhorse Sucker - Moxostoma crassilabre. \*/

13. Redhorse Sucker - M. collapsum. \*/

14. Redhorse Sucker - M. papillosum. \*/

\* Species of Moxostoma have been tentatively identified as these three species. They have been taken in hoop nets, gill nets, and rotenone samples. They are fairly abundant in the river. None of the suckers is of much value as a food fish, although Moxostoma and Minytrema are occasionally sold for food.

15. Spotted Sucker - Minytrema melanops. This sucker is fairly abundant. It is often taken in nets and baskets.
16. Lake Chubsucker - Erimyzon sucetta sucetta. This species is found in backwaters of tributary streams, sloughs, and flood plain lakes.
17. Carp - Cyprinus carpio. The carp appears occasionally in basket catches and seems to be only moderately abundant in the river. It is sometimes sold as food, but it is unimportant commercially when compared with the channel catfish.
18. Silvery Minnow - Hybognathus nuchalis regius. \*/
19. Spottail Shiner - Notropis hudsonius salicarius. \*/
20. Notropis leadsi. These minnows were frequently found in association with the following three species. It is the most abundant of the three.
21. Notropis callisema. These minnows are fairly common in quiet water areas of the river.
22. Notropis niveus. These minnows are fairly common in quiet water areas of the river.
23. Notropis maculatus. These minnows are fairly common in quiet water areas of the river.
24. Notropis patersoni. This minnow is abundant in clear tributary streams. It is found occasionally in both quiet and flowing water in the river.
25. Pugnose minnow - Opsopoeodus emiliae emiliae. This species is relatively scarce but widely distributed in the river.

---

\*/ These are the most abundant forage minnows in the river. They were frequently taken in seine and rotenone collections. H. nuchalis was the more widely dispersed of the two, being found both in quiet water areas and in fast-flowing water. N. hudsonius was found in greater numbers in fast water areas.

26. Golden Shiner - Notemigonus chrysolaemus bosci. This minnow is abundant in flood plain ponds. An occasional specimen was taken in collections made in the river proper.

27. Southern Channel Catfish - Ictalurus lacustris punctatus. This is the most important and intensively fished species in the river. It is the most abundant and valuable of the catfish in the river.

28. White Catfish - Ictalurus catus. This species is caught in baskets with the channel catfish. It is much less abundant than the preceding species.

29. Flat Bullhead - Ameiurus platycephalus. This catfish is regularly caught with the above two species. It is of little value as a food fish.

30. Yellow Bullhead - Ameiurus natalis eximius.

31. Brown Bullhead - Ameiurus nebulosus marmoratus.

The above two species are found in quiet backwaters of the river and in floodplain ponds. Neither species appears to be very abundant.

32. Eastern Madtom - Schilbeodes marginatus marginatus.

33. Madtom - Schilbeodes leptacanthus.

These two diminutive catfish have been taken in brush boxes at several stations in the river and prefer fast water.

34. Tadpole Madtom - Schilbeodes mollis. This species was found in backwater areas of the river.

35. Chain Pickerel - Esox niger. This game-fish is present in tributary streams and quiet backwater areas of the river.

36. Redfin Pickerel - Esox americanus. This smaller pickerel is found in the same habitats as above species.

37. Ricefield Fish - Chologaster cornutus. This small species which burrows in the mud and organic debris of sloughs is very difficult to collect.

It was taken only in rotenone collection 2.

38. Starhead Topminnow - Fundulus dispar lineolatus. Although this species was taken only in flood plain ponds, it no doubt strays into the river proper occasionally.

39. Mosquitofish - Gambusia affinis holbrooki. This species is very abundant in shallow quiet waters of the river and in flood plain ponds.

40. Pirate Perch - Aphredoderus sayanus sayanus. The pirate perch is abundant in tributary streams and occasionally in the river.

41. Needlefish - Strongylura marina. A single specimen of this marine species was taken from a sand bar over 165 miles from the river mouth. Other specimens were observed in Upper Three Runs.

42. Hogchoker - Trinectes maculatus. This small flatfish, although normally considered a marine species, has been found as far as 170 miles up the Savannah River. It is common in the estuarine areas of the river.

43. Brook Silversides - Labidesthes sicculus. Silversides were taken occasionally in both fast and quiet-water areas of the river.

44. Striped Mullet - Mugil cephalus. This marine species is known to ascend the Savannah River, although none were collected by us. It is an important food species in estuarine areas along the coast.

45. Yellow Perch - Perca flavescens. This food fish is found in quiet backwaters from which it is taken occasionally by anglers.

46. Johnny Darter - Boleosoma nigrum subsp. This darter is fairly common on sand bars.



TABLE 22 - SAVANNAH RIVER STUDIES  
 Estuarine Sampling - April 22-24, 1952  
 Analytical Results

| Date Predicted | Tides at<br>S-O.O Time | Description<br>of Station      | Drag<br>No. | Time  | Sample<br>Depth | Temp.<br>°F | O <sub>2</sub><br>ppm | Cl <sub>2</sub><br>ppm | %<br>Sea<br>Water |
|----------------|------------------------|--------------------------------|-------------|-------|-----------------|-------------|-----------------------|------------------------|-------------------|
| Apr. 22        | High: 0634<br>1853     | N. of N. Jetty                 | 1           | 0912  | S               | 56          | ---                   | 10,465                 | 56                |
|                | Low: 0029<br>1239      | Off Tybee Is.                  | 3           | 1009  | S               | 65          | ---                   | 10,829                 | 57                |
|                |                        |                                |             |       | 23              | 64          | ---                   | 6,410                  | 34                |
|                |                        |                                |             | 1024  | S               | --          | ---                   | 15,940                 | 84                |
|                |                        |                                |             |       | 23              | --          | ---                   | 10,404                 | 55                |
|                |                        |                                | 4           | 1044  | S               | --          | ---                   | 15,344                 | 81                |
|                |                        |                                |             |       | 9'              | 65          | ---                   | 5,123                  | 27                |
|                |                        | S. end S. Jetty                | 6           | 1206  | S               | 66          | ---                   | 15,940                 | 84                |
|                |                        |                                |             |       | 12'             | 66          | ---                   | 2,343                  | 12                |
|                |                        |                                | 7           | 1248  | S               | --          | ---                   | 5,317                  | 28                |
|                |                        |                                |             |       | 11'             | 65          | ---                   | 1,530                  | 8                 |
|                |                        | S. Channel                     | 8           | 1343  | S               | --          | ---                   | 1,663                  | 9                 |
|                |                        | Mile 3                         |             |       | 11'             | 65          | ---                   | 977                    | 5                 |
|                |                        |                                |             |       |                 |             |                       | 978                    | 5                 |
| Apr. 23        | High: 0720<br>1937     | N. Channel off<br>Cockspur Is. | 10          | 0858  | S               | 55          | 6.47                  | 6,580                  | 35                |
|                | Low: 0112<br>1323      |                                |             |       | 39'             | 65          | 7.26                  | 15,479                 | 81                |
|                |                        | N. Channel                     | 11          | 0948  | S               | 66          | 6.22                  | 2,974                  | 16                |
|                |                        | Buoys 22-24                    |             |       | 12'             | 65          | 6.60                  | 10,270                 | 54                |
|                |                        | Fields Cut                     | 12          | 1124  | S               | 65          | 5.69                  | 3,132                  | 16                |
|                |                        |                                |             |       | 14'             | 66          | 5.22                  | 4,674                  | 25                |
|                |                        | St. Augustine Cr.              | 14          | 1240  | S               | 55          | 6.78                  | 119                    | 0.6               |
|                |                        | Near Jct. Wil-                 |             |       | 15'             | 65          | 6.37                  | 206                    | 1.1               |
|                |                        | mington R.                     |             |       |                 |             |                       |                        |                   |
|                |                        | Skidway R.                     | 15          | 1402  | S               | 69          | 6.95                  | 5,499                  | 29                |
|                |                        |                                |             |       | 7'              | 69          | 6.63                  | 8,049                  | 42                |
|                |                        |                                |             | 1417  | S               | 70          | 7.10                  | 8,292                  | 44                |
|                |                        |                                |             |       | 19              | 68          | 6.54                  | 8,449                  | 44                |
|                |                        | Wilmington R.                  |             | 1455  | 3.5             | 70          | 8.03                  | 7,964                  | 42                |
|                |                        | off Cen. Ogle-                 |             |       |                 |             |                       |                        |                   |
|                |                        | thorpe Hotel                   |             |       |                 |             |                       |                        |                   |
| Apr. 24        | High: 0802<br>2012     | Wassaw Sound<br>off Wassaw Is. | 18          | (1002 | S               | 69          | 6.86                  | 8,207                  | 43                |
|                | Low: 0158<br>1406      |                                |             | (     | S               | 69          | 6.76                  | 14,738                 | 78                |
|                |                        | Off Petit Chau                 | 20          | 1107  | S               | 69          | 7.00                  | 14,143                 | 74                |
|                |                        | Island                         | 21          | 1137  | S               | --          | 6.86                  | 12,400                 | 66                |
|                |                        |                                |             |       | 20              | 69          | 6.75                  | 13,621                 | 72                |
|                |                        | Tybee RR &                     | 23          | 1253  | --              | 68          | 6.49                  | 9,364                  | 49                |
|                |                        | Lazaretta Cr.                  |             |       |                 |             |                       |                        |                   |
|                |                        | Fields Cut                     |             | (1435 | S               | 69          | 6.45                  | 7,272                  | 38                |
|                |                        |                                | 24          | (     | 10              | 68          | 6.09                  | 7,303                  | 38                |
|                |                        |                                |             | (1516 | S               | 69          | 6.56                  | 7,240                  | 38                |
|                |                        |                                |             |       | 10              | 69          | 6.42                  | 7,296                  | 38                |



TABLE 23 - SAVANNAH RIVER STUDIES - ESTUARINE COLLECTIONS APRIL 1954

| Date                              | 4/22                           |    |    |     |    |    |   |    |     | 4/23 |    |    |    |    |    |    |    |    | 4/24 |    |    |    |    |    |    |  |  | Number of<br>Collect'ns<br>in which<br>Present* |
|-----------------------------------|--------------------------------|----|----|-----|----|----|---|----|-----|------|----|----|----|----|----|----|----|----|------|----|----|----|----|----|----|--|--|-------------------------------------------------|
| Drag No.                          | 1                              | 2  | 3  | 4   | 5  | 6  | 7 | 8  | 9   | 10   | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19   | 20 | 21 | 22 | 23 | 24 |    |  |  |                                                 |
|                                   | Catches not<br>Totally Counted |    |    |     |    |    |   |    |     |      |    |    |    |    |    |    |    |    |      |    |    |    |    |    |    |  |  |                                                 |
| <i>Dasyatis sabina</i>            |                                |    |    |     | 1  |    |   |    |     | 1    |    |    |    |    |    |    |    |    |      |    |    |    |    |    | 2  |  |  |                                                 |
| <i>Lepisosteus osseus</i>         |                                |    |    |     |    |    |   |    |     |      |    | 1  |    |    |    |    |    |    |      |    |    |    |    |    | 1  |  |  |                                                 |
| <i>Brevoortia tyrannus</i>        | 26                             |    |    | 1   | 7  | 4  |   |    |     |      |    | 3  |    |    |    |    |    | 1  |      |    |    |    |    |    | 7  |  |  |                                                 |
| <i>Anchoa mitchilli</i>           | 2                              | 1  |    | 1   | 1  | 12 |   | 1  |     |      |    |    | 13 | 14 | 71 | 60 |    |    |      |    | 1  | 55 | 3  |    | 14 |  |  |                                                 |
| <i>Galeichthys felis</i>          |                                |    |    | 8   |    | 1  |   |    |     | 1    |    |    |    |    |    |    |    |    |      |    | 2  | 1  |    |    | 5  |  |  |                                                 |
| <i>Urophycis regius</i>           | 1                              | 4  |    | 7   | 8  | 1  |   |    |     | 5    |    |    |    |    |    |    |    |    |      |    | 2  |    |    |    | 7  |  |  |                                                 |
| <i>Lophopsetta aquosus</i>        |                                |    |    |     |    |    |   |    |     |      |    |    |    |    |    |    |    | 1  |      | 1  |    |    |    |    | 2  |  |  |                                                 |
| <i>Ancylorsetta quadrocellata</i> |                                |    |    |     |    |    |   |    |     |      |    |    |    |    |    |    |    | 1  |      | 1  | 1  | 3  | 2  |    | 6  |  |  |                                                 |
| <i>Paralichthys dentatus</i>      |                                |    |    |     |    |    |   |    |     |      |    |    |    |    |    |    |    |    |      |    |    |    |    |    | 3  |  |  |                                                 |
| <i>Etropus crossotus</i>          |                                |    |    |     |    |    |   |    |     |      |    |    |    |    |    |    |    | 1  |      |    |    |    |    |    | 1  |  |  |                                                 |
| <i>Trinectes maculatus</i>        |                                |    |    |     |    |    |   |    |     |      |    |    |    |    |    |    |    |    |      |    |    |    |    |    | 5  |  |  |                                                 |
| <i>Symphurus plagiatus</i>        |                                |    |    |     |    |    |   |    |     |      |    |    |    |    |    |    |    |    |      |    |    |    |    |    | 10 |  |  |                                                 |
| <i>Trichiurus lepturus</i>        | 3                              | 3  |    | 1   | 1  |    |   |    |     | 3    |    |    |    |    |    |    |    |    |      |    |    |    |    | 1  | 6  |  |  |                                                 |
| <i>Lernaeus fasciatus</i>         |                                |    |    |     |    |    |   |    |     |      |    |    |    |    |    |    |    |    |      |    |    |    |    |    | 1  |  |  |                                                 |
| <i>Bairdiella chrysura</i>        | 1                              | 1  | 13 | 3   | 5  |    |   |    |     |      |    |    |    |    |    |    |    |    |      |    |    |    |    |    | 13 |  |  |                                                 |
| <i>Stellifer lanceolatus</i>      | 4                              | 38 | 19 | 500 | 32 | 2  | 1 | 43 | 350 |      |    |    |    |    |    |    |    | 1  | 1    |    | 11 | 3  | 5  | 8  | 13 |  |  |                                                 |
| <i>Leiostomus xanthurus</i>       | 4                              | 4  | 19 | 5   | 7  |    | 1 | 1  | 3   | 3    |    |    |    |    |    |    |    | 14 |      | 33 | 9  | 18 | 4  |    | 13 |  |  |                                                 |
| <i>Micropterus undulatus</i>      |                                |    |    |     |    |    |   |    |     | 1    |    |    |    |    |    |    |    |    |      |    |    |    | 1  |    | 4  |  |  |                                                 |
| <i>Menidia menidia americana</i>  |                                |    |    |     |    |    |   |    |     |      |    |    |    |    |    |    |    |    |      |    |    |    |    |    | 3  |  |  |                                                 |
| <i>Cynoscion nebulosus</i>        | 1                              |    |    | 1   |    |    |   |    |     |      |    |    |    |    |    |    |    |    |      |    |    |    |    |    | 3  |  |  |                                                 |
| <i>Alosa marginata</i>            |                                |    |    |     |    |    |   |    |     | 2    |    |    |    |    |    |    |    |    |      |    |    |    |    |    | 3  |  |  |                                                 |
| <i>Trionotus carolinus</i>        |                                |    |    |     |    |    |   |    |     |      |    |    |    |    |    |    |    |    |      |    |    |    |    |    | 1  |  |  |                                                 |
| <i>Vomer setipinnis</i>           |                                |    |    |     |    |    |   |    |     |      |    |    |    |    |    |    |    |    |      |    |    |    |    |    | 2  |  |  |                                                 |
| <i>Peprilus paru</i>              |                                |    |    |     |    |    |   |    |     |      |    |    |    |    |    |    |    |    |      |    |    |    |    |    | 1  |  |  |                                                 |
| <i>Opsanus tau</i>                | 1                              | 2  |    | 1   | 5  |    |   |    |     |      |    |    |    |    |    |    |    | 1  | 6    |    | 1  |    |    |    | 8  |  |  |                                                 |
|                                   |                                |    |    |     |    |    |   |    |     |      |    |    |    |    |    |    | 1  |    |      |    |    |    |    |    | 1  |  |  |                                                 |
| <i>Malaclemys centrata</i>        |                                |    |    |     |    |    |   |    |     |      |    |    |    |    |    |    |    |    |      |    |    |    |    |    | 1  |  |  |                                                 |
| <i>Renilla reniformis</i>         | 1                              | 39 | 11 | 1   | 1  |    |   |    |     |      |    |    |    |    |    |    |    |    |      |    |    |    |    |    | 1  |  |  |                                                 |

TABLE 23 - SAVANNAH RIVER STUDIES - ESTUARINE COLLECTIONS APRIL 1954 (Continued)

| Date                | 4/22                           |   |   |   |   |   |   |   | 4/23 |       |       |    |       |       |    |    | 4/24 |    |    |    |    |   |   |  | Number of<br>Collect'ns<br>in which<br>Present |  |
|---------------------|--------------------------------|---|---|---|---|---|---|---|------|-------|-------|----|-------|-------|----|----|------|----|----|----|----|---|---|--|------------------------------------------------|--|
| Drag No.            | 1                              | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9    | 10-11 | 12-13 | 14 | 15-16 | 17    | 18 | 19 | 20   | 21 | 22 | 23 | 24 |   |   |  |                                                |  |
|                     | Catches not<br>Totally Counted |   |   |   |   |   |   |   |      |       |       |    |       |       |    |    |      |    |    |    |    |   |   |  |                                                |  |
| Gorgonidae          |                                |   |   |   |   |   |   |   |      |       |       |    |       | Pr.** |    |    |      |    |    |    |    | 1 | 1 |  |                                                |  |
| Ctenophora          |                                |   |   |   |   |   |   |   |      |       |       |    |       |       |    |    |      |    |    |    |    |   |   |  |                                                |  |
| Echinoiroides       |                                |   |   |   |   |   |   |   |      |       |       |    |       |       |    |    |      |    |    |    |    |   |   |  |                                                |  |
| Isopoda             |                                |   |   |   |   |   |   |   |      |       |       |    |       |       |    |    |      |    |    |    |    |   |   |  |                                                |  |
| Chloridella eupura  |                                |   |   |   |   |   |   |   |      |       |       |    |       |       |    |    |      |    |    |    |    |   |   |  |                                                |  |
| Palaeomonetes sp.   |                                |   |   |   |   |   |   |   |      |       |       |    |       |       |    |    |      |    |    |    |    |   |   |  |                                                |  |
| Crangonidae         |                                |   |   |   |   |   |   |   |      |       |       |    |       |       |    |    |      |    |    |    |    |   |   |  |                                                |  |
| Penaeus setiferus   |                                |   |   |   |   |   |   |   |      |       |       |    |       |       |    |    |      |    |    |    |    |   |   |  |                                                |  |
| Paguridae           |                                |   |   |   |   |   |   |   |      |       |       |    |       |       |    |    |      |    |    |    |    |   |   |  |                                                |  |
| Callinectes sapidus |                                |   |   |   |   |   |   |   |      |       |       |    |       |       |    |    |      |    |    |    |    |   |   |  |                                                |  |
| Ovalipes occulta    |                                |   |   |   |   |   |   |   |      |       |       |    |       |       |    |    |      |    |    |    |    |   |   |  |                                                |  |
| Loligo sp.          |                                |   |   |   |   |   |   |   |      |       |       |    |       |       |    |    |      |    |    |    |    |   |   |  |                                                |  |

\* Total number of collections - 21 since 3 collections were duplicates

\*\* = Present

Reproduced from  
best available copy.

47. Barratt's Darter - Hololepis barratti. Although this species is primarily a pond and swamp stream form, it was occasionally taken in the river.

48. Banded Pigmy Sunfish - Elassoma zonatum. This small species was found occasionally in back water areas.

49. Largemouth Black Bass - Micropterus salmoides. This widely-distributed game fish is most abundant in flood-plain ponds, sloughs, and quiet water areas of the river. It is one of the important sport species in the river.

50. Blue-spotted Sunfish - Enneacanthus gloriosus. This small sunfish was taken from sloughs along the river.

51. Little Sunfish - Enneacanthus obesus. Although this species was taken only from a flood plain pond, it undoubtedly strays into the river proper.

52. Flier - Centrarchus macropterus. This sunfish is abundant in flood plain ponds. It is occasionally found in sloughs along the river.

53. Warmouth - Chaenobryttus coronarius. \*/

54. White Crappie - Pomoxia annularis. \*/

55. Black Crappie - Pomoxia nigro-maculatus. \*/

56. Yellowbelly Sunfish - Lepomis auritus. \*/

---

\*/ These seven species of panfish are common throughout the river, and are frequently taken by sport fishermen. The bluegill is probably the most abundant of these species.

57. Bluegill - Lepomis macrochirus purpurascens. \*/
58. Florida Longear Sunfish - Lepomis marginatus. \*/
59. Redear Sunfish - Lepomis microlophus. \*/
60. Striped Bass - Morone saxatilis. This anadromous fish is sometimes caught by commercial fishermen in baskets and gill nets. It is sought by sport fishermen in clear tributaries.

#### ESTUARINE STUDIES

Incidental to the collection of materials for radiological analysis from the estuarine areas at the mouth of the Savannah River, during April 1952, observations were made on the organisms encountered. Samples were taken from a 26-foot inboard motor boat employing a small otter-type trawl. This type of net which drags along the bottom is known as a "try net" in the southeastern shrimp fishery.

A series of 24 drags of fifteen minutes each was made in the estuarine areas which might be influenced by materials carried by the Savannah River.

Surface and bottom salinities and temperatures were measured in each area sampled. Beginning with drag number 10, dissolved oxygen was measured at each area sampled. Water samples for these analyses were taken at the beginning of each drag; and in areas where the conditions appeared to vary, at the end of the drag as well. These data are presented in Table 22.

---

\*/ Ibid. foregoing page.

Representatives of the more important species collected were prepared for radiological analyses. Other organisms collected were identified in order to obtain some information of the composition of the fauna of the estuarine area. This limited sampling, at one season, with highly selective gear, could not be expected to give any comprehensive picture of the populations of fish and other animals inhabiting this area. Nevertheless, some information was obtained on the occurrence of some of the important organisms present in the Savannah River estuary. Table 23 shows the numbers and kinds of organisms collected at each station.

The star drum (Stellifer lanceolatus) was the most abundant fish taken in these collections. It is a small fish, of no commercial value. The menhaden (Brevoortia tyrannus), spot (Leiostomus xanthurus), anchovy (Anchoa mitchilli), and silver perch (Bairdilla chrysaura) were all fairly abundant in the collections. Of these, both the menhaden and the spot are of commercial value. The menhaden is used almost entirely for the manufacture of meal and oil. It supports a widespread commercial fishery along the Atlantic coast. The spot is an important food fish, being taken by both commercial and sport fishermen. Other food fish of sport and commercial value occurring less frequently in our catches were the silver squeteague (Cynoscion nothus), and the Atlantic croaker (Micropogonias undulatus).

Blue crabs (Callinectes sapidus) and shrimp (Penaeus setiferus) were the most widespread crustaceans collected. They were taken in most of the areas sampled. Both these crustaceans support extensive fisheries

along the South Atlantic Coast. Some of this fishing takes place in the estuarine areas near the mouth of the Savannah River; commercial shrimping is conducted in Callibogue and Wassaw Sounds, and off Wassaw and Tybee Islands, and crab fishing is conducted in sounds and creeks throughout the area.

In addition to the organisms collected in the trawl samples, other species of commercial importance were observed in the estuarine areas. Extensive oyster beds (Crassostrea virginica) were present on the jetties at the mouth of the river and in areas adjacent to them. These beds are not utilized by man because of pollution in this area. ✓ The striped mullet (Mugil cephalus), which is an important sport and commercial species along the southern coast, is abundant in this area but it was not taken in the trawl samples.



## CHAPTER VII

BACKGROUND RADIOACTIVITY OF WATER AND BIOTA

During this survey, 32 water, 282 fresh-water biota, and 438 estuarine biota samples were collected and shipped to Cincinnati for the assay of gross alpha and beta radioactivity. Radioactive assays were made on the dissolved and suspended solids of water samples and of the flesh and bony tissues of the biological samples.

The gross background radioactivity measurements include the isotopes of the naturally occurring uranium and thorium series, and natural beta emitting isotopes such as  $K^{40}$ . The radon group were probably excluded. Such an assay would also include artificially produced isotopes which had not decayed prior to analysis. To obtain such comparative data, 10 water samples were assayed within 2 weeks of collection, while 23 were assayed within one month, and the remaining nine within four months.

About forty per cent of the biological samples of flesh of fresh water organisms were analyzed during the first month. An additional 40 per cent were analyzed during the next two months, and the remainder in six to seven months. One half of the bone samples of fresh-water fish were analyzed within one month of collection, and the remainder after eight to ten months. The estuarine biological samples were analyzed five to six months after collection.

A resume of the maximum background radioactivity of samples collected from the Savannah River in this survey is presented in Table 30. The results show that the radioactivity density of biological tissues is

substantially greater than the radioactivity density of the water which provides a habitat for the biota.

#### RADIOACTIVITY OF WATER

Five water samples were tested in August 1951, when membrane filters were not available. The analyses were similar except that, when such filters were not available, one liter of water was filtered through a 5.5 cm. Whatman No. 5 filter paper in a small Buchner funnel. The suspended solids that were retained on the paper were dried with infra-red lamps and counted in the internal proportional counter.

The results of the alpha and beta radioactivity assays of the suspended solids and solids in solution in the water samples are presented in Table 24. The radioactivity is reported in micromicrocuries per liter plus or minus a precision of measurement equal to two standard deviations.

In discussing the radioactivity of water samples, it is convenient to refer to the radioactivity associated with the suspended solids as insoluble activity and the activity of the filtrate or dissolved solids as soluble activity.

The suspended solids alpha activity, shown in Table 24, varied from zero to a maximum of  $4.1 \pm 1.7$  micromicrocuries per liter of water. The soluble alpha activity of the sample containing the maximum suspended activity was  $0.1 \pm 1.1$   $\mu\mu\text{c/liter}$ . Thus the total alpha activity of this sample was  $4.2 \pm 2.0$   $\mu\mu\text{c/liter}$  of water. Usually there was more insoluble than soluble alpha activity. However, in six out of 31 samples, the soluble alpha activity exceeded that in the suspended solids. The maximum soluble alpha activity was  $8.4 \pm 2.1$   $\mu\mu\text{c/l.}$  compared to  $0.4$   $\mu\mu\text{c/l.}$  for the suspended solids fraction.

In those samples having a measurable alpha activity, the beta activity was appreciably higher in magnitude than the alpha activity. This is particularly true with respect to soluble activity. Thus, the maximum soluble beta activity was  $31.8 \pm 14.8$   $\mu\text{pc/l.}$  This sample had an insoluble beta activity of  $8.6 \pm 5.1$   $\mu\text{pc/l.}$  The maximum total beta activity occurred for this sample, and was  $40.4 \pm 15.6$   $\mu\text{pc/l.}$

Table 2 of NBS Handbook 52\* suggests the following provisional maximum concentrations of radioactive contaminants in water: for alpha emitters,  $10^{-7}$   $\mu\text{c/ml.}$ , or 100  $\mu\text{pc/l.}$ ; for beta or gamma emitters,  $10^{-7}$   $\mu\text{c/ml.}$ , or 100  $\mu\text{pc/l.}$  These are in addition to natural background. This source further suggests application of a safety factor as large as 10 in the design and operation of permanent installations involving large quantities of radioactivity. Hence, in terms of gross activities, addition of as little as 10  $\mu\text{pc/l.}$  of alpha or beta and gamma contamination is considered limiting.

In this survey, the median gross alpha activity observed in water was 1.0  $\mu\text{pc/l.}$ , with a maximum of 8.8  $\mu\text{pc/l.}$  The median gross beta activity in water was 6.0  $\mu\text{pc/l.}$ , with a maximum of 40.4  $\mu\text{pc/l.}$  Comparison of the suggested maximum permissible additions of radioactive material with the observed gross activities indicates clearly that the suggested standard does not permit major alteration of the natural background radioactivity levels.

\* National Bureau of Standards, Handbook 52 "Maximum Permissible Amounts of Radioisotopes in the Human Body and Maximum Permissible Concentrations in Air and Water." (1953)

#### RADIOACTIVITY OF FRESH WATER BIOTA

The biological samples consist of bone and flesh samples of large fish, whole minnows, insect larvae and algae. The samples were sorted, weighed, and stored in vials in the field laboratory. Some samples were composited in the field. The perishable samples were frozen, packed in dry ice and shipped air express to Cincinnati. The samples were kept in a frozen condition until analyzed.

The radioactivity of the flesh and bony tissue of 40 individual or composite fish samples is presented in Table 25. The radioactivity of 7 composite samples of minnows and 5 algae samples is presented in Table 26, and the radioactivity of 17 samples of brush box organisms is presented in Table 27. The statistically significant results in these Tables show that the radioactivity of all samples exclusive of fish bone was less than  $47 \pm 28$   $\mu\text{pc}$  of alpha activity and less than  $14.3 \pm 5.6$   $\mu\text{pc}$  of beta activity per gram of wet weight sample. The minnows (whole fish samples) and the aquatic insects, all of which probably contained soil, showed a significant alpha activity. The algae showed no significant amount of alpha activity, but did show a measurable beta activity. The maximum alpha activity of fish flesh was  $1.5 \pm 0.9$   $\mu\text{pc/g}$ . The radioactivity of the ash of fish bone was less than the statistical error of the determination which varied from  $\pm 2$  to  $\pm 28$   $\mu\text{pc/gram}$  for the alpha activity and  $\pm 20$  to  $\pm 260$   $\mu\text{pc/g}$ . for beta activity.

#### RADIOACTIVITY OF ESTUARINE BIOTA

In April 1952, 438 samples of crabs, shrimps, oysters and fishes were collected and shipped to Cincinnati. The flesh samples were pro-

cessed for counting in a similar manner to the fresh water samples, except that larger samples amounting to as much as 5 grams of tissue were taken for analysis to improve the counting statistics of the test, with only a slight loss in the efficiency of counting due to self-absorption.

The results of 9 composite flesh samples representing 43 specimen of Blue Crab and 3 composite shell or claw samples representing 15 specimen collected from five areas at the mouth of the Savannah River are presented in Table 28. The radioactivity of the fleshy tissue of 8 composite samples of shrimp from 40 specimen and 2 composite samples of oysters from 10 specimen are also presented in Table 28.

The results of 32 composite flesh samples of fishes representing 135 specimen are presented in Table 29. Also included in Table 29 are the results of 14 composite bone samples from 62 specimen of fishes. The bone and shell of the estuarine samples were not entirely free of muscle tissue when they were processed according to the procedure given for fresh water bony tissue. <sup>\*</sup>/

Considering the results in Tables 28 and 29 the maximum alpha activity in flesh tissue was  $0.22 \pm 0.17 \mu\text{mc/g.}$  in an oyster sample. All but one composite estuarine flesh sample had a small but significant beta activity which may be attributed to  $\text{K}^{40}$  from salt water. The beta activity varied from  $0.5 \pm 0.2 \mu\text{mc/g.}$  in an oyster sample to  $4.5 \pm 1.1 \mu\text{mc/g.}$  in a fish sample. Except for one sample the alpha activity of bony tissue was less than the precision of the assay which varied from  $\pm 2.4$  to  $\pm 15.7 \mu\text{mc/g.}$  of bone ash. Similarly the beta activity of bony

---

<sup>\*</sup>/ See Appendix "E".



tissue was less than the precision of the assay which varied from  $\pm 21$  to  $\pm 110$   $\mu\text{mc/g.}$  of bone ash.

#### MINIMUM DETECTABLE RADIOACTIVITY

Due to the statistical fluctuation of counting, the minimum detectable sample radioactivity (at a 95 per cent confidence level) is taken to be equal to twice the standard deviation of the net counting rate of the sample prior to the application of correction factors for backscatter and self-absorption. The over-all efficiency of counting in a 2 pi chamber was found to be 51 per cent for alpha activity and 73 per cent for beta activity, as determined with the use of uranium and phosphorous  $^{32}$  standards.

Suppose the instrument background for alpha counting is 0.32 cpm (counts per minute) and for beta 64 cpm, then the minimum detectable sample radioactivity would be 0.3  $\mu\text{mc}$  for alpha and 2.5  $\mu\text{mc}$  for beta radioactivity. The minimum detectable radioactivity per liter or gram depends on the size of the sample taken. The minimum detectable level of alpha and beta activity for a given sample type and size is as follows:

|                           | Sample Size | Alpha                    | Beta                      |
|---------------------------|-------------|--------------------------|---------------------------|
| <u>Water Samples</u>      |             |                          |                           |
| Suspended Solids          | 0.5 liters  | 0.6 $\mu\text{mc/liter}$ | 5.0 $\mu\text{mc/liter}$  |
| Dissolved Solids          | 0.2 liters  | 1.5 $\mu\text{mc/liter}$ | 12.5 $\mu\text{mc/liter}$ |
| <u>Biological Samples</u> |             |                          |                           |
| Flesh                     | 200 mg.     | 1.5 $\mu\text{mc/g.}$    | 12.5 $\mu\text{mc/g.}$    |
| Flesh                     | 400 mg.     | 0.75 $\mu\text{mc/g.}$   | 6.2 $\mu\text{mc/g.}$     |
| Flesh                     | 750 mg.     | 0.53 $\mu\text{mc/g.}$   | 4.3 $\mu\text{mc/g.}$     |
| Flesh                     | 2,000 mg.   | 0.15 $\mu\text{mc/g.}$   | 1.25 $\mu\text{mc/g.}$    |
| Bone Ash                  | 20 mg.      | 15 $\mu\text{mc/g.}$     | 125 $\mu\text{mc/g.}$     |
| Bone Ash                  | 40 mg.      | 7.5 $\mu\text{mc/g.}$    | 62 $\mu\text{mc/g.}$      |



TABLE 24 - SAVANNAH RIVER STUDIES  
RADIOACTIVITY OF SAVANNAH RIVER WATER SAMPLES

| Location<br>(River<br>Mile) | Date<br>Collected | Alpha Activity   |                  |                  |                  | Beta Activity    |                  |                  |                  |
|-----------------------------|-------------------|------------------|------------------|------------------|------------------|------------------|------------------|------------------|------------------|
|                             |                   | Suspended        |                  | Dissolved        |                  | Suspended        |                  | Dissolved        |                  |
|                             |                   | $\mu\text{uc/l}$ | $\mu\text{uc/l}$ | $\mu\text{uc/l}$ | $\mu\text{uc/l}$ | $\mu\text{uc/l}$ | $\mu\text{uc/l}$ | $\mu\text{uc/l}$ | $\mu\text{uc/l}$ |
| UTR*                        | 12-26-51          | 3.1              | 5.9              | -0.1             | 1.1              | 8.6              | 5.1              | 31.6             | 14.8             |
| UTR*                        | 10-23-51          | 0.8              | 0.6              | 0                | 1.3              | 5.7              | 5.0              | 2.5              | 10.3             |
| LTR*                        | 12-3-51           | 0.4              | 0.6              | 0                | 1.3              | 0                | 5.2              | 5.6              | 10.0             |
| LTR*                        | 2-6-52            | 0.7              | 0.5              | 0.0              | 1.2              | -0.4             | 4.4              | -4.9             | 9.2              |
| ST*                         | 11-7-51           | 0.2              | 0.1              | 0.2              | 0.5              | 1.5              | 0.6              | -0.8             | 5.5              |
| ST*                         | 2-6-52            | 0.9              | 0.6              | 0.4              | 1.1              | -0.7             | 4.4              | 10.5             | 9.2              |
| Fo*                         | 11-7-51           | 0.2              | 0.1              | 0.2              | 0.5              | 1.0              | 1.1              | -2.9             | 3.4              |
| Fo*                         | 1-23-52           | 0.1              | 0.4              | 0.7              | 0.8              | -2.1             | 4.8              | -3.4             | 7.4              |
| Fo*                         | 2-18-52           | 1.0              | 0.7              | -0.3             | 1.6              | 4.0              | 4.4              | 3.4              | 10.8             |
| 218.5                       | 8-13-51           | 0.2              | 0.2              | 4.4              | 1.5              | 0.0              | 3.2              | 1.0              | 16.0             |
| 218.5                       | 1-8-52            | 1.2              | 1.1              | -0.3             | 1.2              | 10.8             | 10.1             | 7.7              | 7.7              |
| 215.6                       | 8-13-51           | 0.0              | 0.2              | 0.0              | 1.1              | 0.0              | 3.2              | 12.4             | 16.0             |
| 203.3**                     | 8-14-51           | 4.5              | 0.6              | 24.7             | 4.2              | 0.0              | 3.2              | 50.6             | 17.2             |
| 202.9                       | 12-18-51          | 0.6              | 0.6              | 0.1              | 1.2              | 5.7              | 5.0              | 2.5              | 10.8             |
| 202.9                       | 2-13-52           | 0.8              | 0.7              | 0.3              | 1.6              | 3.6              | 4.3              | 3.9              | 10.5             |
| 202.9                       | 3-14-52           | 2.3              | 1.0              | 1.0              | 1.2              | 0.2              | 4.4              | 23.1             | 9.8              |
| 197                         | 12-18-51          | 0.3              | 0.6              | 0.2              | 1.2              | 1.1              | 6.3              | 5.3              | 9.7              |
| 197                         | 2-13-52           | 1.4              | 0.7              | 0.4              | 1.1              | 2.3              | 4.4              | 18.5             | 10.8             |
| 189                         | 8-2-51            | 0.4              | 0.4              | 8.4              | 1.1              | 0.0              | 3.2              | 0                | 16.0             |
| 172                         | 10-23-51          | 0.3              | 0.5              | 0.8              | 1.6              | 0                | 5.1              | 4.4              | 10.3             |
| 172                         | 12-28-51          | 3.9              | 1.5              | 1.6              | 2.6              | 6.2              | 4.1              | 6.8              | 29.0             |
| 172                         | 4-3-52            | 1.2              | 0.8              | 0.5              | 1.5              | 12.3             | 4.6              | 12.6             | 10.5             |
| 167                         | 8-2-51            | 0.3              | 0.2              | 0.4              | 1.1              | 0.0              | 3.2              | 0                | 16.0             |
| 152                         | 12-3-51           | 0                | 0.5              | 0                | 1.6              | 0                | 5.2              | 0                | 11.5             |
| 152                         | 1-9-52            | 4.1              | 1.7              | 0.1              | 1.1              | 5.2              | 8.9              | -3.7             | 7.7              |
| 152                         | 4-3-52            | 1.3              | 0.7              | 1.7              | 2.2              | 13.2             | 6.3              | 12.6             | 11.1             |
| 52                          | 12-6-51           | 0.8              | 0.6              | 0                | 1.4              | 2.9              | 5.2              | 0                | 10.9             |
| 52                          | 2-20-52           | 1.7              | 0.8              | 0.0              | 1.2              | 3.8              | 4.3              | 0.0              | 10.5             |
| 52                          | 4-9-52            | 1.4              | 0.7              | 2.5              | 2.3              | -4.6             | 6.0              | 13.2             | 11.1             |
| 11 <sup>x</sup>             | 12-6-51           | 0.4              | 0.6              | 0                | 1.3              | 0                | 5.2              | 9.6              | 8.5              |
| 11                          | 3-12-52           | 0.9              | 0.7              | 0.0              | 1.6              | 5.9              | 4.3              | -5.              | 10.              |
| 11                          | 4-9-52            | 1.4              | 0.7              | 0.2              | 1.4              | -4.2             | 6.0              | 3.7              | 11.1             |

\* UTR - Upper Three Runs, LTR - Lower Three Runs,  
ST - Steel Creek, Fo - Four-Mile Branch.

\*\* Laboratory contamination suspected.

<sup>x</sup> Excluded because high salinity caused large counting error.

Table No. 25

## The Radioactivity of Fresh Water Fish

Savannah River 1951-1952

| Specimen<br>Kind             | Numbers<br>per<br>Sample | Station<br>River<br>Mile | Month<br>Coll-<br>ected | Radioactivity of flesh (1)        |                                  | Radioactivity of bone (2)         |                                  |
|------------------------------|--------------------------|--------------------------|-------------------------|-----------------------------------|----------------------------------|-----------------------------------|----------------------------------|
|                              |                          |                          |                         | Alpha<br>pnc/g <sub>f</sub> pnc/g | Beta<br>pnc/g <sub>f</sub> pnc/g | Alpha<br>pnc/g <sub>f</sub> pnc/g | Beta<br>pnc/g <sub>f</sub> pnc/g |
| <u>Gar Family</u>            |                          |                          |                         |                                   |                                  |                                   |                                  |
| <u>Lepisosteus osseus</u>    | 3                        | 146.4                    | Jan.                    | 0.2                               | 0.4                              | -0.5                              | 2.7                              |
| "                            | 4                        | 146.4                    | Jan.                    | 0.0                               | 0.2                              | -1.3                              | 2.2                              |
| <u>Bowfin Family</u>         |                          |                          |                         |                                   |                                  |                                   |                                  |
| <u>Amia calva</u>            | 2                        | 146.4                    | Jan.                    | 0.0                               | 0.3                              | -                                 | -                                |
| <u>Henning (Siad) Family</u> |                          |                          |                         |                                   |                                  |                                   |                                  |
| <u>Alosa sapidissima</u>     | 5                        | 146.4                    | Jan.                    | 0.0                               | 0.4                              | 1.4                               | 2.5                              |
| <u>Sucker Family</u>         |                          |                          |                         |                                   |                                  |                                   |                                  |
| <u>Minytrema melanops</u>    | 3                        | 164.7                    | Jan.                    | 0.5                               | 0.4                              | -0.5                              | 5.2                              |
| "                            | 10                       | 137.5                    | Dec.                    | 0.2                               | 0.4                              | 2                                 | 25                               |
| "                            | 1                        | 136.4                    | Dec.                    | 0.3                               | 0.3                              | 6                                 | 24                               |
| "                            | 1                        | 172.5                    | Jan.                    | 0.1                               | 0.3                              | -                                 | -                                |
| "                            | 1                        | 170.3                    | Jan.                    | 0.1                               | 0.3                              | -                                 | -                                |
| "                            | 5                        | 164.7                    | Jan.                    | 0.0                               | 0.3                              | 5.8                               | 5.8                              |
| "                            | 5                        | 164.7                    | Jan.                    | 0.1                               | 0.3                              | -0.2                              | 4.8                              |
| "                            | 1                        | 154.7                    | Dec.                    | 0.1                               | 0.3                              | -                                 | -                                |
| "                            | 1                        | 137.5                    | Dec.                    | 0.1                               | 0.3                              | -5                                | 20                               |
| "                            | 1                        | 136.4                    | Dec.                    | 0.0                               | 0.3                              | -6                                | 20                               |
| "                            | 4                        |                          | Jan.                    | 0.0                               | 0.3                              | 0.0                               | 2.0                              |
| <u>Composite A</u>           |                          |                          |                         |                                   |                                  |                                   |                                  |
| <u>Catfish Family</u>        |                          |                          |                         |                                   |                                  |                                   |                                  |
| <u>Ameiurus nebulosus</u>    | 3                        | 175                      | Jan.                    | 0.1                               | 0.3                              | -5.5                              | 4.2                              |
| "                            | 1                        | 172.5                    | Jan.                    | 1.5                               | 0.9                              | -                                 | -                                |
| "                            | 1                        | 171.6                    | Jan.                    | 0.2                               | 0.2                              | 12                                | 13                               |
| "                            | 3                        | 170.3                    | Jan.                    | 0.1                               | 0.3                              | -4.1                              | 4.5                              |
| "                            | 1                        | 170.3                    | Jan.                    | -0.2                              | 0.3                              | -                                 | -                                |
| <u>Ictalurus catus</u>       |                          |                          |                         |                                   |                                  |                                   |                                  |
|                              |                          |                          |                         |                                   |                                  |                                   |                                  |

(continued)

Table No. 25 (continued)

## The Radioactivity of Fresh Water Fish

Savannah River 1951-1952

| Specimen                     | Station        | Month          | Radioactivity of flesh (1) |               | Radioactivity of bone (2) |               |
|------------------------------|----------------|----------------|----------------------------|---------------|---------------------------|---------------|
| Kind                         | River-<br>Mile | Col-<br>lected | Alpha<br>mpc/g             | Beta<br>mpc/g | Alpha<br>mpc/g            | Beta<br>mpc/g |
| Catfish Family (Cont'd)      |                |                |                            |               |                           |               |
| <i>Ictalurus lacustris</i>   | 170.3          | Jan.           | 0.1                        | 0.6           | -                         | -             |
| "                            | 171.6          | Jan.           | 0.0                        | 0.2           | -1.7                      | 2.7           |
| "                            | 172.5          | Jan.           | 0.1                        | 0.6           | -                         | -             |
| Ameiurus platycephalus       |                |                |                            |               |                           |               |
| "                            | 164.7          | Jan.           | 0.1                        | 0.3           | 0.9                       | 4.8           |
| "                            | 163.8          | Dec.           | 0.2                        | 0.3           | 27                        | 13            |
| <i>Ictalurus catus</i>       | 164.7          | Jan.           | 0.4                        | 0.6           | -3.3                      | -100          |
| <i>Ictalurus lacustris</i>   | 164.7          | Jan.           | 0.2                        | 0.3           | -2.7                      | 4.5           |
| "                            | 164.7          | Jan.           | 0.2                        | 0.3           | -2.7                      | 1.9           |
| "                            | 163.8          | Dec.           | 0.1                        | 0.4           | -1.4                      | 2.0           |
| Ameiurus platycephalus       |                |                |                            |               |                           |               |
| "                            | 154.7          | Dec.           | 0.2                        | 0.4           | 8                         | 22            |
| <i>Ictalurus catus</i>       | 154.7          | Dec.           | 0.1                        | 0.3           | 31                        | 75            |
| <i>Ictalurus lacustris</i>   | 155.8          | Dec.           | -0.1                       | 0.4           | 19                        | 3             |
| "                            | 154.7          | Jan.           | 0.4                        | 0.5           | 18                        | 80            |
| Ameiurus platycephalus       |                |                |                            |               |                           |               |
| "                            | 137.5          | Jan.           | 0.2                        | 0.4           | 19                        | 25            |
| "                            | 136.4          | Jan.           | 0.2                        | 0.4           | -11                       | -140          |
| <i>Ictalurus catus</i>       | 137.5          | Dec.           | 0.2                        | 0.3           | -3                        | 220           |
| <i>Ictalurus lacustris</i>   | 137.5          | Jan.           | -0.1                       | 0.4           | 27                        | 32            |
| "                            | 136.4          | Jan.           | 0.1                        | 0.4           | 11                        | 23            |
| Composite B                  |                | Jan.           | 0.1                        | 0.4           | -11                       | 22            |
| Sunfish (Bass) Family        |                |                |                            |               |                           |               |
| <i>Micropterus salmoides</i> | 146.4          | Jan.           | -0.1                       | 0.3           | 2.7                       | 2.7           |

NOTE: (1) Per gram of tissue. (2) Per gram of ignited bone.

TABLE NO. 20

## The Radioactivity of Fresh Water Minnows and Algae

Savannah River 1951-1952

| Specimen                    |                          | Station<br>River<br>Mile | Month<br>Col-<br>lected | Radioactivity*            |                          |
|-----------------------------|--------------------------|--------------------------|-------------------------|---------------------------|--------------------------|
| Kind                        | Numbers<br>per<br>Sample |                          |                         | Alpha<br>unc/ $\mu$ unc/g | Beta<br>unc/ $\mu$ unc/g |
| <u>Minnows</u>              |                          |                          |                         |                           |                          |
| <i>Hybognathus nuchalis</i> | 5                        | 167.7                    | Jan.                    | 0.6                       | 0.3                      |
| "                           | 5                        | 167.7                    | Jan.                    | 0.3                       | 0.3                      |
| "                           | 2                        | 167.7                    | Jan.                    | 2.6                       | 1.0                      |
| <i>Notropis audsonius</i>   | 2                        | 167.7                    | Jan.                    | 3.2                       | 1.1                      |
| "                           | 1                        | 167.7                    | Jan.                    | 0.8                       | 1.6                      |
| "                           | 5                        | 167.7                    | Jan.                    | 0.7                       | 0.4                      |
| <i>Notropis</i> sp.         | 10                       | 148.3                    | Jan.                    | -0.1                      | 0.2                      |
| <u>Algae</u>                |                          |                          |                         |                           |                          |
| <i>Vaucheria</i> sp.        |                          | 171.0                    | Jan.                    | 0.4                       | 1.3                      |
| <i>Stigeoclonium</i> sp.    |                          | 155.4                    | Jan.                    | -0.5                      | 0.6                      |
| <i>Vaucheria</i> sp.        |                          | 155.4                    | Jan.                    | -0.1                      | 0.7                      |
| <i>Audouinella</i> sp.      |                          | 155.4                    | Jan.                    | -0.4                      | 0.6                      |
| Algae                       |                          | 148.8                    | Jan.                    | 3.4                       | 3.6                      |
|                             |                          |                          |                         | 7.9                       | 15.3                     |
|                             |                          |                          |                         | 13.4                      | 5.4                      |
|                             |                          |                          |                         | 0.8                       | 8.2                      |
|                             |                          |                          |                         | 14.3                      | 5.6                      |
|                             |                          |                          |                         | -9.3                      | 27.1                     |

Note. \* Per gram of whole fish or moist (blotted) sample.

TABLE NO. 27

## The Radioactivity of Fresh Water Brush Box Organisms

Savannah River 1951-1952

| Specimen                      |         |                 | Radioactivity       |                    |            |
|-------------------------------|---------|-----------------|---------------------|--------------------|------------|
| Kind                          | Station | Month Collected | Alpha<br>mc/g-unc/g | Beta<br>mc/g-unc/g |            |
| Annelida (Aquatic earthworms) | Fo      | Dec.            | 6.4                 | 11.2               | -25 109    |
| Coleoptera (Beetles)          | UTR     | Dec.            | 47                  | 28                 | 156 217    |
| Diptera (Two-winged flies)    | Fo      | Dec.            | 0.0                 | 20                 | -129 187   |
| "                             | UTR     | Dec.            | 1.9                 | 3.8                | 6.7 37     |
| "                             | UTR     | Nov.            | 3.0                 | 12.0               | 53 134     |
| Ephemera (Mayflies)           | Fo      | Dec.            | -0.4                | 2.5                | -8.9 30.6  |
| "                             | UTR     | Dec.            | 1.3                 | 1.6                | 4.7 13.9   |
| "                             | UTR     | Nov.            | 1.3                 | 1.7                | 7.3 15.8   |
| Neuroptera (Lace-wings)       | UTR     | Dec.            | 1.6                 | 0.9                | 2.6 5.7    |
| "                             | 163.8   | Dec.            | 0.5                 | 0.5                | 0.5 4.0    |
| Odonata (Dragonflies)         | UTR     | Nov.            | 2.6                 | 2.1                | 1.3 17.2   |
| Plecoptera (Stoneflies)       | UTR     | Dec.            | 0.3                 | 0.7                | -5 5.4     |
| "                             | 163.8   | Dec.            | 0.2                 | 2.5                | 3.1 30     |
| Trichoptera (Caddis flies)    | Fo      | Dec.            | 3.9                 | 4.7                | -23.1 39.2 |
| "                             | UTR     | Dec.            | 4.8                 | 3.9                | 11.5 32    |
| "                             | 163.8   | Dec.            | 0.3                 | 0.6                | 7.9 5.5    |
| "                             | UTR     | Nov.            | 3.1                 | 2.4                | 7.8 19.0   |

Note: \* Four Mile Creek (Fo), Upper Three Runs (UTR).

\*\* Per gram of moist (blotted) sample.

TABLE 28 - The Radioactivity of Crabs, Oysters and Shrimp  
Found in Savannah River Estuaries April 1952

| Specimen<br>King                    | Numbers<br>per<br>Sample | Station* | Radioactivity of Flesh (1)              |                                        | Radioactivity of Shell (2)              |                                        |
|-------------------------------------|--------------------------|----------|-----------------------------------------|----------------------------------------|-----------------------------------------|----------------------------------------|
|                                     |                          |          | Alpha<br>$\mu\text{mc/g}\mu\text{mc/g}$ | Beta<br>$\mu\text{mc/g}\mu\text{mc/g}$ | Alpha<br>$\mu\text{mc/g}\mu\text{mc/g}$ | Beta<br>$\mu\text{mc/g}\mu\text{mc/g}$ |
| Crustaceae                          |                          |          |                                         |                                        |                                         |                                        |
| Callinectes sapidus,<br>(Blue crab) | 5                        | A        | 0.04                                    | 0.06                                   | 0.5                                     | 2.9                                    |
| " "                                 | 5                        | A        | 0.04                                    | 0.06                                   | 2.2                                     | 3.2                                    |
| " "                                 | 4                        | B        | -0.04                                   | 0.15                                   |                                         |                                        |
| " "                                 | 5                        | B        | -0.11                                   | 0.11                                   |                                         |                                        |
| " "                                 | 4                        | B        | 0.02                                    | 0.13                                   |                                         |                                        |
| " "                                 | 5                        | C-D      | 0.11                                    | 0.16                                   | 1.0                                     | 5.7                                    |
| " "                                 | 5                        | D        | 0.01                                    | 0.14                                   |                                         |                                        |
| " "                                 | 5                        | E        | -0.11                                   | 0.12                                   |                                         |                                        |
| " "                                 | 5                        | E        | -0.02                                   | 0.15                                   |                                         |                                        |
| Penaeo setiferus,<br>(Shrimp)       | 5                        | A        | 0.03                                    | 0.07                                   | 1.6                                     | 0.5                                    |
| " "                                 | 5                        | A        | 0.03                                    | 0.07                                   | 1.6                                     | 0.5                                    |
| " "                                 | 5                        | A        | 0.00                                    | 0.07                                   | 1.8                                     | 0.5                                    |
| " "                                 | 5                        | B        | -0.02                                   | 0.08                                   | 2.3                                     | 0.5                                    |
| " "                                 | 5                        | B        | -0.02                                   | 0.08                                   | 2.4                                     | 0.6                                    |
| " "                                 | 5                        | C        | 0.09                                    | 0.05                                   | 2.2                                     | 0.5                                    |
| " "                                 | 5                        | C        | 0.04                                    | 0.05                                   | 2.6                                     | 0.5                                    |
| " "                                 | 5                        | D        | 0.00                                    | 0.07                                   | 1.8                                     | 0.5                                    |
| Mollusca                            |                          |          |                                         |                                        |                                         |                                        |
| Ostrea virginica,<br>(Oyster)       | 5                        | A        | 0.19                                    | 0.06                                   | 0.9                                     | 0.2                                    |
| " "                                 | 5                        | A        | 0.13                                    | 0.05                                   | 0.9                                     | 0.2                                    |

Note: \* Mouth of Savannah River off Savannah Beach (A); Off Long Island at midlength (B); at Fields

Cut (C), near mouth of Wilmington and Skidaway Rivers (D), Wassaw Sound off Cabbage Island (E).

(1) Per gram of flesh.

(2) Per gram of ignited shell or claw.



TABLE NO. 29

## The Radioactivity of Fish Found in the

Savannah River Estuaries April 22-24, 1952

| Specimen<br>Kind                                 | Numbers<br>per<br>Sample | Station* | Radioactivity of Flesh(1)  |                           | Radioactivity of Bone(2)   |                           |
|--------------------------------------------------|--------------------------|----------|----------------------------|---------------------------|----------------------------|---------------------------|
|                                                  |                          |          | Alpha<br>unc/g $\pm$ unc/g | Beta<br>unc/g $\pm$ unc/g | Alpha<br>unc/g $\pm$ unc/g | Beta<br>unc/g $\pm$ unc/g |
| <i>Phycis regius</i> (Hake)                      | 3                        | A        | -0.03                      | 0.06                      | -4.4                       | 15.7                      |
|                                                  | 3                        | A        | 0.02                       | 0.14                      | -2.8                       | 4.4                       |
|                                                  | 3                        | B        | 0.00                       | 0.10                      | 1.8                        | 5.7                       |
| <i>Brevoortia smithi</i><br>(Menhaden)           | 5                        | A        | 0.22                       | 0.17                      | -3.1                       | 2.4                       |
|                                                  | 5                        | A        | -0.01                      | 0.09                      | -1.1                       | 2.7                       |
|                                                  | 5                        | A/D      | 0.08                       | 0.07                      | -                          | -                         |
|                                                  | 5                        | C/E      | 0.01                       | 0.05                      | -                          | -                         |
| <i>Bairdella chrysura</i> ,<br>(Silver perch)    | 5                        | A        | -0.06                      | 0.10                      | -                          | -                         |
|                                                  | 4                        | C        | 0.04                       | 0.08                      | -                          | -                         |
|                                                  | 5                        | C        | 0.06                       | 0.07                      | -                          | -                         |
|                                                  | 5                        | D        | -0.02                      | 0.06                      | -                          | -                         |
|                                                  | 4                        | D        | -0.03                      | 0.08                      | -                          | -                         |
|                                                  | 5                        | E        | 0.03                       | 0.05                      | 1.7                        | 3.9                       |
|                                                  | 5                        | E        | 0.05                       | 0.05                      | 0.3                        | 2.5                       |
| <i>Leiostomus xanthurus</i><br>(Spotted croaker) | 5                        | A        | 0.03                       | 0.07                      | -0.3                       | 2.5                       |
|                                                  | 5                        | A        | 0.01                       | 0.07                      | -0.3                       | 2.5                       |
|                                                  | 5                        | E        | 0.09                       | 0.07                      | -                          | -                         |
|                                                  | 4                        | D        | 0.03                       | 0.10                      | -                          | -                         |
|                                                  | 4                        | D        | 0.03                       | 0.10                      | -                          | -                         |
|                                                  | 5                        | E        | -0.06                      | 0.07                      | 4.1                        | 3.4                       |
|                                                  | 5                        | E        | -0.07                      | 0.07                      | 0.0                        | 2.7                       |
|                                                  | 5                        | E        | -                          | -                         | -                          | -                         |

(Continued)

TABLE NO. 25 (Cont)

## The Radioactivity of Fish Found in the

Savannah River Estuaries April 22-24, 1952

| Specimen Kind                              | Numbers per Sample | Station* | Radioactivity of Flesh(1)     |                              | Radioactivity of Bone(2)      |                              |
|--------------------------------------------|--------------------|----------|-------------------------------|------------------------------|-------------------------------|------------------------------|
|                                            |                    |          | Alpha<br>muc/ $\frac{1}{2}$ g | Beta<br>muc/ $\frac{1}{2}$ g | Alpha<br>muc/ $\frac{1}{2}$ g | Beta<br>muc/ $\frac{1}{2}$ g |
| Menticirrhus americanus,<br>(King croaker) | 4                  | A        | 0.06                          | 2.8                          | -2.0                          | -57                          |
| Micropogon undulatus,<br>(croaker)         | 3                  | D        | -0.01                         | 1.7                          | -                             | -                            |
|                                            | 3                  | D        | 0.00                          | 1.8                          | -                             | -                            |
| Peprilus paru,<br>(Harvest fish)           | 5                  | A        | 0.22                          | 2.3                          | 1.2                           | 3.2                          |
|                                            | 4                  | A        | 0.23                          | 2.9                          | 2.2                           | -12.2                        |
|                                            | 3                  | A/D      | 0.13                          | 2.9                          | -                             | -12.5                        |
|                                            | 4                  | B        | 0.05                          | 3.0                          | -                             | -                            |
|                                            | 4                  | B        | 0.06                          | 2.0                          | -                             | -                            |
|                                            | 4                  | D        | 0.06                          | 1.7                          | -                             | -                            |
| Cynoscion regalis,<br>(Weak fish)          | 4                  | D        | 0.10                          | 1.5                          | -                             | -                            |
|                                            | 1                  | E        | 0.04                          | 2.2                          | -                             | -                            |

Note: \* Mouth of Savannah River off Savannah Beach (A), Off Long Island at midlength (B), at Fields Cut(C), near mouth of Wilmington and Skidaway Rivers (D), Wassaw Sound off Cabbage Island (E).

(1) Per gram of flesh.

(2) Per gram of ignited bone.

TABLE 30  
The Maximum Radioactivity Density  
of the Savannah River Samples

1952-1953

| Sample type                  | Alpha Radioactivity |        | Beta Radioactivity |        |
|------------------------------|---------------------|--------|--------------------|--------|
|                              | umc/g $\pm$ umc/g   |        | umc/g $\pm$ umc/g  |        |
| <u>Fresh Water</u>           |                     |        |                    |        |
| In suspended solids          | 0.0041*             | 0.0017 | 0.0132*            | 0.0006 |
| In dissolved solids          | 0.0084*             | 0.0021 | 0.0318*            | 0.0148 |
| <u>Fresh Water Organisms</u> |                     |        |                    |        |
| Fish Flesh                   | 1.5                 | 0.9    | 11.6               | 4.5    |
| Minnows (whole fish)         | 3.2                 | 1.1    | 8.7                | 5.0    |
| Aquatic Insects              | 47.                 | 28.    | 8.6                | 5.7    |
| Algae                        | 3.4                 | 3.6    | 14.3               | 5.6    |
| <u>Estuarine Organisms</u>   |                     |        |                    |        |
| Crab flesh                   | 0.11                | 0.16   | 3.0                | 0.9    |
| Shrimp flesh                 | 0.09                | 0.06   | 2.6                | 0.5    |
| Oyster flesh                 | 0.19                | 0.06   | 0.9                | 0.2    |
| Fish flesh                   | 0.22                | 0.17   | 4.5                | 1.1    |

Note: \*Per milliliter of water

APPENDIX "A"

SUMMARY OF INDIVIDUAL RESULTS - SANITARY EXAMINATIONS

**SAVANNAH RIVER STUDIES**  
**Summary of Individual Results - Sanitary Examinations**

| Sampling Point    | Date<br>1951 | Disch.<br>at<br>Augusta<br>Sec-Ft | O.C.<br>Moore<br>ppm | 5-Day<br>20°C<br>BOD<br>ppm | Temp.<br>°C | Turb.<br>ppm | Color<br>ppm | Dissolved<br>Oxygen |       | pH  | Alkal.<br>ppm | Chloride<br>ppm | Coliforms<br>MPN per<br>100 ml |        | S.O. |
|-------------------|--------------|-----------------------------------|----------------------|-----------------------------|-------------|--------------|--------------|---------------------|-------|-----|---------------|-----------------|--------------------------------|--------|------|
|                   |              |                                   |                      |                             |             |              |              | ppm                 | % Sat |     |               |                 | Ga.                            | 100 ml |      |
| S 230.7           | M.S. 8/9     | 2,960                             | 6.0                  | 0.25                        | ---         | 70           | 15           | 7.0                 | ---   | 7.1 | 2             | 4.1             | 230                            |        |      |
|                   | M.S. 8/14    | 3,800                             | 4.1                  | 0.31                        | 30          | 42           | 15           | 7.0                 | 92    | 7.3 | 19            | 3.2             | 930                            |        |      |
|                   | M.S. 8/23    | 2,790                             | 5.0                  | 0.62                        | 30          | 80           | 20           | 7.0                 | 91    | 7.3 | 17            | 3.3             | 230                            |        |      |
|                   | M.S. 9/18    | 4,100                             | 6.5                  | 0.31                        | 23          | 93           | 20           | 8.0                 | 92    | 7.6 | 19            | 3.6             | 2,300                          |        |      |
| *S 223.6; Agc 3.4 | 8/9          |                                   | 2.9                  | 0.23                        | ---         | 66           | 50           | 6.8                 | ---   | 7.2 | 17            | 3.0             | 210                            |        |      |
|                   | 8/14         |                                   | 4.4                  | 0.49                        | 30          | 37           | 10           | 7.1                 | 93    | 7.2 | 17            | 3.2             | 930                            |        |      |
|                   | 8/23         |                                   | 4.9                  | 0.51                        | 29          | 85           | 20           | 6.9                 | 88    | 7.3 | 17            | 3.2             | 1,500                          |        |      |
|                   | 9/18         |                                   | 5.1                  | 0.41                        | 24          | 86           | 20           | 7.4                 | 87    | 7.5 | 18            | 3.4             | 4,300                          |        |      |
| S 218.5           | Ga. 7/30     | 4,800                             | 8.4                  | 0.43                        | ---         | 48           | 40           | 7.2                 | ---   | 7.4 | 19            | 2.2             | 750                            |        |      |
|                   | M.S. 7/30    | 4,800                             | 6.7                  | 0.47                        | ---         | 37           | 32           | 7.4                 | ---   | 7.5 | 20            | 2.8             | 230                            |        |      |
|                   | S.O. 7/30    | 4,800                             | 8.3                  | 0.19                        | ---         | 36           | 28           | 7.3                 | ---   | 7.5 | 19            | 2.7             | 430                            |        |      |
|                   | Ga. 8/13     | 3,580                             | 5.1                  | 0.26                        | 30          | 36           | 10           | 6.8                 | 89    | 7.4 | 19            | 2.4             | 91                             |        |      |
|                   | M.S. 8/13    | 3,580                             | 4.4                  | 0.37                        | 30          | 34           |              | 7.0                 | 91    | 7.4 | 19            | 2.8             | 360                            |        |      |
|                   | S.O. 8/13    | 3,580                             | ---                  | 0.43                        | 30          | 32           |              | 7.0                 | 91    | 7.5 | 18            | 2.7             | 2,400                          |        |      |
|                   | M.S. 8/22    | 3,470                             | 5.4                  | 0.20                        | 30          | 65           | 20           | 6.7                 | 88    | 7.4 | 17            | 3.2             | 430                            |        |      |
|                   | M.S. 9/11    | 2,800                             | 4.2                  | 0.44                        | 27          | 27           | 10           | 7.3                 | 90    | 7.6 | 18            | 2.9             | 2,300                          |        |      |
|                   | M.S. 9/24    | 4,720                             | 1.6                  | 0.48                        | 25          | 48           | 20           | 8.0                 | 95    | 7.5 | 20            | 3.5             | 930                            |        |      |
|                   | M.S. 10/8    | 2,560                             | 7.4                  | 0.27                        | 25          | 83           | 28           | 7.6                 | 90    | 7.4 | 17            | 4.2             | 2,300                          |        |      |
|                   | M.S. 10/15   | 2,360                             | 3.0                  | 0.26                        | 19          | 37           | 33           | 8.8                 | 94    | 7.5 | 20            | 3.5             | 91                             |        |      |
|                   | M.S. 10/24   | 1,960                             | 4.0                  | 0.22                        | 21          | 33           | 28           | 8.5                 | 94    | 7.5 | 19            | 3.8             | 93                             |        |      |
|                   | M.S. 10/29   | 3,840                             | 4.2                  | 0.37                        | 19          | 28           | 17           | 8.6                 | 92    | 7.5 | 19            | 3.9             | 930                            |        |      |
|                   | M.S. 12/4    | 4,840                             | 2.9                  | 0.31                        | 11          | 8            | 28           | 11.0                | 99    | 6.9 | 16            | 3.9             | 430                            |        |      |
|                   | M.S. 12/10   | 5,460                             | 9.5                  | 0.55                        | 14          | 180          | 140          | 10.2                | 98    | 7.1 | 13            | 3.6             | 2,400                          |        |      |
|                   | M.S. 12/19   | 9,990                             | 6.1                  | 0.37                        | 6           | 65           | 56           | 12.4                | 99    | 7.1 | 15            | 3.4             | 4,600                          |        |      |
|                   | M.S. 12/26   | 14,300                            | 20                   | 0.85                        | 7           | 310          | 136          | 12.5                | 103   | 6.7 | 7             | 2.6             | 24,000                         |        |      |

Preceding page blank

\* By waterworks intake in Augusta Canal. This is 3.4 miles downstream from source of Canal at S 223.6





**SAVANNAH RIVER STUDIES**  
**Summary of Individual Results - Sanitary Examinations**

| Sampling Point                  | Date         | Disch.<br>at<br>Augusta<br>Sec-Ft | O.C.<br>Moore<br>ppm | 5-Day<br>20°C<br>BOD<br>ppm | Temp.<br>°C | Turb.<br>ppm | Color<br>ppm | Dissolved<br>Oxygen |       | pH  | Alkal.<br>ppm | Chloride<br>ppm | Coliforms<br>MPN per<br>Ga. 100 ml | S.C.   |
|---------------------------------|--------------|-----------------------------------|----------------------|-----------------------------|-------------|--------------|--------------|---------------------|-------|-----|---------------|-----------------|------------------------------------|--------|
|                                 |              |                                   |                      |                             |             |              |              | ppm                 | % Sat |     |               |                 |                                    |        |
| S 217.2;<br>Hc 0.01<br>(Cont'd) | 1951<br>9/24 | M.S.                              | 6.0                  | 1.0                         | 24          | 81           | 20           | 7.8                 | 91    | 7.7 | 21            | 3.4             | 93,000                             |        |
|                                 | 10/15        | M.S.                              | 5.2                  | 1.2                         | 19          | 40           | 31           | 8.7                 | 81    | 7.4 | 19            | 3.8             | 430,000                            |        |
|                                 | 12/10        | M.S.                              | 15                   | 0.87                        | 13          | 190          | 108          | 10.1                | 95    | 6.8 | 11            | 5.0             | 93,000                             |        |
|                                 | 1952<br>1/8  | M.S.                              | 9.4                  | 1.2                         | 8           | 73           | 104          | 11.6                | 98    | 7.0 | 9             | 3.3             | 24,000                             |        |
| S 215.6                         | 1951<br>7/30 | Ga.                               | 4.2                  | 0.44                        | --          | 60           | 32           | 6.9                 | --    | 7.2 | 20            | 1.7             | 24,000                             |        |
|                                 | 7/30         | M.S.                              | 5.5                  | 0.35                        | --          | 60           | 37           | 6.9                 | --    | 7.3 | 20            |                 | 46,000                             |        |
|                                 | 7/30         | S.C.                              | 7.6                  | 0.40                        | --          | 60           | 38           | 6.9                 | --    | 7.3 | 20            |                 | 46,000                             |        |
|                                 | 8/13         | Ga.                               | 3.8                  | 0.46                        | 31          | 36           | 12           | 6.6                 | 88    | 7.4 | 18            | 2.2             | 43,000                             | 46,000 |
|                                 | 8/13         | M.S.                              | 3.4                  | 0.56                        | 31          | 37           | 12           | 6.6                 | 88    | 7.4 | 19            | 2.0             | 93,000                             |        |
|                                 | 8/13         | S.C.                              | 3.0                  | 0.54                        | 31          | 37           | 12           | 6.6                 | 88    | 7.4 | 19            | 2.2             | 43,000                             |        |
|                                 | 8/22         | Ga.                               | 5.5                  | 0.07                        | 30          | 80           | 20           | 6.3                 | 84    | 7.3 | 16            | 3.0             | 43,000                             | 43,000 |
|                                 | 8/22         | M.S.                              | 6.2                  | 0.44                        | 30          | 80           | 20           | 6.7                 | 88    | 7.3 | 15            | 3.2             | 93,000                             |        |
|                                 | 8/22         | S.C.                              | 5.0                  | 0.16                        | 30          | 83           | 20           | 6.4                 | 84    | 7.3 | 14            | 2.7             | 43,000                             |        |
|                                 | 9/11         | M.S.                              | 4.5                  | 0.73                        | 27          | 27           | 10           | 7.2                 | 90    | 7.6 | 18            | 3.2             | 240,000                            | 23,000 |
|                                 | 9/24         | M.S.                              | 0.83                 | 0.77                        | 24          | 55           | 25           | 7.8                 | 91    | 7.6 | 19            | 3.8             | 23,000                             | 43,000 |
|                                 | 8/9          | M.S.                              | 42                   | 11                          | --          | 39           | 110          | 0                   | --    | 8.3 | 100           | 5.5             | 43,000                             |        |
|                                 | 8/15         | M.S.                              | 31                   | 11                          | 28          | 32           | 80           | 2.5                 | 32    | 8.5 | 50            | 5.4             | 430,000                            |        |
| S 213.36;<br>Hc 0.01            | 8/22         | M.S.                              | 41                   | >23                         | 28          | 25           | 80           | 0                   | 0     | 8.5 | 41            | 8.3             | 430,000                            |        |
|                                 | 9/11         | M.S.                              | 27                   | 6.5                         | 25          | <25          | 70           | 0.16                | 2     | 7.8 | 69            | 5.5             | 430,000                            |        |
|                                 | 9/26         | M.S.                              | 20                   | 7.4                         | 22          | 7            | 80           | 2.9                 | 33    | 7.8 | 57            | 5.3             | 43,000                             |        |
|                                 | 10/8         | M.S.                              | 20                   | 2.7                         | 22          | 7            | 60           | 3.4                 | 38    | 7.7 | 60            | 5.5             | 240,000                            |        |
|                                 | 12/4         | M.S.                              | 44                   | 14                          | 14          | 4            | 58           | 2.5                 | 24    | 7.7 | 61            | 5.5             | 9,300                              |        |
|                                 | 1952<br>1/2  | M.S.                              | 32                   | 8.9                         | 14          | 4            | 72           | 2.8                 | 27    | 7.5 | 44            | 4.5             | 230                                |        |
|                                 | 2/11         | M.S.                              | 16                   | 4.5                         | 13          | 2            | 50           | 4.2                 | 40    | 7.8 | 45            | 5.1             | 5,000                              |        |
|                                 | 3/5          | M.S.                              | 15                   | 4.0                         | 12          | 93           | 80           | 6.3                 | 58    | 7.2 | 13            | 3.4             | 7,500                              |        |

**SAVANNAH RIVER STUDIES**  
**Summary of Individual Results - Sanitary Examinations**

| Sampling Point | Date  | Disch.<br>at<br>Augusta<br>Sec-Ft | O.C.<br>Moore<br>ppm | 5-Day<br>20°C<br>BOD<br>ppm | Temp.<br>°C | Turb.<br>ppm | Color<br>ppm | Dissolved<br>Oxygen |       | pH  | Alkal.<br>ppm | Chloride<br>ppm | Coliforms<br>MPN per<br>Ga. 100 ml S.C. |         |
|----------------|-------|-----------------------------------|----------------------|-----------------------------|-------------|--------------|--------------|---------------------|-------|-----|---------------|-----------------|-----------------------------------------|---------|
|                |       |                                   |                      |                             |             |              |              | ppm                 | % Sat |     |               |                 |                                         |         |
|                |       |                                   |                      |                             |             |              |              |                     |       |     |               |                 |                                         |         |
| 1951           |       |                                   |                      |                             |             |              |              |                     |       |     |               |                 |                                         |         |
| S 212.0        | 8/13  | 3,580                             | 3.2                  | 0.47                        | 31          | 36           | 12           | 6.4                 | 86    | 7.2 | 18            | 2.5             | 93,000                                  |         |
| Ga.            | 8/13  | 3,580                             | 2.9                  | 0.54                        | 31          | 37           |              | 6.3                 | 84    | 7.3 | 20            | 2.4             | 43,000                                  |         |
| M.S.           | 8/13  | 3,580                             | 4.2                  | 0.70                        | 31          | 38           |              | 6.4                 | 86    | 7.3 | 20            | 2.6             |                                         | 43,000  |
| S.C.           | 8/22  | 3,470                             | 5.3                  | 0.64                        | 30          | 80           | 25           | 6.1                 | 80    | 7.4 | 15            | 3.3             | 150,000                                 |         |
| Ga.            | 8/22  | 3,470                             | 5.4                  | 0.96                        | 30          | 82           |              | 6.1                 | 80    | 7.3 | 16            | 3.2             | 75,000                                  |         |
| M.S.           | 8/22  | 3,470                             | 5.6                  | 0.61                        | 30          | 81           |              | 6.1                 | 80    | 7.3 | 15            | 3.3             |                                         | 93,000  |
| S.O.           | 9/12  | 2,080                             | 5.0                  | 1.1                         | 28          | <25          | 15           | 5.5                 | 70    | 7.7 | 23            | 4.0             | 240,000                                 |         |
| Ga.            | 9/12  | 2,080                             | 4.0                  | 1.1                         | 28          | <25          |              | 6.0                 | 76    | 7.7 | 21            | 3.9             | 460,000                                 |         |
| M.S.           | 9/12  | 2,080                             | 3.6                  | 1.2                         | 28          | <25          |              | 6.1                 | 77    | 7.7 | 21            | 3.4             |                                         | 460,000 |
| S.O.           | 9/26  | 6,030                             | 6.3                  | 0.83                        | 27          | 85           | 43           | 6.8                 | 84    | 7.5 | 19            | 3.9             | 230,000                                 |         |
| Ga.            | 9/26  | 6,030                             | 7.6                  | 1.1                         | 26          | 83           |              | 7.0                 | 85    | 7.5 | 20            | 4.2             | 23,000                                  |         |
| M.S.           | 9/26  | 6,030                             | 6.1                  | 1.1                         | 26          | 67           |              | 6.8                 | 83    | 7.5 | 24            | 3.6             |                                         | 23,000  |
| S.C.           | 10/8  | 2,560                             | 10                   | 0.49                        | 26          | 78           | 25           | 6.9                 | 84    | 7.3 | 20            | 4.5             | 93,999                                  | 93,000  |
| M.S.           | 10/15 | 2,360                             | 6.1                  | 0.51                        | 19          | 32           |              | 8.0                 | 86    | 7.5 | 22            | 10              | 43,000                                  | 93,000  |
| M.S.           | 10/24 | 1,960                             | 6.9                  | 1.2                         | 19          | 28           |              | 7.1                 | 76    | 7.4 | 24            | 4.4             | 150,000                                 | 240,000 |
| M.S.           | 10/29 | 3,840                             | 5.6                  | 0.57                        | 19          | 25           | 26           | 7.7                 | 83    | 7.5 | 23            | 3.8             | 24,000                                  | 9,300   |
| M.S.           | 12/4  | 4,840                             | 7.5                  | 0.16                        | 11          | 11           |              | 10.8                | 97    | 7.0 | 15            | 4.2             | 43,000                                  | 24,000  |
| M.S.           | 12/10 | 5,460                             | 10                   | 0.66                        | 14          | 180          |              | 9.6                 | 93    | 7.0 | 13            | 3.6             | 43,000                                  | 43,000  |
| M.S.           | 12/19 | 9,990                             | 5.4                  | 0.50                        | 6           | 62           | 50           | 12.3                | 99    | 7.0 | 13            | 3.3             | 43,000                                  | 9,300   |
| M.S.           | 12/26 | 14,300                            | 18                   | 1.0                         | 7           | 330          |              | 12.4                | 102   | 7.0 | 8             | 3.0             | 24,000                                  | 4,300   |
| M.S.           | 1952  |                                   |                      |                             |             |              |              |                     |       |     |               |                 |                                         | 2,300   |
| M.S.           | 1/2   | 14,000                            | 13                   | 1.1                         | 9           | 155          | 128          | 12.2                | 105   | 7.0 | 8             | 2.9             | 4,300                                   | 910     |
| M.S.           | 1/8   | 12,800                            | 7.9                  | 0.51                        | 8           | 68           |              | 11.7                | 99    | 7.2 | 10            | 3.4             | 15,000                                  | 750     |
| M.S.           | 1/14  | 11,400                            | 7.5                  | 0.57                        | 10          | 63           |              | 11.5                | 101   | 6.8 | 10            | 3.3             | 4,300                                   | 1,500   |
| M.S.           | 1/21  | 5,410                             | 9.4                  | 0.72                        | 11          | 40           | 88           | 10.5                | 95    | 7.2 | 14            | 4.2             | 46,000                                  | 24,000  |
| M.S.           | 2/11  | 5,260                             | 5.6                  | 0.54                        | 11          | 63           |              | 11.0                | 99    | 7.4 | 13            | 3.2             | 230                                     | 230     |
| M.S.           | 3/5   | 37,900                            | 24                   | 2.1                         | 11          | 440          |              | 10.9                | 98    | 6.9 | 9             | 3.5             | 4,300                                   | 2,300   |

**SAVANNAH RIVER STUDIES**  
**Summary of Individual Results - Sanitary Examinations**

| Sampling Point                   | Date         | Disch.<br>at<br>Augusta<br>Sec-Ft | O.C.<br>Moore<br>ppm | 5-Day<br>20°C<br>BOD<br>ppm | Temp.<br>°C | Turb.<br>ppm | Color<br>ppm | Dissolved<br>Oxygen |       | pH  | Alkal.<br>ppm | Chloride<br>ppm | Coliforms<br>MPN per<br>Ga. 100 ml | S.C.   |
|----------------------------------|--------------|-----------------------------------|----------------------|-----------------------------|-------------|--------------|--------------|---------------------|-------|-----|---------------|-----------------|------------------------------------|--------|
|                                  |              |                                   |                      |                             |             |              |              | ppm                 | % Sat |     |               |                 |                                    |        |
| S 217.2;<br>HQO 0.01<br>(Cont'd) | 1951<br>9/24 |                                   | 6.0                  | 1.0                         | 24          | 81           | 20           | 7.8                 | 91    | 7.7 | 21            | 3.4             | 93,000                             |        |
|                                  | 10/15        |                                   | 5.2                  | 1.2                         | 19          | 40           | 31           | 8.7                 | 81    | 7.4 | 19            | 3.8             | 430,000                            |        |
|                                  | 12/10        |                                   | 15                   | 0.87                        | 13          | 190          | 108          | 10.1                | 95    | 6.8 | 11            | 5.0             | 93,000                             |        |
|                                  | 1952<br>1/8  |                                   | 9.4                  | 1.2                         | 8           | 73           | 104          | 11.6                | 98    | 7.0 | 9             | 3.3             | 24,000                             |        |
| S 215.6                          | 1951<br>7/30 | 4,800                             | 4.2                  | 0.44                        | --          | 60           | 32           | 6.9                 | --    | 7.2 |               |                 | 24,000                             |        |
|                                  | 7/30         | 4,800                             | 5.5                  | 0.35                        | --          | 60           | 37           | 6.9                 | --    | 7.3 | 20            | 1.7             | 46,000                             |        |
|                                  | 7/30         | 4,800                             | 7.6                  | 0.40                        | --          | 60           | 38           | 6.9                 | --    | 7.3 |               |                 | 43,000                             | 46,000 |
|                                  | 8/13         | 3,580                             | 3.8                  | 0.46                        | 31          | 36           | 12           | 6.6                 | 88    | 7.4 | 18            | 2.2             | 93,000                             |        |
|                                  | 8/13         | 3,580                             | 3.4                  | 0.56                        | 31          | 37           |              | 6.6                 | 88    | 7.4 | 19            | 2.0             | 43,000                             |        |
|                                  | 8/13         | 3,580                             | 3.0                  | 0.54                        | 31          | 37           |              | 6.6                 | 88    | 7.4 | 19            | 2.2             | 43,000                             |        |
|                                  | 8/22         | 3,470                             | 5.5                  | 0.07                        | 30          | 80           | 20           | 6.3                 | 84    | 7.3 | 16            | 3.0             | 93,000                             |        |
|                                  | 8/22         | 3,470                             | 6.2                  | 0.44                        | 30          | 80           |              | 6.7                 | 88    | 7.3 | 15            | 3.2             | 43,000                             |        |
|                                  | 8/22         | 3,470                             | 5.0                  | 0.16                        | 30          | 83           |              | 6.4                 | 84    | 7.3 | 14            | 2.7             | 240,000                            | 23,000 |
|                                  | 9/11         | 2,800                             | 4.5                  | 0.73                        | 27          | 27           | 10           | 7.2                 | 90    | 7.6 | 18            | 3.2             | 23,000                             | 43,000 |
|                                  | 9/24         | 4,720                             | 0.83                 | 0.77                        | 24          | 55           | 25           | 7.8                 | 91    | 7.6 | 19            | 3.8             | 43,000                             |        |
|                                  | 8/9          |                                   | 42                   | 11                          | --          | 39           | 110          | 0                   | --    | 8.3 | 100           | 5.5             | 43,000                             |        |
|                                  | 8/15         |                                   | 31                   | 11                          | 28          | 32           | 80           | 2.5                 | 32    | 8.5 | 50            | 5.4             | 430,000                            |        |
| S 213.36;<br>Ho 2.7              | 8/22         |                                   | 41                   | >23                         | 28          | 25           | 80           | 0                   | 0     | 8.5 | 41            | 8.3             | 430,000                            |        |
|                                  | 9/11         |                                   | 27                   | 6.5                         | 25          | <25          | 70           | 0.16                | 2     | 7.8 | 69            | 5.5             | 430,000                            |        |
|                                  | 9/26         |                                   | 20                   | 7.4                         | 22          | 7            | 80           | 2.9                 | 33    | 7.8 | 57            | 5.3             | 43,000                             |        |
|                                  | 10/8         |                                   | 20                   | 2.7                         | 22          | 7            | 60           | 3.4                 | 38    | 7.7 | 60            | 5.5             | 240,000                            |        |
|                                  | 12/4         |                                   | 44                   | 14                          | 14          | 4            | 58           | 2.5                 | 24    | 7.7 | 61            | 5.5             | 9,300                              |        |
|                                  | 1952<br>1/2  |                                   | 32                   | 8.9                         | 14          | 4            | 72           | 2.8                 | 27    | 7.5 | 44            | 4.5             | 230                                |        |
|                                  | 2/11         |                                   | 16                   | 4.5                         | 13          | 2            | 50           | 4.2                 | 40    | 7.8 | 45            | 5.1             | 15,000                             |        |
|                                  | 3/5          |                                   | 15                   | 4.0                         | 12          | 93           | 80           | 6.3                 | 58    | 7.2 | 13            | 3.4             | 7,500                              |        |

**SAVANNAH RIVER STUDIES**  
**Summary of Individual Results - Sanitary Examinations**

| Sampling Point | Date  | Disch.<br>at<br>Augusta<br>Sec-Ft | O.C.<br>Moore<br>ppm | 5-Day<br>20°C<br>BOD<br>ppm | Temp.<br>°C | Turb.<br>ppm | Color<br>ppm | Dissolved<br>Oxygen |       | pH  | Alkal.<br>ppm | Chloride<br>ppm | Coliforms<br>MPN per<br>Ga. 100 ml S.C. |         |
|----------------|-------|-----------------------------------|----------------------|-----------------------------|-------------|--------------|--------------|---------------------|-------|-----|---------------|-----------------|-----------------------------------------|---------|
|                |       |                                   |                      |                             |             |              |              | ppm                 | % Sat |     |               |                 |                                         |         |
|                |       |                                   |                      |                             |             |              |              |                     |       |     |               |                 |                                         |         |
| 1951           |       |                                   |                      |                             |             |              |              |                     |       |     |               |                 |                                         |         |
| S 212.0        | 8/13  | 3,580                             | 3.2                  | 0.47                        | 31          | 36           | 12           | 6.4                 | 86    | 7.2 | 18            | 2.5             | 93,000                                  |         |
| Ga.            | 8/13  | 3,580                             | 2.9                  | 0.54                        | 31          | 37           |              | 6.3                 | 84    | 7.3 | 20            | 2.4             | 43,000                                  | 43,000  |
| M.S.           | 8/13  | 3,580                             | 4.2                  | 0.70                        | 31          | 38           |              | 6.4                 | 86    | 7.3 | 20            | 2.6             |                                         |         |
| S.C.           | 8/22  | 3,470                             | 5.3                  | 0.64                        | 30          | 80           | 25           | 6.1                 | 80    | 7.4 | 15            | 3.3             | 150,000                                 |         |
| Ga.            | 8/22  | 3,470                             | 5.4                  | 0.96                        | 30          | 82           |              | 6.1                 | 80    | 7.3 | 16            | 3.2             | 75,000                                  |         |
| M.S.           | 8/22  | 3,470                             | 5.6                  | 0.61                        | 30          | 81           |              | 6.1                 | 80    | 7.3 | 15            | 3.3             | 93,000                                  |         |
| S.C.           | 9/12  | 2,080                             | 5.0                  | 1.1                         | 28          | <25          | 15           | 5.5                 | 70    | 7.7 | 23            | 4.0             | 240,000                                 |         |
| Ga.            | 9/12  | 2,080                             | 4.0                  | 1.1                         | 28          | <25          |              | 6.0                 | 76    | 7.7 | 21            | 3.9             | 460,000                                 | 460,000 |
| M.S.           | 9/12  | 2,080                             | 3.6                  | 1.2                         | 28          | <25          |              | 6.1                 | 77    | 7.7 | 21            | 3.4             |                                         |         |
| S.O.           | 9/26  | 6,030                             | 6.3                  | 0.83                        | 27          | 85           | 43           | 6.8                 | 84    | 7.5 | 19            | 3.9             | 230,000                                 |         |
| Ga.            | 9/26  | 6,030                             | 7.6                  | 1.1                         | 26          | 83           | 40           | 7.0                 | 85    | 7.5 | 20            | 4.2             | 23,000                                  |         |
| M.S.           | 9/26  | 6,030                             | 6.1                  | 1.1                         | 26          | 67           | 40           | 6.8                 | 83    | 7.5 | 24            | 3.6             |                                         | 23,000  |
| S.C.           | 10/8  | 2,560                             | 10                   | 0.49                        | 26          | 78           | 25           | 6.9                 | 84    | 7.3 | 20            | 4.5             | 93,999                                  | 93,000  |
| M.S.           | 10/15 | 2,360                             | 6.1                  | 0.51                        | 19          | 32           | 37           | 8.0                 | 86    | 7.5 | 22            | 10              | 43,000                                  | 43,000  |
| M.S.           | 10/24 | 1,960                             | 6.9                  | 1.2                         | 19          | 28           | 27           | 7.1                 | 76    | 7.4 | 24            | 4.4             | 150,000                                 | 240,000 |
| M.S.           | 10/29 | 3,840                             | 5.6                  | 0.57                        | 19          | 25           | 26           | 7.7                 | 83    | 7.5 | 23            | 3.8             | 24,000                                  | 9,300   |
| M.S.           | 12/4  | 4,840                             | 7.5                  | 0.16                        | 11          | 11           | 30           | 10.8                | 97    | 7.0 | 15            | 4.2             | 43,000                                  | 24,000  |
| M.S.           | 12/10 | 5,460                             | 10                   | 0.66                        | 14          | 180          | 120          | 9.6                 | 53    | 7.0 | 13            | 3.6             | 43,000                                  | 43,000  |
| M.S.           | 12/19 | 9,990                             | 5.4                  | 0.50                        | 6           | 62           | 50           | 12.3                | 99    | 7.0 | 13            | 3.3             | 43,000                                  | 9,300   |
| M.S.           | 12/26 | 14,300                            | 18                   | 1.0                         | 7           | 330          | 128          | 12.4                | 102   | 7.0 | 8             | 3.0             | 24,000                                  | 4,300   |
| 1952           |       |                                   |                      |                             |             |              |              |                     |       |     |               |                 |                                         |         |
| M.S.           | 1/2   | 14,000                            | 13                   | 1.1                         | 9           | 155          | 128          | 12.2                | 105   | 7.0 | 8             | 2.9             | 4,300                                   | <360    |
| M.S.           | 1/8   | 12,800                            | 7.9                  | 0.51                        | 8           | 68           | 108          | 11.7                | 99    | 7.2 | 10            | 3.4             | 15,000                                  | 750     |
| M.S.           | 1/14  | 11,400                            | 7.5                  | 0.57                        | 10          | 63           | 116          | 11.5                | 101   | 6.8 | 10            | 3.3             | 4,300                                   | 1,500   |
| M.S.           | 1/21  | 5,410                             | 9.4                  | 0.72                        | 11          | 40           | 88           | 10.5                | 95    | 7.2 | 14            | 4.2             | 46,000                                  | 24,000  |
| M.S.           | 2/11  |                                   | 5.6                  | 0.54                        | 11          | 63           | 120          | 11.0                | 99    | 7.4 | 13            | 3.2             | 230                                     | 230     |
| M.S.           | 3/5   |                                   |                      | 2.1                         | 11          | 440          | 80           | 10.9                | 98    | 6.9 | 9             | 3.5             | 4,300                                   | 2,300   |

**SAVANNAH RIVER STUDIES**  
**Summary of Individual Results - Sanitary Examinations**

| Sampling Point | Date         | Disch.<br>at<br>Augusta<br>Sec-Ft | O.C.<br>Moore BOD<br>ppm | 5-Day<br>20°C<br>BOD<br>ppm | Temp.<br>°C | Turb.<br>ppm | Color<br>ppm | Dissolved<br>Oxygen |       | pH  | Alkal.<br>ppm | Chloride<br>ppm | Coliforms<br>MPN per<br>100 ml S.C. |        |
|----------------|--------------|-----------------------------------|--------------------------|-----------------------------|-------------|--------------|--------------|---------------------|-------|-----|---------------|-----------------|-------------------------------------|--------|
|                |              |                                   |                          |                             |             |              |              | ppm                 | % Sat |     |               |                 | Ga.                                 | S.C.   |
| S 203.3        | 1951<br>8/1  | 4,440                             | 5.7                      | 0.40                        | --          | 54           | 37           | 6.4                 | --    | 7.2 | 20            | 3.0             | 2,900                               |        |
|                | 8/1          | 4,440                             | 5.1                      | 0.35                        | --          | 54           |              | 6.4                 | --    | 7.6 | 20            | 5.2             | 43,000                              |        |
|                | 8/1          | 4,440                             | 5.9                      | 0.49                        | --          | 52           |              | 6.6                 | --    | 7.3 | 20            | 3.3             |                                     | 24,000 |
|                | 8/14         | 3,800                             | 4.6                      | 0.49                        | 30          | 31           | 15           | 6.2                 | 81    | 7.4 | 20            | 3.0             | 24,000                              |        |
|                | 8/14         | 3,800                             | 4.0                      | 0.33                        | 30          | 31           |              | 6.0                 | 79    | 7.3 | 19            | 3.2             | 46,000                              |        |
|                | 8/14         | 3,800                             | 5.6                      | 0.40                        | 30          | 31           |              | 5.9                 | 77    | 7.4 | 21            | 3.0             | 46,000                              | 45,000 |
| S 202.9        | 8/30         | 1,890                             | 3.2                      | 0.30                        | 29          | 54           | 20           | 7.5                 | 97    | 7.2 | 18            | 4.0             | 2,300                               |        |
|                | 8/30         | 1,790                             | 1.7                      | 0.41                        | 29          | 56           | 20           | 7.5                 | 97    | 7.3 | 18            | 3.8             | 9,300                               |        |
|                | 8/30         | 1,890                             | 2.0                      | 0.37                        | 29          | 47           | 20           | 7.4                 | 95    | 7.3 | 17            | 3.4             |                                     |        |
|                | 9/12         | 2,080                             | 1.4                      | 0.47                        | 28          | <25          | 10           | 7.9                 | 100   | 7.7 | 19            | 3.3             | 24,000                              | 1,500  |
|                | 10/1         | 4,800                             | 11                       | 0.42                        | 23          | 170          | 48           | 8.2                 | 95    | 7.4 | 17            | 3.6             | 24,000                              | 4,300  |
|                | 10/9         | 2,560                             | 7.7                      | 0.50                        | 23          | 75           | 25           | 8.2                 | 95    | 7.3 | 19            | 4.0             | 23,000                              | 23,000 |
|                | 10/16        | 2,280                             | 4.5                      | 0.62                        | 20          | 31           | 30           | 8.9                 | 97    | 7.5 | 24            | 4.0             | 43,000                              | 24,000 |
|                | 10/31        | 2,860                             | 5.0                      | 0.70                        | 19          | 25           | 19           | 8.9                 | 95    | 7.5 | 22            | 4.6             | 240,000                             | 93,000 |
|                | 12/5         | 5,550                             | 5.8                      | 0.20                        | 11          | 28           | 32           | 11.1                | 100   | 7.1 | 17            | 4.3             | 43,000                              | 43,000 |
|                | 12/11        | 6,130                             | 11                       | 0.61                        | 12          | 150          | 128          | 10.2                | 94    | 6.7 | 11            | 3.2             | 24,000                              | 24,000 |
|                | 12/18        | 9,150                             | 8.3                      | 0.61                        | 7           | 51           | 56           | 11.8                | 97    | 7.2 | 17            | 3.7             | 7,500                               | 2,300  |
|                | 12/27        | 14,300                            | 16                       | 1.1                         | 6           | 290          | 144          | 12.1                | 97    | 6.5 | 7.0           | 2.6             | 15,000                              | 46,000 |
|                | 1952<br>1/10 | 11,600                            | 12                       | 0.32                        | 9           | 100          | 156          | 11.3                | 98    | 6.8 | 11            | 3.0             | 2,300                               | 9,300  |
|                | 1/16         | 8,320                             | 11                       | 0.68                        | 12          | 52           | 124          | 11.0                | 101   | 7.2 | 10            | 3.2             | 23,000                              | 23,000 |
|                | 1/22         | 5,040                             | 8.0                      | 0.43                        | 11          | 36           | 88           | 11.2                | 101   | 6.9 | 14            | 3.7             | 23,000                              | 4,300  |
|                | 1/29         | 6,730                             | 8.8                      | 0.92                        | 11          | 49           | 94           | 10.8                | 97    | 7.2 | 14            | 4.3             | 43,000                              | 43,000 |
|                | 2/13         | 5,080                             | 10                       | 1.1                         | 12          | 50           | 112          | 11.0                | 101   | 7.3 | 15            | 3.5             | 9,300                               | 23,000 |
|                | 3/6          | 38,600                            | 11                       | 1.4                         | 11          | 205          | 112          | 11                  | 99    | 7.0 | 9             | 3.2             | 4,300                               | 1,500  |
|                | 3/14         | 33,600                            | 11                       | 1.0                         | 12          | 175          | 160          | 10.8                | 100   | 7.1 | 13            | 3.3             | 2,300                               | 2,300  |



**SAVANNAH RIVER STUDIES**  
**Summary of Individual Results - Sanitary Examinations**

| Sampling Point                                                      | Date  | Disch.<br>at<br>Augusta<br>Sec-Ft | O.C.<br>; ore<br>ppm | 5-Day<br>20°C<br>BOD<br>ppm | Temp.<br>°C | Turb.<br>ppm | Color<br>ppm | Dissolved<br>Oxygen |        | pH  | Alkal.<br>ppm | Chloride<br>ppm | Coliforms<br>MPN per<br>Ga. 100 ml | S.C. |
|---------------------------------------------------------------------|-------|-----------------------------------|----------------------|-----------------------------|-------------|--------------|--------------|---------------------|--------|-----|---------------|-----------------|------------------------------------|------|
|                                                                     |       |                                   |                      |                             |             |              |              | ppm                 | % Sat. |     |               |                 |                                    |      |
| * S 202.65;<br>Bu 0.02                                              | 1951  |                                   |                      |                             |             |              |              |                     |        |     |               |                 |                                    |      |
|                                                                     | 8/1   |                                   | 33                   | 7.4                         | --          | 61           | 45           | 3.2                 | --     | 6.6 | 19            | 2.9             | 2,100,000                          |      |
|                                                                     | 8/14  |                                   | 34                   | 21                          | 29          | 31           | 45           | 2.1                 | 27     | 6.7 | 17            | 3.1             | 11,000,000                         |      |
|                                                                     | 8/23  |                                   | 29                   | 10                          | 27          | 40           | 35           | 1.8                 | 22     | 7.2 | 28            | 3.3             | 9,300,000                          |      |
|                                                                     | 8/30  |                                   | 32                   | 11                          | 27          | 31           | 43           | 1.0                 | 12     | 7.3 | 35            | 12              | 2,300,000                          |      |
|                                                                     | 9/10  |                                   | 17                   | 3.9                         | 23          | <25          | 38           | 4.2                 | 48     | 7.1 | 23            | 7.3             | 430,000                            |      |
|                                                                     | 9/25  |                                   | 33                   | 5.4                         | 24          | 150          | 40           | 4.4                 | 51     | 6.9 | 9             | 4.9             | 1,500,000                          |      |
|                                                                     | 10/1  |                                   | 28                   | 9.1                         | 22          | 93           | 55           | 3.7                 | 42     | 7.0 | 23            | 7.3             | 3,900,000                          |      |
|                                                                     | 10/9  |                                   | 34                   | 10                          | 20          | 33           | 54           | 3.7                 | 40     | 7.1 | 29            | 10              | 15,000,000                         |      |
|                                                                     | 10/16 |                                   | 30                   | 11                          | 19          | 13           | 46           | 4.4                 | 47     | 7.0 | 27            | 9.7             | 4,300,000                          |      |
|                                                                     | 10/31 |                                   | 37                   | 12                          | 20          | 65           | 46           | 4.2                 | 46     | 7.0 | 23            | 10              | 4,300,000                          |      |
|                                                                     | 12/11 |                                   | 30                   | 15                          | 11          | 38           | 80           | 5.1                 | 46     | 6.7 | 16            | 7.8             | 2,100,000                          |      |
|                                                                     | 12/18 |                                   | 24                   | 12                          | 7           | 29           | 50           | 6.9                 | 57     | 6.9 | 16            | 6.7             | 930,000                            |      |
|                                                                     | 12/27 |                                   | 22                   | 5.6                         | 9           | 62           | 92           | 5.2                 | 45     | 6.3 | 9             | 8.1             | 36,000                             |      |
|                                                                     | 1952  |                                   |                      |                             |             |              |              |                     |        |     |               |                 |                                    |      |
|                                                                     | 1/10  |                                   | 30                   | 9.2                         | 12          | 45           | 84           | 4.0                 | 37     | 6.7 | 16            | 7.9             | 2,400,000                          |      |
|                                                                     | 1/16  |                                   | 35                   | 7.4                         | 14          | 58           | 50           | 3.7                 | 36     | 7.2 | 14            | 8.0             | 430,000                            |      |
|                                                                     | 1/22  |                                   | 41                   | 12                          | 13          | 67           | 58           | 4.8                 | 45     | 6.7 | 14            | 7.7             | 2,300,000                          |      |
|                                                                     | 1/29  |                                   | 26                   | 8.2                         | 11          | 70           | 70           | 5.3                 | 48     | 6.9 | 15            | 8.6             | 430,000                            |      |
|                                                                     | 3/13  |                                   | 24                   | 8.9                         | 13          | 37           | 56           | 2.3                 | 23     | 6.9 | 16            | 7.3             | 430,000                            |      |
| S 202.65; Bu 2.2<br>(Butler Creek at<br>Old Savannah<br>Rd. Bridge) | 1951  |                                   |                      |                             |             |              |              |                     |        |     |               |                 |                                    |      |
|                                                                     | 8/9   |                                   | 10                   | 0.27                        | --          | 25           | 50           | 6.4                 | --     | 6.4 | 6.0           | 5.0             | 1,500                              |      |
|                                                                     | 8/14  |                                   | 9.6                  | 0.62                        | 29          | <25          | 50           | 6.3                 | 84     | 6.3 | 6.2           | 3.7             | 9,300                              |      |
|                                                                     | 8/23  |                                   | 9.0                  | 0.46                        | 27          | <25          | 45           | 6.8                 | 84     | 6.3 | 5.3           | 3.7             | 2,100                              |      |
|                                                                     | 8/30  |                                   | 6.6                  | 0.61                        | 26          | <25          | 48           | 7.1                 | 86     | 6.3 | 5.2           | 4.0             | 1,500                              |      |
|                                                                     | 9/10  |                                   | 8.7                  | 0.41                        | 24          | <25          | 50           | 7.4                 | 86     | 6.6 | 4.2           | 4.1             | 2,300                              |      |
|                                                                     | 9/25  |                                   | 5.8                  | 0.69                        | 22          | 10           | 50           | 6.0                 | 68     | 6.1 | 4.3           | 4.4             | 2,300                              |      |
|                                                                     | 10/9  |                                   | 12                   | 0.47                        | 21          | 6            | 48           | 7.7                 | 86     | 6.3 | 5.5           | 7.3             | 2,300                              |      |
|                                                                     | 12/27 |                                   | 8.2                  | 0.38                        | 9           | 2            | 29           | 10.7                | 92     | 4.9 | 2.0           | 3.4             | 210                                |      |

\* Butler Creek, 100' before its confluence with the Savannah River at S 202.65



**SAVANNAH RIVER STUDIES**  
Summary of Individual Results - Sanitary Examinations

| Sampling Point                               | Date      | Disch.<br>at<br>Augusta<br>Sec-Ft | O.C.<br>Moore<br>ppm | 5-Day<br>20°C<br>BOD<br>ppm | Temp.<br>°C | Turb.<br>ppm | Color<br>ppm | Dissolved<br>Oxygen |       | pH  | Alkal.<br>ppm | Chloride<br>ppm | Coliforms<br>MPN per |         | S.C.   |
|----------------------------------------------|-----------|-----------------------------------|----------------------|-----------------------------|-------------|--------------|--------------|---------------------|-------|-----|---------------|-----------------|----------------------|---------|--------|
|                                              |           |                                   |                      |                             |             |              |              | ppm                 | % Sat |     |               |                 | OO                   | l       |        |
|                                              |           |                                   |                      |                             |             |              |              |                     |       |     |               |                 |                      |         |        |
| 1952                                         |           |                                   |                      |                             |             |              |              |                     |       |     |               |                 |                      |         |        |
| S 202.65;<br>Bu 2.2<br>(Cont'd)              | 1/24      |                                   | 11                   | .47                         | 9           | 2            | 54           | 10.1                | 87    | 6.0 | 2.9           | 3.3             |                      | 930     |        |
|                                              | 2/28      |                                   | 8.8                  | 0.85                        | 9           | 2            | 36           | 10.5                | 91    | 6.1 | 3.2           | 4.6             |                      | 93      |        |
|                                              | 4/2       |                                   | 11                   | 0.61                        | 17          | 2            | 52           | 8.6                 | 88    | 6.1 | 5.3           | 4.0             |                      | 930     |        |
|                                              | 4/10      |                                   | 8.7                  | 0.51                        | 17          | 1            | 26           | 8.4                 | 86    | 6.0 | 6.0           | 4.7             |                      | 2,300   |        |
| * S 202.65;<br>Bu 0.27;<br>Be 2.5;<br>Ro 3.6 | 8/9/51    |                                   | 17                   | 2.0                         | --          | <25          | 15           | 3.4                 | --    | 6.7 | 20            | 5.2             |                      | 430,000 |        |
|                                              | 8/14      |                                   | 10                   | 1.5                         | 27          | <25          | 45           | 4.0                 | 50    | 6.7 | 13            | 4.7             |                      | 93,000  |        |
|                                              | 8/23      |                                   | 24                   | 8.9                         | 26          | <25          | 70           | 1.4                 | 16    | 6.5 | 11            | 3.5             |                      | 93,000  |        |
|                                              | 8/30      |                                   | 18                   | 4.6                         | 27          | <25          | 60           | 1.6                 | 20    | 6.6 | 18            | 7.3             |                      | 230,000 |        |
|                                              | 9/10      |                                   | 20                   | 4.8                         | 24          | <25          | 50           | 2.3                 | 27    | 7.0 | 18            | 6.1             |                      | 39,000  |        |
|                                              | 9/25      |                                   | 26                   | 3.7                         | 22          | 90           | 70           | 5.0                 | 56    | 6.7 | 4.8           | 5.6             |                      | 240,000 |        |
|                                              | 10/9      |                                   | 16                   | 2.8                         | 20          | 5            | 56           | 5.0                 | 55    | 6.7 | 11            | 6.1             |                      | 43,000  |        |
|                                              | 12/27     |                                   | 24                   | 8.7                         | 10          | 8            | 56           | 8.8                 | 78    | 6.2 | 7.0           | 6.7             |                      | 240,000 |        |
| 1952                                         |           |                                   |                      |                             |             |              |              |                     |       |     |               |                 |                      |         |        |
| 1/24                                         |           | 36                                | 8.7                  | 9                           | 14          | 64           | 8.8          | 8.8                 | 76    | 6.2 | 4.5           | 6.1             |                      | 43,000  |        |
|                                              | 2/28      |                                   | 29                   | 8.3                         | 10          | 42           | 56           | 8.4                 | 74    | 6.5 | 6.8           | 6.7             |                      | 93,000  |        |
|                                              | 4/2       |                                   | 30                   | 4.8                         | 17          | 145          | 120          | 6.0                 | 62    | 6.4 | 6.9           | 5.3             |                      | 460,000 |        |
|                                              | 4/10      |                                   | 32                   | .9                          | 17          | 13           | 80           | 6.4                 | 66    | 6.5 | 12            | 6.8             |                      | 430,000 |        |
| 1951                                         |           |                                   |                      |                             |             |              |              |                     |       |     |               |                 |                      |         |        |
| S 201.65                                     | Ga. 8/1   | 4,440                             | 5.0                  | 0.41                        | --          | 58           |              | 6.9                 | --    | 7.3 | 20            | 3.0             | 75,000               | 93,000  | 75,000 |
|                                              | M.S. 8/1  | 4,440                             | 5.0                  | 0.57                        | --          | 58           | 40           | 7.1                 | --    | 7.2 | 18            | 3.4             |                      |         |        |
|                                              | S.C. 8/1  | 4,440                             | 4.8                  | 0.46                        | --          | 54           |              | 7.1                 | --    | 7.2 | 20            | 3.5             | 24,000               |         | 24,000 |
|                                              | Ga. 8/14  | 800                               | .4                   | .65                         | 31          | 33           |              | .3                  | 37    | 7.3 | 19            | 2.9             |                      |         |        |
|                                              | M.S. 8/14 | 3,800                             | 4.2                  | 0.70                        | 31          | 32           | 20           | 7.4                 | 98    | 7.3 | 19            | 2.7             | 150,000              |         | 43,000 |
|                                              | S.C. 8/14 | 3,800                             | 3.7                  | 0.8                         | 31          | 32           |              | 7.3                 | 98    | 7.3 | 19            | 2.8             |                      |         |        |
|                                              | Ga. 8/23  | 2,150                             | 3.0                  | 0.48                        | 29          | 48           | 25           | 7.0                 | 90    | 7.2 | 18            | 3.3             | 93,000               | 240,000 | 43,000 |
|                                              | M.S. 8/29 | 50                                | 2.9                  | 0.47                        | 29          | 47           | 25           | 7.2                 | 92    | 7.2 | 17            | 3.5             |                      |         |        |
|                                              | S.C. 8/29 | 2,150                             | 2.7                  | 0.35                        | 29          | 51           | 25           | 7.3                 | 94    | 7.2 | 17            | 3.2             |                      |         |        |

\* Rocky Creek at Old Savannah Rd. Bridge.

SAVANNAH RIVER STUDIES  
Summary of Individual Results - Sanitary Examinations

| Sampling Point  | Date  | Disch.<br>at<br>Augusta<br>Sec-Ft | O.C.<br>Moore<br>ppm | 5-Day<br>20°C<br>BOD<br>ppm | Temp.<br>°C | Turb.<br>ppm | Color<br>ppm | Dissolved<br>Oxygen |       | pH  | Alkal.<br>ppm | Chloride<br>ppm | Coliforms<br>MPN per<br>100 ml |         | S.C.    |
|-----------------|-------|-----------------------------------|----------------------|-----------------------------|-------------|--------------|--------------|---------------------|-------|-----|---------------|-----------------|--------------------------------|---------|---------|
|                 |       |                                   |                      |                             |             |              |              | ppm                 | % Sat |     |               |                 | Ga.                            |         |         |
| 1951            |       |                                   |                      |                             |             |              |              |                     |       |     |               |                 |                                |         |         |
| S 201.65 M.S.   | 9/12  | 2,080                             | 3.5                  | 0.94                        | 28          | <25          | 10           | 7.9                 | 99    | 7.6 | 20            | 3.3             | 93,000                         | 93,000  | 93,000  |
| (Cont'd) M.S.   | 10/1  | 4,800                             | 11                   | 0.29                        | 23          | 165          | 45           | 8.2                 | 94    | 7.4 | 18            | 4.0             | 93,000                         | 23,000  | 240,000 |
| M.S.            | 12/18 | 9,150                             | 6.8                  | 0.42                        | 7           | 50           | 50           | 12                  | 96    | 7.1 | 13            | 3.2             | 24,000                         | 24,000  | 43,000  |
| 1952            |       |                                   |                      |                             |             |              |              |                     |       |     |               |                 |                                |         |         |
| M.S.            | 1/16  | 8,320                             | 11                   | 0.79                        | 12          | 48           | 116          | 10.7                | 99    | 7.0 | 10            | 3.7             | 150,000                        | 460,000 | 9,300   |
| M.S.            | 3/6   | 38,600                            | 12                   | 1.1                         | 11          | 200          | 120          | 11.0                | 100   | 6.8 | 7.5           | 3.1             | 2,300                          | 2,300   | 3,900   |
| 1951            |       |                                   |                      |                             |             |              |              |                     |       |     |               |                 |                                |         |         |
| S 198.4; Sp 3.7 | 8/9   |                                   | 12                   | 0.27                        | --          | <25          | 33           | 5.9                 | --    | 6.4 | 12            | 4.6             |                                | 290     |         |
| 1951            |       |                                   |                      |                             |             |              |              |                     |       |     |               |                 |                                |         |         |
| S 197.0 Ga.     | 8/1   | 4,440                             | 5.5                  | 0.94                        | --          | 48           | 35           | 7.0                 | --    | 7.2 | 18            | 3.2             | 24,000                         | 150,000 |         |
| M.S.            | 8/1   | 4,440                             | 6.3                  | 0.31                        | --          | 48           |              | 7.0                 | --    | 7.2 | 18            | 3.5             |                                |         |         |
| S.C.            | 8/1   | 4,440                             | 4.7                  | 0.52                        | --          | 46           |              | 7.0                 | --    | 7.2 | 18            | 3.5             |                                |         | 210,000 |
| Ga.             | 8/15  | 4,720                             | 4.9                  | 0.99                        | 30          | 38           | 15           | 6.8                 | 89    | 7.4 | 18            | 3.5             |                                | 43,000  |         |
| M.S.            | 8/15  | 4,720                             | 4.4                  | 0.44                        | 30          | 36           |              | 6.6                 | 86    | 7.3 | 18            | 3.2             |                                |         |         |
| S.C.            | 8/15  | 4,720                             | 4.5                  | 0.88                        | 30          | 35           |              | 6.9                 | 90    | 7.3 | 19            | 3.5             |                                |         | 43,000  |
| Ga.             | 8/29  | 2,150                             | 9.3                  | 0.58                        | 28          | 59           | 25           | 6.8                 | 86    | 7.1 | 18            | 3.1             | 240,000                        |         |         |
| M.S.            | 8/29  | 2,150                             | 3.8                  | 0.58                        | 28          | 58           |              | 6.9                 | 87    | 7.1 | 18            | 3.3             | 93,000                         | 93,000  |         |
| S.C.            | 8/29  | 2,150                             | 2.6                  | 0.62                        | 28          | 58           |              | 6.9                 | 87    | 7.1 | 18            | 3.5             |                                |         | 39,000  |
| M.S.            | 9/13  | 1,830                             | 4.7                  | 1.0                         | 27          | <25          | 10           | 7.3                 | 90    | 7.6 | 21            | 3.4             | 93,000                         | 43,000  | 150,000 |
| M.S.            | 10/1  | 4,800                             | 11                   | 0.49                        | 23          | 160          | 50           | 7.9                 | 91    | 7.4 | 19            | 4.0             | 240,000                        | 75,000  | 75,000  |
| M.S.            | 10/9  | 2,560                             | 7.2                  | 0.82                        | 22          | 78           | 40           | 7.8                 | 88    | 7.2 | 18            | 3.6             | 240,000                        | 240,000 | 460,000 |
| M.S.            | 10/16 | 2,280                             | 5.3                  | 1.3                         | 20          | 33           | 38           | 8.3                 | 90    | 7.4 | 26            | 4.0             | 430,000                        | 75,000  | 460,000 |
| M.S.            | 10/31 | 2,860                             | 5.4                  | 0.60                        | 21          | 27           | 21           | 8.5                 | 95    | 7.5 | 21            | 4.7             | 240,000                        | 240,000 | 430,000 |
| M.S.            | 12/5  | 5,550                             | 6.6                  | 0.44                        | 11          | 27           | 31           | 10.8                | 97    | 7.0 | 15            | 3.8             | 240,000                        | 93,000  | 43,000  |
| M.S.            | 12/11 | 6,130                             | 11                   | 0.45                        | 13          | 150          | 124          | 9.9                 | 93    | 6.8 | 11            | 3.3             | 23,000                         | 15,000  | 43,000  |
| M.S.            | 12/18 | 9,150                             | 6.9                  | 0.73                        | 7           | 47           | 50           | 11.7                | 96    | 7.1 | 13            | 3.7             | 24,000                         | 21,000  | 9,300   |
| M.S.            | 12/27 | 14,300                            | 16                   | 0.94                        | 6           | 290          | 152          | 12.0                | 96    | 6.3 | 6             | 3.2             | 24,000                         | 2,300   | 7,500   |

\* Spirit Creek at Old Savannah Road Bridge.

**SAVANNAH RIVER STUDIES**  
**Summary of Individual Results - Sanitary Examinations**

| Sampling Point | Date      | Disch.<br>at<br>Augusta<br>Sec-Ft | O.C.<br>Moore<br>ppm | 5-Day<br>B.O.D.<br>20°C<br>ppm | Temp.<br>°C | Turb.<br>ppm | Color<br>ppm | Dissolved<br>Oxygen |       | pH  | Alkal.<br>ppm | Chloride<br>ppm | Ga.<br>100 ml | Coliforms<br>MPN per<br>100 ml | S.C.    |
|----------------|-----------|-----------------------------------|----------------------|--------------------------------|-------------|--------------|--------------|---------------------|-------|-----|---------------|-----------------|---------------|--------------------------------|---------|
|                |           |                                   |                      |                                |             |              |              | ppm                 | % Sat |     |               |                 |               |                                |         |
| 1952           |           |                                   |                      |                                |             |              |              |                     |       |     |               |                 |               |                                |         |
| S 197.0        | M.S. 1/10 | 11,600                            | 12                   | 0.68                           | 9           | 96           | 148          | 11.2                | 97    | 6.8 | 11            | 3.1             | 9,300         | 9,300                          | 4,300   |
| (Cont'd)       | M.S. 1/16 | 8,320                             | 10                   | 0.67                           | 12          | 50           | 108          | 10.8                | 100   | 7.2 | 11            | 3.3             | 15,000        | 43,000                         | 43,000  |
|                | M.S. 1/22 | 5,040                             | 8.8                  | 0.67                           | 11          | 41           | 84           | 10.5                | 95    | 6.9 | 13            | 3.8             | 150,000       | 75,000                         | 43,000  |
|                | M.S. 1/29 | 6,730                             | 8.1                  | 1.2                            | 11          | 48           | 92           | 10.4                | 94    | 7.3 | 13            | 4.0             | 9,300         | 15,000                         | 15,000  |
|                | M.S. 2/13 | 5,080                             | 9.2                  | 1.0                            | 12          | 52           | 112          | 10.7                | 99    | 7.3 | 14            | 3.6             | 43,000        | 23,000                         | 9,300   |
|                | M.S. 3/6  | 38,600                            | 15                   | 1.5                            | 11          | 215          | 120          | 10.9                | 98    | 7.1 | 9.5           | 3.2             | 2,300         | 930                            | 2,300   |
| 1951           |           |                                   |                      |                                |             |              |              |                     |       |     |               |                 |               |                                |         |
| S 192.0        | Ga. 8/2   | 4,260                             | 5.5                  | 0.51                           | --          | 64           | 38           | 6.4                 | --    | 7.1 | 20            | 3.5             | 430,000       |                                | 150,000 |
|                | S.O. 8/2  | 4,260                             | 5.7                  | 0.51                           | --          | 62           |              | 6.4                 | --    | 7.0 | 20            | 3.6             | 24,000        |                                |         |
|                | Ga. 8/15  | 4,720                             | 4.0                  | 0.55                           | 30          | 38           | 15           | 6.7                 | 88    | 7.2 | 18            | 3.3             |               |                                | 93,000  |
|                | S.C. 8/15 | 4,720                             | 5.1                  | 0.54                           | 30          | 37           |              | 6.6                 | 86    | 7.1 | 20            | 3.6             | 240,000       |                                |         |
|                | Ga. 9/6   | 2,170                             | 5.0                  | 0.64                           | 29          | <25          | 15           | 7.0                 | 89    | 7.4 | 20            | 3.3             |               |                                | 9,300   |
|                | S.C. 9/6  | 2,170                             | 4.0                  | 0.46                           | 29          | <25          |              | 6.9                 | 88    | 7.5 | 18            | 4.0             | 75,000        |                                | 23,000  |
|                | M.S. 9/13 | 1,830                             | 1.9                  | 0.59                           | 27          | <25          | 10           | 7.1                 | 88    | 7.6 | 20            | 3.3             |               |                                |         |
| S 189.0        | Ga. 8/2   | 4,260                             | 5.1                  | 0.59                           | --          | 58           | 33           | 6.6                 | --    | 7.0 | 20            | 3.6             | 430,000       |                                | 930,000 |
|                | S.O. 8/2  | 4,260                             | 4.4                  | 0.35                           | --          | 56           |              | 6.3                 | --    | 7.2 | 21            | 3.6             |               |                                |         |
| S 187.0        | Ga. 8/16  | 3,670                             | 5.9                  | 0.51                           | 29          | 29           | 15           | 6.6                 | 85    | 7.4 | 18            | 3.2             | 93,000        |                                |         |
|                | S.C. 8/16 | 3,670                             | 6.2                  | 0.71                           | 29          | 26           |              | 6.6                 | 85    | 7.3 | 18            | 3.5             |               |                                | 93,000  |
|                | M.S. 9/6  | 2,170                             | 3.9                  | 1.7                            | 29          | <25          | 15           | 6.9                 | 89    | 7.5 | 18            | 3.5             | 23,000        |                                |         |
|                | M.S. 9/13 | 1,830                             | 7.0                  | 0.54                           | 27          | <25          | 10           | 7.0                 | 87    | 7.7 | 19            | 3.5             | 75,000        |                                | 93,000  |
| S 182.0        | Ga. 8/2   | 4,260                             | 3.9                  | 0.31                           | --          | 57           | 30           | 6.4                 | --    | 7.0 | 20            | 3.3             | 43,000        |                                | 15,000  |
|                | S.O. 8/2  | 4,260                             | 4.7                  | 0.38                           | --          | 55           |              | 6.5                 | --    | 7.2 | 19            | 3.5             | 24,000        |                                |         |
|                | Ga. 8/16  | 3,670                             | 7.0                  | 0.44                           | 29          | 29           | 15           | 6.1                 | 79    | 7.3 | 18            | 3.6             |               |                                | 21,000  |
|                | S.O. 8/16 | 3,670                             | 5.4                  | 0.87                           | 29          | 28           |              | 6.6                 | 85    | 7.4 | 18            | 3.3             |               |                                |         |
|                | M.S. 9/6  | 2,170                             | 1.6                  | 0.52                           | 29          | 25           | 15           | 6.7                 | 86    | 7.6 | 18            | 3.6             | 9,300         |                                |         |

SAVANNAH RIVER STUDIES  
Summary of Individual Results - Sanitary Examinations

| Sampling Point      | Date       | Disch.<br>at<br>Augusta<br>Sec-Ft | 5-Day<br>20°C<br>BOD<br>ppm | Temp.<br>°C | Turb.<br>ppm | Color<br>ppm | Dissolved<br>Oxygen |       | pH  | Alkal.<br>ppm | Chloride<br>ppm | Coliforms<br>MPN per<br>100 ml |         |
|---------------------|------------|-----------------------------------|-----------------------------|-------------|--------------|--------------|---------------------|-------|-----|---------------|-----------------|--------------------------------|---------|
|                     |            |                                   |                             |             |              |              | ppm                 | % Sat |     |               |                 | Ga.                            | S.C.    |
| S 177.0             | M.S. 8/2   | 4,260                             | 0.36                        | --          | 49           | 30           | 6.5                 | --    | 6.9 | 21            | 3.4             | 24,000                         |         |
|                     | Ga. 8/16   | 3,670                             | 0.62                        | 29          | 29           | 15           | 6.5                 | 84    | 7.4 | 18            | 3.4             | 4,300                          |         |
|                     | S.C. 8/16  | 3,670                             | 5.0                         | 29          | 27           |              | 6.8                 | 87    | 7.3 | 18            | 3.4             |                                | 24,000  |
|                     | Ga. 9/6    | 2,170                             | 1.8                         | 29          | <25          | 15           | 6.9                 | 88    | 7.5 | 18            | 3.5             | 15,000                         |         |
|                     | S.C. 9/6   | 2,170                             | 2.1                         | 29          | 26           |              | 7.0                 | 90    | 7.6 | 19            | 3.5             | >110,000                       |         |
| M.S. 9/13           |            | 1,830                             | 3.2                         | 27          | <25          | 10           | 7.0                 | 87    | 7.6 | 19            | 3.4             | 24,000                         | 9,300   |
|                     |            |                                   |                             |             |              |              |                     |       |     |               |                 |                                |         |
| *S176.67;<br>BG 1.9 | 8/30       |                                   | 2.4                         | 27          | <25          | 25           | 6.4                 | 79    | 7.4 | 42            | 4.0             | 2,300                          |         |
|                     | 9/10       |                                   | 0.54                        | 24          | <25          | 30           | 6.7                 | 79    | 7.4 | 38            | 3.6             | 24,000                         |         |
|                     | 9/18       |                                   | 0.66                        | 22          | <25          | 57           | 6.2                 | 71    | 7.5 | 25            | 3.9             | 4,300                          |         |
| S 172.0             | M.S. 8/2   | 4,260                             | 0.28                        | --          | 49           | 30           | 6.2                 | --    | 7.2 | 20            | 3.3             | 93,000                         |         |
|                     | Ga. 8/16   | 3,670                             | 0.68                        | 30          | 30           | 15           | 6.8                 | 90    | 7.4 | 18            | 3.6             | 46,000                         |         |
|                     | S.C. 8/16  | 3,670                             | 5.0                         | 30          | 29           |              | 6.6                 | 87    | 7.4 | 18            | 3.5             |                                | 15,000  |
|                     | Ga. 9/10   | 2,940                             | 9.5                         | 27          | 33           | 10           | 7.5                 | 93    | 7.5 | 21            | 3.8             | 23,000                         |         |
|                     | S.C. 9/10  | 2,940                             | 7.6                         | 27          | 29           | 10           | 7.6                 | 94    | 7.5 | 19            | 3.8             |                                |         |
|                     | M.S. 9/13  | 1,830                             | 2.0                         | 27          | <25          | 10           | 7.2                 | 89    | 7.7 | 18            | 3.5             | 9,300                          | 43,000  |
|                     | M.S. 10/3  | 3,350                             | 6.6                         | 23          | 195          | 35           | 7.4                 | 85    | 7.3 | 16            | 3.8             | 240,000                        | 39,000  |
|                     | M.S. 10/10 | 2,680                             | 8.3                         | 21          | 76           | 30           | 7.6                 | 84    | 7.4 | 18            | 3.6             | 93,000                         | 93,000  |
|                     | M.S. 10/23 | 1,890                             | 5.3                         | 21          | 27           | 20           | 7.6                 | 85    | 7.3 | 21            | 4.8             | 93,000                         | 93,000  |
|                     | M.S. 11/7  | 4,260                             | 6.9                         | 13          | 49           | 30           | 9.4                 | 89    | 7.7 | 81            | 3.9             | 43,000                         | 23,000  |
|                     | M.S. 12/5  | 5,550                             | 5.0                         | 12          | 33           | 35           | 10.4                | 96    | 7.1 | 17            | 4.1             | 93,000                         | 210,000 |
|                     | M.S. 12/20 | 8,420                             | 6.9                         | 6           | 87           | 52           | 11.9                | 95    | 7.0 | 15            | 3.6             | 24,000                         | 3,900   |
|                     | M.S. 12/28 | 14,200                            | 1.0                         | 6           | 245          | 96           | 12.0                | 96    | 6.6 | 7             | 2.9             | 2,300                          | 2,300   |
|                     | 1952       |                                   |                             |             |              |              |                     |       |     |               |                 |                                |         |
|                     | M.S. 1/3   | 13,900                            | 15                          | 0.91        | 10           | 148          | 11.5                | 102   | 6.9 | 8             | 3.4             | 280                            | 4,300   |
|                     | M.S. 1/9   | 12,800                            | 9.9                         | 0.53        | 9            | 96           | 11.3                | 98    | 6.9 | 8             | 3.2             | 4,300                          |         |
|                     |            |                                   |                             |             |              |              |                     |       |     |               |                 |                                |         |

\* Boggy Gut Creek at River Road Bridge.

**SAVANNAH RIVER STUDIES**  
**Summary of Individual Results - Sanitary Examinations**

| Sampling Point                                     | Date         | Disch.<br>at<br>Augusta<br>Sec-Ft  | O.C.<br>Moore<br>ppm | 5-Day<br>20°C<br>BOD<br>ppm | Temp.<br>°C | Turb.<br>ppm | Color<br>ppm | Dissolved<br>Oxygen |       | pH  | Alkal.<br>ppm | Chloride<br>ppm | Coliforms<br>MPN per<br>100 ml | S.C.    |
|----------------------------------------------------|--------------|------------------------------------|----------------------|-----------------------------|-------------|--------------|--------------|---------------------|-------|-----|---------------|-----------------|--------------------------------|---------|
|                                                    |              |                                    |                      |                             |             |              |              | ppm                 | % Sat |     |               |                 |                                |         |
| S 172.0 M.S.<br>(Cont'd) M.S.<br>M.S.<br>M.S.      | 1952<br>1/15 | 13,200                             | 9.5                  | 0.65                        | 11          | 90           | 100          | 11.0                | 99    | 7.1 | 12            | 3.3             | 4,300                          | 2,300   |
|                                                    | 1/23         | 5,030                              | 9.2                  | 0.87                        | 11          | 61           | 84           | 10.4                | 94    | 6.9 | 14            | 3.7             | 93,000                         | 120,000 |
|                                                    | 2/18         | 6,030                              | 13                   | 0.86                        | 10          | 45           | 113          | 10.7                | 95    | 7.0 | 15            | 4.1             | 15,000                         | 2,100   |
|                                                    | 4/3          | 33,200                             | 10                   | 0.92                        | 14          | 130          | 240          | 8.6                 | 82    | 6.9 | 11            | 3.1             | 2,300                          | 4,300   |
| S 171.59;<br>UTR 0.01                              | 1951<br>8/16 |                                    | 22                   | 0.97                        | 24          | <25          | 55           | 6.9                 | 80    | 6.5 | 7.7           | 3.3             | 9,300                          | 9,300   |
|                                                    | 9/10         |                                    | 5.5                  | 0.12                        | 20          | <25          | 20           | 8.5                 | 93    | 6.7 | 5.4           | 3.3             | 430                            | 1,500   |
|                                                    | 9/13         |                                    | 4.0                  | 0.06                        | 25          | <25          | 20           | 8.0                 | 95    | 7.0 | 7.1           | 3.2             | 2,300                          | 2,300   |
|                                                    | 10/3         |                                    | 5.1                  | 0.34                        | 21          | 4            | 30           | 8.1                 | 90    | 6.7 | 8.0           | 3.9             | 930                            | 2,100   |
|                                                    | 10/10        |                                    | 5.1                  | 0.61                        | 16          | 2            | 22           | 9.1                 | 91    | 6.7 | 7.1           | 3.0             | 2,100                          | 2,100   |
|                                                    | 10/23        |                                    | 13                   | 0.47                        | 19          | 3            | 44           | 7.8                 | 84    | 6.7 | 8.9           | 5.8             | 930                            | 1,500   |
|                                                    | 11/7         |                                    | 13                   | 0.67                        | 13          | 2            | 56           | 8.6                 | 82    | 6.5 | 6.1           | 3.6             | 1,500                          | 4,300   |
|                                                    | 12/5         |                                    | 14                   | 1.1                         | 14          | 2            | 66           | 8.3                 | 80    | 6.4 | 6.0           | 3.9             | 4,300                          | 930     |
|                                                    | 12/20        |                                    | 12                   | 0.73                        | 7           | 3            | 56           | 10.5                | 86    | 5.7 | 4.0           | 3.5             | 930                            | 930     |
|                                                    | 12/28        |                                    | 11                   | 0.74                        | 9           | 8            | 72           | 7.9                 | 68    | 6.1 | 5.0           | 3.3             | 91                             | 240     |
|                                                    | 1952<br>1/3  |                                    | 10                   | 0.48                        | 16          | 2            | 52           | 7.1                 | 71    | 6.6 | 6.0           | 3.5             | 9,300                          | 9,300   |
|                                                    | 1/9          | Disch.<br>at Bur-<br>2/18 tons Py. | 9.9                  | 0.65                        | 11          | 1            | 44           | 8.8                 | 79    | 6.3 | 5.0           | 3.0             | 24,000                         | 9,300   |
|                                                    | 1/23         |                                    | 15                   | 0.61                        | 13          | 5            | 64           | 8.9                 | 84    | 5.5 | 3.5           | 3.4             | 9,300                          | 9,300   |
| S 167.0 M.S.<br>Ga.<br>S.C.<br>Ga.<br>S.C.<br>M.S. | 1951<br>8/2  | 5,290                              | 8.1                  | 0.33                        | --          | 45           | 25           | 6.4                 | --    | 7.2 | 20            | 3.5             | 9,300                          | 23,000  |
|                                                    | 8/16         | 4,930                              | 5.1                  | 0.51                        | 29          | 32           | 20           | 6.6                 | 84    | 7.3 | 18            | 3.4             | 15,000                         | 15,000  |
|                                                    | 8/16         | 4,930                              | 5.0                  | 0.43                        | 29          | 31           | 20           | 6.5                 | 84    | 7.4 | 18            | 3.3             | 23,000                         | 23,000  |
|                                                    | 9/10         | 2,210                              | 6.1                  | 1.3                         | 27          | 27           | 10           | 7.7                 | 96    | 7.5 | 19            | 3.6             | 15,000                         | 15,000  |
|                                                    | 9/10         | 2,210                              | 6.7                  | 1.4                         | 27          | 26           | 10           | 7.7                 | 96    | 7.5 | 19            | 3.8             | 15,000                         | 15,000  |
|                                                    | 9/13         | 2,800                              | 3.4                  | 0.29                        | 27          | <25          | 10           | 7.1                 | 87    | 7.6 | 17            | 3.3             | 15,000                         | 15,000  |



SAVANNAH RIVER STUDIES  
Summary of Individual Results - Sanitary Examinations

| Sampling Point        | Date  | Disch.<br>at<br>Burton's<br>Sec-Ft/y | O.C.<br>Moore<br>ppm | 5-Day<br>20°C<br>BOD<br>ppm | Temp.<br>°C | Turb.<br>ppm | Color<br>ppm | Dissolved<br>Oxygen |       | pH  | Alkal.<br>ppm | Chloride<br>ppm | Ga.<br>100 ml | Coliforms<br>MPN per<br>100 ml | S.C.    |
|-----------------------|-------|--------------------------------------|----------------------|-----------------------------|-------------|--------------|--------------|---------------------|-------|-----|---------------|-----------------|---------------|--------------------------------|---------|
|                       |       |                                      |                      |                             |             |              |              | ppm                 | % Sat |     |               |                 |               |                                |         |
| 1951                  |       |                                      |                      |                             |             |              |              |                     |       |     |               |                 |               |                                |         |
| *S 164.26;<br>Fo 0.01 | 8/20  |                                      | 6.9                  | 0.32                        | 27          | <25          | 35           | 5.7                 | 70    | 7.0 | 15            | 3.5             |               | 930                            |         |
|                       | 9/4   |                                      | 6.0                  | 0.25                        | 27          | <25          | 25           | 6.1                 | 76    | 7.0 | 14            | 3.5             |               | 430                            |         |
|                       | 9/17  |                                      | 31                   | 1.6                         | 22          | 220          | 56           | 5.1                 | 57    | 6.5 | 5.9           | 3.4             |               | 4,500                          |         |
|                       | 10/3  |                                      | 7.5                  | 0.42                        | 21          | 7            | 50           | 6.7                 | 74    | 6.9 | 13            | 4.3             |               | 230                            |         |
|                       | 10/10 |                                      | 9.8                  | 0.32                        | 18          | 12           | 44           | 7.2                 | 76    | 6.9 | 13            | 3.9             |               | 2,300                          |         |
|                       | 10/23 |                                      | 12                   | 1.3                         | 20          | 26           | 38           | 6.2                 | 67    | 6.9 | 13            | 5.6             |               | 1,500                          |         |
|                       | 11/7  |                                      | 21                   | 0.77                        | 12          | 9            | 74           | 7.7                 | 71    | 6.7 | 8.9           | 4.7             |               | 2,300                          |         |
|                       | 12/5  |                                      | 10                   | 0.56                        | 13          | 2            | 43           | 7.5                 | 71    | 6.4 | 8.3           | 3.7             |               | 430                            |         |
|                       | 12/20 |                                      | 14                   | 0.88                        | 6           | 3            | 58           | 10.3                | 82    | 6.6 | 12            | 4.5             |               | 430                            |         |
| 1952                  |       |                                      |                      |                             |             |              |              |                     |       |     |               |                 |               |                                |         |
|                       | 1/23  |                                      | 9.4                  | 0.73                        | 13          | 4            | 72           | 8.2                 | 78    | 6.7 | 11            | 4.1             |               | 930                            |         |
|                       | 2/18  |                                      | 14                   | 0.62                        | 10          | 13           | 110          | 9.3                 | 82    | 6.6 | 8.2           | 4.0             |               | 430                            |         |
| S 162.0               | 1951  |                                      |                      |                             |             |              |              |                     |       |     |               |                 |               |                                |         |
|                       | 8/6   | 4,480                                | 7.7                  | 0.26                        | 21          | 58           | 25           | 6.0                 | --    | 7.2 | 20            | 3.3             | 15,000        |                                | 110,000 |
|                       | 8/6   | 4,480                                | 6.9                  | 0.39                        | 21          | 54           |              | 6.1                 | --    | 7.1 | 19            | 3.3             |               |                                |         |
|                       | 8/20  | 4,480                                | 4.0                  | 0.42                        | 31          | 41           | 15           | 6.4                 | 86    | 7.3 | 18            | 3.3             | 39,000        |                                | 9,300   |
|                       | 8/20  | 4,480                                | 5.4                  | 0.77                        | 31          | 42           |              | 6.7                 | 30    | 7.3 | 18            | 3.4             | 43,000        |                                |         |
|                       | 9/4   | 2,720                                | 5.4                  | 0.76                        | 29          | 38           | 15           | 7.1                 | 91    | 7.3 | 18            | 3.8             |               |                                |         |
|                       | 9/4   | 2,720                                | 8.4                  | 0.55                        | 29          | 40           | 15           | 7.0                 | 91    | 7.4 | 17            | 3.6             | 93,000        |                                | 23,000  |
|                       | 9/17  | 4,300                                | 15                   | 0.89                        | 24          | 43           | 35           | 6.4                 | 75    | 7.3 | 16            | 4.1             |               |                                | 75,000  |
|                       | 9/17  | 4,300                                | 15                   | 1.0                         | 24          | 42           | 35           | 6.5                 | 77    | 7.3 | 15            | 3.2             |               |                                |         |
| 1952                  |       |                                      |                      |                             |             |              |              |                     |       |     |               |                 |               |                                |         |
| S 157.0               | 8/6   | 4,480                                | 6.8                  | 0.52                        | --          | 56           | 38           | 6.2                 | --    | 7.1 | 19            | 3.2             | 46,000        |                                |         |
|                       | 8/6   | 4,480                                | 7.8                  | 0.27                        | --          | 60           |              | 6.2                 | --    | 7.2 | 19            | 3.0             |               |                                |         |
|                       | 8/20  | 4,480                                | 4.9                  | 0.45                        | 31          | 39           | 15           | 6.5                 | 87    | 7.3 | 18            | 3.6             |               | 7,500                          |         |
|                       | 9/4   | 2,720                                | 4.7                  | 0.76                        | 29          | 40           | 15           | 7.0                 | 91    | 7.3 | 17            | 3.9             |               | 43,000                         |         |
|                       | 9/17  | 4,300                                | 15                   | 0.62                        | 24          | 45           | 35           | 6.2                 | 73    | 7.4 | 16            | 3.4             |               | 93,000                         |         |

\* Fourmile Branch, 0.01 mile above confluence.



**SAVANNAH RIVER STUDIES**  
**Summary of Individual Results - Sanitary Examinations**

| Sampling Point      | Date         | Disch.<br>at Bur-<br>tons Fy.<br>Sec-Ft | O.C.<br>Moore<br>ppm | 5-Day<br>20°C<br>BOD<br>ppm | Temp.<br>°C | Turb.<br>ppm | Color<br>ppm | Dissolved<br>Oxygen |       | pH  | Alkal.<br>ppm | Chloride<br>ppm | Coliforms<br>MPN per<br>Ga. 100 ml | S.C.   |
|---------------------|--------------|-----------------------------------------|----------------------|-----------------------------|-------------|--------------|--------------|---------------------|-------|-----|---------------|-----------------|------------------------------------|--------|
|                     |              |                                         |                      |                             |             |              |              | ppm                 | % Sat |     |               |                 |                                    |        |
| S 155.2:<br>St O.OI | 1951<br>8/20 |                                         | 5.1                  | 0.50                        | 27          | <25          | 25           | 5.8                 | 72    | 7.3 | 18            | 3.4             | 4,300                              |        |
|                     | 9/4          |                                         | 6.9                  | 0.44                        | 26          | <25          | 25           | 6.8                 | 82    | 7.5 | 35            | 4.1             | 4,300                              |        |
|                     | 9/18         |                                         | 16                   | 2.6                         | 21          | <25          | 48           | 7.9                 | 88    | 7.6 | 25            | 4.6             | 2,300                              |        |
|                     | 9/25         |                                         | 12                   | 0.84                        | 22          | 8            | 40           | 7.3                 | 83    | 7.6 | 26            | 4.5             | 7,500                              |        |
|                     | 10/3         |                                         | 5.1                  | 0.31                        | 21          | 7            | 30           | 6.6                 | 74    | 7.5 | 33            | 4.4             | 930                                |        |
|                     | 10/10        |                                         | 6.1                  | 0.52                        | 21          | 12           | 24           | 8.8                 | 97    | 7.5 | 32            | 3.6             | 210                                |        |
|                     | 10/23        |                                         | 23                   | 1.2                         | 21          | 8            | 72           | 6.7                 | 74    | 7.2 | 21            | 8.1             | 930                                |        |
|                     | 11/7         |                                         | 14                   | 0.57                        | 14          | 5            | 44           | 7.7                 | 74    | 7.0 | 22            | 5.8             | 430                                |        |
|                     | 12/3         |                                         | 18                   | 0.93                        | 11          | 13           | 78           | 8.6                 | 78    | 6.7 | 15            | 5.0             | 1,500                              |        |
|                     | 12/20        |                                         | 21                   | 0.85                        | 5           | 7            | 112          | 9.3                 | 73    | 6.5 | 13            | 5.3             | 2,400                              |        |
|                     | 1952<br>1/23 |                                         | 22                   | 0.80                        | 12          | 3            | 172          | 6.8                 | 63    | 6.9 | 13            | 4.6             | 1,500                              |        |
|                     | 2/6          |                                         | 16                   | 0.65                        | 10          | 3            | 160          | 7.3                 | 64    | 7.0 | 14            | 5.0             | 430                                |        |
| S 152.0             | 2/18         |                                         | 20                   | 0.79                        | 9           | 3            | 150          | 8.5                 | 73    | 6.8 | 13            | 4.5             | 930                                |        |
|                     | 1951<br>8/6  | 4,480                                   | 5.8                  | 0.39                        | --          | 58           | 40           | 6.4                 | --    | 7.2 | 19            | 3.5             | 46,000                             | 46,000 |
|                     | 8/6          | 4,480                                   | 6.1                  | 0.53                        | --          | 56           |              | 6.4                 | --    | 7.2 | 19            | 3.2             | 4,300                              | 15,000 |
|                     | 8/20         | 4,480                                   | 3.5                  | 0.70                        | 31          | 38           | 15           | 6.6                 | 88    | 7.4 | 18            | 3.3             |                                    |        |
|                     | 9/4          | 2,720                                   | 8.2                  | 0.79                        | 29          | 46           | 15           | 7.2                 | 92    | 7.3 | 19            | 4.1             |                                    |        |
|                     | 9/17         | 4,300                                   | 12                   | 0.92                        | 24          | 60           | 30           | 6.5                 | 77    | 7.4 | 16            | 3.5             | 9,300                              |        |
|                     | 9/17         | 4,300                                   | 14                   | 0.93                        | 24          | 56           | 30           | 6.6                 | 78    | 7.4 | 16            | 3.5             |                                    |        |
|                     | 10/3         | 5,560                                   | 8.1                  | 0.49                        | 23          | 155          | 40           | 7.3                 | 84    | 7.4 | 18            | 3.7             | 75,000                             | 23,000 |
|                     | 10/10        | 3,230                                   | 7.2                  | 0.32                        | 21          | 82           | 29           | 7.8                 | 86    | 7.3 | 16            | 3.9             | 43,000                             | 23,000 |
|                     | 10/23        | 2,640                                   | 6.2                  | 0.57                        | 21          | 24           | 25           | 7.6                 | 87    | 7.4 | 23            | 5.2             | 24,000                             | 15,000 |
|                     | 11/7         | 6,280                                   | 6.0                  | 0.39                        | 14          | 29           | 33           | 9.3                 | 90    | 7.2 | 18            | 3.9             | 9,300                              | 43,000 |
|                     | 12/3         | 4,120                                   | 6.7                  | 0.50                        | 11          | 27           | 36           | 10.3                | 93    | 7.1 | 17            | 3.8             | 15,000                             | 15,000 |
|                     | 12/20        | 9,300                                   | 7.0                  | 0.78                        | 6           | 64           | 44           | 11.6                | 93    | 7.0 | 16            | 4.0             | 93,000                             | 24,000 |

\* Steel Creek, 50 ft. above confluence.

SAVANNAH RIVER STUDIES  
Summary of Individual Results - Sanitary Examinations

| Sampling Point           | Date        | Disch.<br>at Bur-<br>tons Ft<br>Sec-Ft | O.O.<br>Moore<br>ppm | 5-Day<br>20°C<br>BOD<br>ppm | Temp.<br>°C | Turb.<br>ppm | Color<br>ppm | Dissolved<br>Oxygen |       | pH  | Alkal.<br>ppm | Chloride<br>ppm | Coliforms<br>MPN per<br>100 ml |        |
|--------------------------|-------------|----------------------------------------|----------------------|-----------------------------|-------------|--------------|--------------|---------------------|-------|-----|---------------|-----------------|--------------------------------|--------|
|                          |             |                                        |                      |                             |             |              |              | ppm                 | % Sat |     |               |                 | Ga.                            | S.O.   |
| S 152.0 M.S.<br>(Cont'd) | 1952<br>1/3 | 12,800                                 | 13                   | 0.68                        | 10          | 165          | 144          | 11.4                | 100   | 7.0 | 7             | 3.4             | 9,300                          | 930    |
|                          | 1/9         | 12,400                                 | 11                   | 0.69                        | 12          | 90           | 108          | 11.2                | 103   | 7.0 | 9             | 3.7             | 4,300                          | 15,000 |
|                          | 1/15        | 10,900                                 | 10                   | 0.66                        | 11          | 75           | 104          | 10.8                | 98    | 7.0 | 12            | 3.8             | 9,300                          | 9,300  |
|                          | 1/23        | 6,550                                  | 9.3                  | 0.38                        | 11          | 53           | 88           | 9.7                 | 88    | 6.9 | 13            | 4.0             | 15,000                         | 23,000 |
|                          | 2/6         | 9,200                                  | 10                   | 0.82                        | 11          | 80           | 120          | 10.0                | 90    | 7.2 | 13            | 4.6             | 9,300                          | 43,000 |
|                          | 2/18        | 8,330                                  | 10                   | 1.4                         | 11          | 42           | 100          | 10.3                | 93    | 7.0 | 15            | 4.1             | 9,300                          | 23,000 |
| S 142.0 M.S.             | 4/3         | 33,000                                 | 8.2                  | 0.46                        | 15          | 95           | 230          | 8.1                 | 80    | 7.0 | 11            | 3.2             | 750                            | 2,300  |
|                          | 1951<br>8/6 | 4,480                                  | 5.3                  | 0.40                        | --          | 59           | 32           | 6.4                 | --    | 7.2 | 19            | 3.2             | 9,300                          | 9,300  |
|                          | 8/20        | 4,480                                  | 4.4                  | 0.52                        | 31          | 37           | 15           | 6.6                 | 88    | 7.3 | 18            | 3.3             | 4,300                          | 4,300  |
|                          | 9/4         | 2,720                                  | 4.1                  | 0.89                        | 29          | 49           | 15           | 7.3                 | 94    | 7.2 | 18            | 4.1             | 24,000                         | 24,000 |
|                          | 9/17        | 4,300                                  | 12                   | 0.73                        | 24          | 61           | 25           | 6.7                 | 79    | 7.3 | 17            | 3.4             | 9,300                          | 9,300  |
|                          | 8/21        |                                        | 9.5                  | 0.40                        | 24          | <25          | 20           | 7.0                 | 82    | 7.4 | 48            | 2.7             | 360                            | 360    |
| *S 137.32;<br>LWR 0.01   | 9/18        |                                        | 19                   | 0.54                        | 21          | <25          | 40           | 7.6                 | 24    | 7.8 | 41            | 4.1             | 11,000                         | 11,000 |
|                          | 9/25        |                                        | 31                   | 1.4                         | 22          | 7            | 80           | 6.2                 | 70    | 7.4 | 26            | 4.9             | 2,300                          | 2,300  |
|                          | 10/4        |                                        | 4.7                  | 0.21                        | 20          | 2            | 25           | 7.7                 | 84    | 8.0 | 47            | 3.9             | 2,300                          | 2,300  |
|                          | 10/11       |                                        | 7.2                  | 0.20                        | 16          | 1            | 19           | 8.8                 | 94    | 7.8 | 48            | 4.2             | 2,300                          | 2,300  |
|                          | 10/18       |                                        | 6.7                  | 0.19                        | 17          | 2            | 18           | 8.2                 | 84    | 7.7 | 47            | 4.0             | 430                            | 430    |
|                          | 10/25       |                                        | 21                   | 0.76                        | 20          | 2            | 55           | 6.6                 | 72    | 7.4 | 37            | 3.3             | 2,400                          | 2,400  |
|                          | 11/1        |                                        | 17                   | 0.61                        | 20          | 2            | 48           | 6.6                 | 72    | 7.6 | 42            | 5.2             | 2,100                          | 2,100  |
|                          | 12/3        |                                        | 9.8                  | 0.59                        | 11          | 2            | 26           | 6.8                 | 61    | 7.5 | 54            | 4.8             | 430                            | 430    |
|                          | 1952<br>1/4 |                                        | 14                   | 0.37                        | 14          | 1            | 54           | 8.2                 | 79    | 7.6 | 36            | 4.2             | 430                            | 430    |
|                          | 1/9         |                                        | 12                   | 0.31                        | 11          | 1            | 38           | 9.7                 | 88    | 7.3 | 34            | 3.8             | 240                            | 240    |
|                          | 2/6         |                                        | 21                   | 1.3                         | 10          | 2            | 88           | 10.1                | 89    | 7.4 | 25            | 4.6             | 230                            | 230    |
|                          | 3/13        |                                        | 21                   | 0.69                        | 19          | 2            | 80           | 7.7                 | 82    | 7.2 | 13            | 4.3             | 430                            | 430    |

\* Lower Three Runs, 50 ft. above confluence.

**SAVANNAH RIVER STUDIES**  
**Summary of Individual Results - Sanitary Examinations**

| Sampling Point         | Date        | Disch.<br>at Bur-<br>tons Fy.<br>Sec-Ft | O.C.<br>Moore<br>ppm | 5-Day<br>20°C<br>BOD<br>ppm | Temp.<br>°C | Turb.<br>ppm | Color<br>ppm | Dissolved<br>Oxygen |       | pH  | Alkal.<br>ppm | Chloride<br>ppm | Coliforms<br>MPN per<br>Ga. 100 ml |
|------------------------|-------------|-----------------------------------------|----------------------|-----------------------------|-------------|--------------|--------------|---------------------|-------|-----|---------------|-----------------|------------------------------------|
|                        |             |                                         |                      |                             |             |              |              | ppm                 | % Sat |     |               |                 |                                    |
| S 132.0 M.S.           | 1951<br>8/6 | 4,480                                   | 5.3                  | 0.25                        | --          | 58           | 37           | 6.4                 | --    | 7.3 | 19            | 3.5             | 9,300                              |
| (1) S 129.25 M.S.      | 1952<br>1/4 | 13,100                                  | 12                   | 0.94                        | 10          | 160          | 132          | 11.0                | 97    | 7.0 | 8.6           | 3.0             | 3,900                              |
| M.S.                   | 1/9         | 12,400                                  | 14                   | 0.52                        | 11          | 125          | 136          | 11.3                | 102   | 6.9 | 0.71          | 3.4             | 9,300                              |
| M.S.                   | 2/6         | 9,200                                   | 12                   | 1.51                        | 11          | 83           | 100          | 10.0                | 90    | 7.2 | 14            | 4.2             | 9,300                              |
| M.S.                   | 3/13        | 34,400                                  | 11                   | 0.91                        | 14          | 92           | 160          | 8.3                 | 80    | 6.9 | 13            | 3.2             | 430                                |
| S 122.0 M.S.           | 1951<br>8/6 | 4,480                                   | 4.8                  | 0.85                        | --          | 59           | 35           | 6.6                 | --    | 7.2 | 19            | 3.4             | 7,500                              |
| M.S.                   | 8/21        | 4,660                                   | 8.1                  | 0.34                        | 29          | 36           | 15           | 6.9                 | 88    | 7.4 | 21            | 3.2             | 4,300                              |
| M.S.                   | 9/5         | 3,140                                   | 5.8                  | 0.66                        | 29          | 43           | 15           | 7.0                 | 90    | 7.5 | 20            | 4.1             | 3,900                              |
| M.S.                   | 9/19        | 4,840                                   | 8.8                  | 0.29                        | 23          | 65           | 30           | 6.8                 | 78    | 7.3 | 17            | 3.5             | 9,300                              |
| S 112.0 M.S.           | 8/8         | 4,480                                   | 4.9                  | 0.60                        | --          | 53           | 15           | 6.7                 | --    | 7.3 | 20            | 3.5             | 1,100,000                          |
| M.S.                   | 8/21        | 4,660                                   | 7.5                  | 0.62                        | 30          | 34           | 15           | 6.9                 | 90    | 7.4 | 20            | 3.2             | 3,900                              |
| M.S.                   | 9/5         | 3,140                                   | 5.2                  | 0.69                        | 29          | 47           | 15           | 7.3                 | 93    | 7.5 | 21            | 3.9             | 9,300                              |
| M.S.                   | 9/19        | 4,840                                   | 6.6                  | 0.39                        | 23          | 56           | 30           | 6.8                 | 79    | 7.3 | 17            | 3.8             | 4,300                              |
| (2) S 105.0;<br>Br 4.0 | 8/9         |                                         | 5.5                  | 0.24                        | 27          | <25          | 30           | 7.0                 | 86    | 7.5 | 42            | 3.2             | 430                                |
| (3) S 105.0;<br>Br 0.5 | 8/21        |                                         | 8.1                  | 1.5                         | 32          | 48           | 15           | 4.1                 | 56    | 7.1 | 23            | 3.4             | 36                                 |
|                        | 9/5         |                                         | 5.6                  | 0.21                        | 26          | <25          | 20           | 7.5                 | 92    | 7.7 | 39            | 3.6             | 930                                |
|                        | 9/19        |                                         | 6.9                  | 0.15                        | 21          | <25          | 30           | 7.5                 | 84    | 7.6 | 34            | 4.1             | 930                                |
| (4) S 105.0;<br>Br 1.2 | 8/9         |                                         | <0.10                | 0.06                        | 19          | <25          | <5           | 5.1                 | 55    | 7.6 | 113           | 4.7             | 36                                 |

(1) Bridge on Ga. Highway 301, known as Burtons Ferry Bridge.

(2) Brier Creek at Brannons Bridge.

(3) Brier Creek 0.5 mile above confluence.

(4) Blue Springs flows into Beaverdam Creek.

SAVANNAH RIVER STUDIES  
Summary of Individual Results - Sanitary Examinations

| Sampling Point | Date | Disch.<br>at Bur-<br>tons Ft.<br>Sec-Ft | O.C.<br>Moore<br>ppm | 5-Day<br>20°C<br>BOD<br>ppm | Temp.<br>°C | Turb.<br>ppm | Color<br>ppm | Dissolved<br>Oxygen |       | pH | Alkal.<br>ppm | Chloride<br>ppm | Coliforms<br>MPN per<br>100 ml |      |
|----------------|------|-----------------------------------------|----------------------|-----------------------------|-------------|--------------|--------------|---------------------|-------|----|---------------|-----------------|--------------------------------|------|
|                |      |                                         |                      |                             |             |              |              | ppm                 | % Sat |    |               |                 | Ga.                            | S.O. |
| 1951           |      |                                         |                      |                             |             |              |              |                     |       |    |               |                 |                                |      |
| S 102.0        | M.S. | 8/8                                     | 4,480                | 6.4                         | 0.54        | —            | 51           | 15                  | 6.8   | —  | 7.3           | 20              | 24,000                         |      |
|                | M.S. | 8/21                                    | 4,660                | 10.9                        | 0.96        | 30           | 33           | 20                  | 7.1   | 93 | 7.2           | 20              | 360                            |      |
|                | M.S. | 9/5                                     | 3,140                | 5.2                         | 0.73        | 29           | 44           | 15                  | 7.2   | 93 | 7.6           | 21              | 4,300                          |      |
|                | M.S. | 9/19                                    | 4,840                | 11                          | 0.30        | 23           | 53           | 30                  | 6.7   | 77 | 7.4           | 18              | 9,300                          |      |
| Disch. near    |      |                                         |                      |                             |             |              |              |                     |       |    |               |                 |                                |      |
| Olyo Sec-Ft    |      |                                         |                      |                             |             |              |              |                     |       |    |               |                 |                                |      |
| S 92.0         | M.S. | 8/8                                     | 4,950                | 7.2                         | 0.21        | —            | 53           | 15                  | 6.4   | —  | 7.3           | 20              | 9,300                          |      |
|                | M.S. | 8/21                                    | 5,040                | 9.8                         | 0.79        | 29           | 33           | 20                  | 6.9   | 89 | 7.3           | 20              | 910                            |      |
|                | M.S. | 9/5                                     | 3,210                | 11                          | 0.61        | 29           | 46           | 15                  | 7.1   | 91 | 7.6           | 22              | 4,300                          |      |
|                | M.S. | 9/19                                    | 4,770                | 9.1                         | 0.47        | 23           | 52           | 30                  | 6.9   | 80 | 7.5           | 18              | 9,300                          |      |
| S 82.0         | M.S. | 8/8                                     | 4,950                | 4.3                         | 0.04        | —            | 55           | 15                  | 6.5   | —  | 7.3           | 20              | 9,300                          |      |
| S 72.0         | M.S. | 8/8                                     | 4,950                | 4.0                         | 0.46        | —            | 54           | 15                  | 6.7   | —  | 7.4           | 20              | 15,000                         |      |
|                | M.S. | 8/21                                    | 5,040                | 9.9                         | 1.0         | 30           | 38           | 20                  | 7.3   | 95 | 7.3           | 21              | 1,500                          |      |
|                | M.S. | 9/5                                     | 3,210                | 9.2                         | 0.72        | 29           | 57           | 15                  | 7.3   | 94 | 7.6           | 25              | 930                            |      |
|                | M.S. | 9/19                                    | 4,770                | 8.4                         | 0.49        | 23           | 60           | 20                  | 7.3   | 84 | 7.5           | 21              | 1,500                          |      |
| S 62.0         | M.S. | 8/8                                     | 4,950                | 6.8                         | 0.43        | 21           | 54           | 15                  | 6.8   | —  | 7.4           | 20              | 15,000                         |      |
| S 52.0         | M.S. | 8/8                                     | 4,950                | 7.6                         | 0.26        | —            | 55           | 15                  | 6.6   | —  | 7.4           | 21              | 910                            |      |
|                | M.S. | 8/21                                    | 5,040                | 8.3                         | 0.64        | 30           | 40           | 20                  | 6.9   | 91 | 7.3           | 21              | 1,500                          |      |
|                | M.S. | 9/5                                     | 3,210                | 4.9                         | 0.36        | 29           | 51           | 15                  | 7.2   | 93 | 7.7           | 25              | 4,300                          |      |
|                | M.S. | 9/19                                    | 4,770                | 5.3                         | 0.24        | 23           | 50           | 15                  | 7.4   | 85 | 7.6           | 23              | 930                            |      |
|                | M.S. | 10/4                                    | 6,260                | 8.8                         | 0.31        | 26           | 122          | 34                  | 7.1   | 87 | 7.6           | 19              | 24,000                         |      |
|                | M.S. | 10/11                                   | 3,870                | 8.0                         | 0.13        | 21           | 108          | 33                  | 7.8   | 87 | 7.5           | 23              | 9,300                          |      |
|                | M.S. | 10/18                                   | 3,780                | 16                          | 0.05        | 21           | 74           | 24                  | 8.3   | 92 | 7.6           | 25              | 24,000                         |      |
|                | M.S. | 10/25                                   | 3,140                | 8.8                         | 0.37        | 23           | 52           | 20                  | 8.2   | 94 | 7.5           | 25              | 2,400                          |      |
|                | M.S. | 11/1                                    | 4,770                | 12                          | 0.31        | 21           | 52           | 26                  | 8.1   | 90 | 7.6           | 23              | 46,000                         |      |
|                |      |                                         |                      |                             |             |              |              |                     |       |    |               |                 | 9,300                          |      |
|                |      |                                         |                      |                             |             |              |              |                     |       |    |               |                 | 24,000                         |      |
|                |      |                                         |                      |                             |             |              |              |                     |       |    |               |                 | 1,500                          |      |
|                |      |                                         |                      |                             |             |              |              |                     |       |    |               |                 | 9,300                          |      |
|                |      |                                         |                      |                             |             |              |              |                     |       |    |               |                 | 46,000                         |      |
|                |      |                                         |                      |                             |             |              |              |                     |       |    |               |                 | 9,300                          |      |

9,300  
24,000  
2,400  
46,000  
9,300

28

**SAVANNAH RIVER STUDIES**  
**Summary of Individual Results - Sanitary Examinations**

| Sampling Point          | Date                  | Disch.<br>near<br>Clyo<br>Sec-Ft | O.C.<br>Moore<br>ppm | 5-Day<br>BOD<br>20°C<br>ppm | Temp.<br>°C | Turb.<br>ppm | Color<br>ppm | Dissolved<br>Oxygen |       | pH  | Alkal.<br>ppr | Chloride<br>ppm | Ga.<br>100 ml | Coliforms<br>MPN per<br>100 ml | S.C.   |
|-------------------------|-----------------------|----------------------------------|----------------------|-----------------------------|-------------|--------------|--------------|---------------------|-------|-----|---------------|-----------------|---------------|--------------------------------|--------|
|                         |                       |                                  |                      |                             |             |              |              | ppm                 | % Sat |     |               |                 |               |                                |        |
| S 52.0 M.S.<br>(Cont'd) | 1951<br>12/6          | 5,220                            | 9.4                  | 0.56                        | 14          | 32           | 37           | 9.8                 | 95    | 7.4 | 21            | 5.0             | 930           |                                | 750    |
|                         | 1952<br>1/31          | 7,820                            | 9.8                  | 0.61                        | 10          | 60           | 80           | 9.8                 | 86    | 6.9 | 20            | 4.6             | 4,300         |                                | 24,000 |
|                         | M.S.<br>2/20          | 11,200                           | 17                   | 1.9                         | 11          | 49           | 96           | 9.6                 | 87    | 7.1 | 14            | 16              | 4,300         |                                | 4,300  |
|                         | M.S.<br>3/12          | 21,400                           | 20                   | 0.97                        | 14          | 75           | 160          | 7.8                 | 75    | 7.1 | 14            | 5.2             | 750           | 430                            | 750    |
| S 48.2;<br>Eb 10.0(1)   | 1951<br>4/9           | 36,500                           | 14                   | 0.67                        | 16          | 55           | 160          | 8.1                 | 81    | 7.1 | 17            | 3.9             | 430           | 430                            | 93     |
|                         | 1951<br>8/9           |                                  | 71                   | 1.8                         | 26          | <25          | 275          | 4.8                 | 58    | 5.7 | 10            | 5.6             |               | 9,300                          |        |
|                         | S 48.2;<br>Eb 4.3(2)  |                                  | 77                   | 1.9                         | 26          | <25          | 300          | 3.2                 | 39    | 5.6 | 10            | 5.6             |               | 2,300                          |        |
|                         | S 48.2;<br>Eb 0.25(3) |                                  |                      |                             |             |              |              |                     |       |     |               |                 |               |                                |        |
|                         | 8/21                  |                                  | 87                   | 1.2                         | 29          | <25          | 300          | 2.1                 | 27    | 5.2 | 9             | 2.6             |               | 730                            |        |
|                         | 9/5                   |                                  | 22                   | 0.69                        | 31          | 64           | 100          | 4.8                 | 64    | 7.3 | 20            | 3.6             |               |                                |        |
|                         | 9/19                  |                                  | 6.9                  | 0.62                        | 21          | 31           | 35           | 6.9                 | 76    | 7.4 | 22            | 3.3             |               | 430                            |        |
|                         | 10/4                  |                                  | 49                   | 0.81                        | 24          | 12           | 132          | 2.4                 | 28    | 7.2 | 15            | 32              |               | 930                            |        |
|                         | 1952<br>1/31          |                                  | 52                   | 0.87                        | 10          | 2            | 230          | 3.0                 | 26    | 5.0 | 6.4           | 7.9             |               | 43                             |        |
|                         | 2/20                  |                                  | 68                   | 2.2                         | 13          | 2            | 300          | 5.5                 | 52    | 5.6 | 4.9           | 6.8             |               | 2,400                          |        |
|                         | S 42.0 M.S.           | 8/8                              | 4,950                | 7.5                         | 0.25        | 22           | 53           | 15                  | 6.5   | 7.4 | 22            | 3.4             |               | 910                            |        |
|                         | S 32.0 M.S.           | 8/8                              | 4,950                | 6.3                         | 0.42        |              | 48           | 18                  | 6.5   | 7.4 | 22            | 3.5             |               | 910                            |        |
|                         | M.S.<br>8/21          | 5,040                            | 8.7                  | 1.09                        | 31          | 32           | 30           | 6.9                 | 92    | 7.2 | 22            | 3.3             |               | 430                            |        |

- (1) Ebenezer Creek at bridge on Ga. Highway 119.  
(2) Ebenezer Creek at bridge on road nearest Savannah River.  
(3) Ebenezer Creek, 0.25 mile above confluence (boat sampling)

SAVANNAH RIVER STUDIES  
Summary of Individual Results - Sanitary Examinations

| Sampling Point | Date  | Disch.<br>near<br>Glyo<br>Sec-Ft | O.C.<br>Moore<br>ppm | 5-Day<br>20°C<br>BOD<br>ppm | Temp.<br>°C | Turb.<br>ppm | Color<br>ppm | Dissolved<br>Oxygen |       | pH  | Alkal.<br>ppm | Chloride<br>ppm | Ga.<br>100 ml | Coliforms<br>MPN per<br>100 ml | S.C.   |
|----------------|-------|----------------------------------|----------------------|-----------------------------|-------------|--------------|--------------|---------------------|-------|-----|---------------|-----------------|---------------|--------------------------------|--------|
|                |       |                                  |                      |                             |             |              |              | ppm                 | % Sat |     |               |                 |               |                                |        |
| S 11.0         | 1951  |                                  |                      |                             |             |              |              |                     |       |     |               |                 |               |                                |        |
|                | 10/4  | 5,310                            |                      | 0.69                        | 25          | 49           | 15           | 4.0                 | 51    | 8.0 | 67            | 7,890           | 7,500         |                                |        |
|                | 10/4  | 5,310                            |                      | 0.58                        | 25          | 46           | 15           | 3.8                 | 50    | 8.0 | 67            | 7,710           | 23,000        |                                |        |
|                | 10/4  | 5,310                            |                      | 0.56                        | 25          | 50           | 15           | 3.6                 | 46    | 8.0 | 67            | 7,450           |               |                                |        |
|                | 10/11 | 3,780                            |                      | 1.0                         | 23          | 74           | 19           | 0.0                 | 0     | 7.6 | 47            | 3,040           | 43,000        | 4,300                          |        |
|                | 10/11 | 3,780                            |                      | 1.1                         | 23          | 70           | 18           | 0.0                 | 0     | 7.6 | 46            | 2,890           |               |                                |        |
|                | 10/11 | 3,780                            |                      | 1.3                         | 23          | 73           | 19           | 0.68                | 8     | 7.6 | 46            | 3,040           | 43,000        |                                |        |
|                | 10/18 | 3,290                            |                      | 0.51                        | 21          | 31           | 11           | 5.1                 | 63    | 7.6 | 80            | 10,000          | 4,300         | 43,000                         |        |
|                | 10/18 | 3,290                            |                      | 1.0                         | 21          | 33           | 12           | 5.1                 | 62    | 7.6 | 79            | 9,850           |               |                                |        |
|                | 10/18 | 3,290                            |                      | 0.53                        | 21          | 31           | 13           | 5.1                 | 62    | 7.6 | 78            | 9,680           | 2,300         |                                |        |
|                | 10/25 | 3,210                            |                      | 1.0                         | 23          | 14           | 17           | 3.7                 | 45    | 7.2 | 52            | 4,190           | 24,000        | 730                            |        |
|                | 10/25 | 3,210                            |                      | 1.0                         | 23          | 15           | 17           | 4.4                 | 53    | 7.2 | 51            | 3,980           | 15,000        |                                |        |
|                | 10/25 | 3,210                            |                      | 0.98                        | 23          | 14           | 19           | 4.0                 | 48    | 7.2 | 51            | 3,810           |               | 2,400                          |        |
|                | 11/1  | 4,680                            |                      | 1.2                         | 23          | 65           | 13           | 2.7                 | 34    | 7.7 | 77            | 7,680           | 2,400         |                                |        |
|                | 11/1  | 4,680                            |                      | 0.86                        | 23          | 85           | 14           | 3.1                 | 39    | 7.7 | 79            | 8,110           |               |                                |        |
|                | 11/1  | 4,680                            |                      | 1.3                         | 23          | 260          | 14           | 2.8                 | 34    | 7.7 | 81            | 8,320           |               | 4,600                          |        |
|                | 12/6  | 6,760                            |                      | 0.82                        | 14          | 126          | 13           | 2.8                 | 27    | 7.5 | 50            | 2,720           | 4,300         | 9,300                          |        |
|                | 1952  |                                  |                      |                             |             |              |              |                     |       |     |               |                 |               |                                |        |
|                | 1/4   | 14,500                           |                      | 0.78                        | 12          | 120          | 28           | 6.3                 | 58    | 7.4 | 26            | 1,620           | 21,000        | 43,000                         | 9,300  |
|                | 2/20  | 9,900                            |                      | 1.2                         | 13          | 46           | 32           | 6.8                 | 67    | 7.4 | 37            | 3,220           | 23,000        | 23,000                         | 23,000 |
|                | 3/12  | 38,600                           |                      | 1.1                         | 15          | 170          | 32           | 6.2                 | 61    | 7.6 | 29            | 2,490           | 9,300         | 23,000                         | 23,000 |
|                | 4/9   | 36,500                           |                      | 2.8                         | 17          | 190          | 220          | 6.9                 | 71    | 7.2 | 19            | 13              | 4,300         | 4,300                          | 9,300  |



APPENDIX "B"

SUMMARY OF INDIVIDUAL RESULTS

MINERAL EXAMINATIONS

SAVANNAH RIVER STUDIES  
Summary of Individual Results - Mineral Examinations  
(Results in ppm)

| Station   | Date     | Suspended Solids |      | Silica           | Sulfate         | Magnesium | Sol. + Pot. | Total Dis. | Iron  | Fluoride | Manganese | Orthophosphate  | Calcium | Bicarbonate      | Chloride | Iron Al. Oxides | Aluminum |
|-----------|----------|------------------|------|------------------|-----------------|-----------|-------------|------------|-------|----------|-----------|-----------------|---------|------------------|----------|-----------------|----------|
|           |          | Total            | Ash  | SiO <sub>2</sub> | SO <sub>4</sub> | Mg        | Na+K        | Solids     | Fe    | F        | Mn        | PO <sub>4</sub> | Ca      | HCO <sub>3</sub> | Cl       | Fe+Al           | Al       |
| S 218.5   | 12/10/51 | 94.4             | 66.4 | 27.0             | 6.2             | 1.4       | 3.0         | 62.4       | 4.4   | 0.05     | 0.65      | 0.08            | 2.7     | 13               | 3.6      | 24.5            | 9.5      |
|           | 1/8      | 34.0             | 25.2 | 23.2             | 3.6             | 1.4       | 3.9         | 74.8       | 3.6   | 0.10     | 0.10      | 0.02            | 2.6     | 11               | 3.2      | 17.0            | 6.2      |
| S 202.9   | 10/31    | 19.6             | 12.0 | 15.3             | 3.1             | 1.1       | 4.4         | 54.4       | 0.02  | 0.15     | 0.15      | 0.01            | 2.8     | 22               | 4.6      | 3.3             | 1.6      |
|           | 12/18    | 26.0             | 12.8 | 16.9             | 4.9             | 1.7       | 3.7         | 48.4       | 0.80  | 0.18     | 0.80      | 0.07            | 3.2     | 17               | 3.7      | 5.0             | 2.0      |
|           | 1/29     | 30.0             | 22.0 | 19.6             | 2.9             | 1.2       | 3.4         | 64.4       | 2.2   | 0.20     | 1.2       | 0.003           | 4.3     | 14               | 4.3      | 9.2             | 3.2      |
|           | 2/13     | 30.4             | 20.4 | 21.4             | 3.4             | 1.0       | 3.1         | 79.2       | 3.0   | 0.15     | 1.6       | 0.01            | 4.3     | 15               | 3.5      | 13.3            | 4.7      |
|           | 3/13     | 108              | 89.2 | 7.2              | 10.2            | 0.93      | 4.5         | 28.0       | 0.15  | nil      | 0.16      | 0.02            | 3.7     | 13               | 3.3      | 1.2             | 0.49     |
| S 197     | 10/31    | 19.0             | 11.4 | 14.5             | 2.9             | 1.2       | 4.2         | 54         | 0.02  | 0.15     | 0.01      | 0.13            | 3.0     | 21               | 4.7      | 2.9             | 1.4      |
|           | 12/18    | 22.4             | 11.6 | 16.5             | 4.2             | 1.6       | 3.8         | 51.2       | 0.95  | 0.17     | 0.80      | 0.07            | 3.1     | 13               | 3.7      | 5.3             | 1.8      |
|           | 1/29     | 28.8             | 21.2 | 19.3             | 3.3             | 1.1       | 3.2         | 64.8       | 2.6   | 0.20     | 1.2       | 0.004           | 4.4     | 13               | 4.0      | 11.9            | 4.2      |
|           | 2/13     | 36.4             | 26.4 | 21.9             | 3.6             | 1.2       | 3.8         | 72.4       | 2.8   | 0.15     | 1.6       | 0.06            | 4.3     | 16               | 3.6      | 14.9            | 5.7      |
| S 172     | 10/23    | 20.4             | 14.8 | 10.3             | 7.2             | 0.26      | 8.4         | 56         | 0.02  | 0.15     | 0.20      | 0.05            | 4.2     | 21               | 4.8      | 1.8             | 0.80     |
|           | 12/28    | 124              | 106  | 23.6             | 4.0             | 1.6       | 3.8         | 126        | 3.8   | 0.10     | 2.8       | 0.02            | 2.6     | 7.2              | 2.9      | 20.8            | 8.1      |
|           | 1/15     | 48.4             | 40.8 | 19.7             | 4.1             | 1.2       | 4.4         | 72.8       | 2.8   | 0.10     | 1.3       | 0.013           | 2.7     | 12               | 3.3      | 13.6            | 5.1      |
|           | 4/3      | 64.8             | 51.2 | 7.8              | 4.8             | 0.94      | 1.1         | 44.4       | 0.56  | —        | 0.24      | 0.008           | 3.2     | 11               | 3.1      | 0.98            | 0.08     |
| S 171.59: | 10/23    | 5.8              | 1.8  | 7.4              | 1.6             | 0.24      | 5.9         | 40         | 0.02  | 0.15     | 0.07      | 0.07            | 2.3     | 8.9              | 5.8      | 0.5             | 0.14     |
| UTR 0.01  | 12/28    | 4.8              | 3.6  | 12.5             | 3.6             | 1.5       | 1.8         | 56.8       | 0.85  | nil      | 2.4       | 0.001           | 2.6     | 4.9              | 3.3      | 5.4             | 2.1      |
|           | 1/23     | 5.6              | 3.6  | 9.6              | 2.1             | 0.80      | 5.0         | 24.8       | 1.2   | 0.15     | 1.2       | 0.003           | 2.5     | 3.5              | 3.4      | 2.8             | 0.53     |
| S 164.26: | 11/7     | 10.8             | 4.4  | 9.8              | 2.8             | 1.2       | 5.7         | 60.4       | 0.005 | 0.15     | 0.04      | 0.007           | 2.9     | 8.9              | 4.7      | 5.3             | 2.7      |
| Fe 0.01   | 1/23     | 1.6              | nil  | 8.7              | 1.4             | 1.2       | 0.54        | 36.8       | 0.24  | 0.15     | 0.80      | 0.004           | 3.7     | 11               | 4.1      | 2.7             | 1.2      |
|           | 2/18     | 16.0             | 8.0  | 11.7             | 2.1             | 0.58      | 1.7         | 44.4       | 3.5   | nil      | 0.52      | 0.05            | 3.7     | 8.2              | 4.0      | 9.0             | 2.0      |

\* Fusion

**SAVANNAH RIVER STUDIES**  
**Summary of Individual Results Mineral Examinations**  
 (Results in ppm)

| Station              | Date<br>1951<br>1952 | Suspended Solids |      | Silica<br>SiO <sub>2</sub> | Sulfate<br>SO <sub>4</sub> | Magnesium<br>Mg | Sodium + Potassium<br>Na+K | Total Dissolved Solids | Iron<br>Fe | Fluoride<br>F | Manganese<br>Mn | Orthophosphate<br>PO <sub>4</sub> | Total phosphate<br>PO <sub>4</sub> | Calcium<br>Ca | Bicarbonate<br>HCO <sub>3</sub> | Chloride<br>Cl | Iron-Aluminum<br>Fe+Al<br>Oxides | Aluminum<br>Al |
|----------------------|----------------------|------------------|------|----------------------------|----------------------------|-----------------|----------------------------|------------------------|------------|---------------|-----------------|-----------------------------------|------------------------------------|---------------|---------------------------------|----------------|----------------------------------|----------------|
|                      |                      | Total            | Ash  |                            |                            |                 |                            |                        |            |               |                 |                                   |                                    |               |                                 |                |                                  |                |
| S 155.2;<br>St 0.01  | 11/7                 | 9.6              | 2.8  | 10.5                       | 3.3                        | 1.1             | 7.1                        | 60                     | 0.015      | 0.13          | 0.03            | 0.03                              | 0.28                               | 8.1           | 22                              | 5.8            | 2.0                              | 0.93           |
|                      | 2/6                  | 5.6              | 2.8  | 11.6                       | 3.3                        | 0.78            | 1.6                        | 134                    | 3.4        | 0.15          | 0.56            | 0.02                              | 0.02                               | 6.5           | 14                              | 5.0            | 6.0                              | 0.63           |
| S 152                | 12/3                 | 16.0             | 10.4 | 13.4                       | 3.4                        | 1.4             | 2.2                        | 48.8                   | 0.003      | 0.15          | 1.1             | 0.07                              | 0.14                               | 3.2           | 17                              | 3.8            | 2.1                              | 1.0            |
|                      | 1/9                  | 50.8             | 42.0 | 21.8                       | 4.4                        | 1.3             | 4.5                        | 62.4                   | 2.9        | 0.10          | 1.2             | 0.003                             | 0.04                               | 2.6           | 9.2                             | 3.7            | 15.2                             | 5.8            |
|                      | 4/3                  | 42.8             | 34.0 | 7.2                        | 5.1                        | 1.3             | 0.73                       | 34.8                   | 0.69       | 0.05          | 0.24            | 0.01                              | 0.03                               | 3.2           | 11                              | 3.1            | 1.4                              | 0.18           |
| S 140.9;<br>LTS 5.75 | 12/3                 | 3.2              | 11.1 | 14.8                       | 4.6                        | 1.7             | 4.2                        | 87.2                   | 0.013      | 0.12          | 1.2             | 0.04                              | 0.14                               | 17.1          | 54                              | 4.8            | 0.50                             | 0.20           |
|                      | 2/6                  | 0.40             | 11.1 | 12.4                       | 2.1                        | 0.71            | 0.77                       | 128                    | 0.46       | 0.14          | 0.96            | 0.003                             | 0.005                              | 14.5          | 25                              | 4.6            | 2.1                              | 0.73           |
| S 132                | 1/4                  | 102              | 80.8 | 19.1                       | 4.3                        | 1.2             | 3.6                        | 65.2                   | 2.9        | 0.15          | 0.90            | 0.034                             | 0.02                               | 2.5           | 8.6                             | 3.0            | 14.7                             | 5.5            |
| S 129.25             | 3/13                 | 37.2             | 26.0 | 7.6                        | 11.9                       | 0.88            | 5.5                        | 32.8                   | 0.65       | 0.05          | 0.20            | 0.007                             | 0.01                               | 4.2           | 13                              | 3.2            | 1.7                              | 0.40           |
| S 52                 | 12/6                 | 15.6             | 10.4 | 13.6                       | 6.3                        | 1.6             | 3.7                        | 29.2                   | 0.003      | 11.1          | 1.4             | 0.04                              | 0.04                               | 5.1           | 21                              | 5.0            | 3.9                              | 2.0            |
|                      | 2/20                 | 34.0             | 25.6 | 17.6                       | 4.0                        | 1.0             | 2.5                        | 74.0                   | 2.2        | 0.10          | 0.28            | 0.03                              | 0.11                               | 5.6           | 14                              | 16.2           | 9.3                              | 3.2            |
|                      | 4/9                  | 26.8             | 21.6 | —                          | 5.7                        | 0.83            | —                          | 30.4                   | 0.08       | 0.02          | 0.07            | 0.007                             | 0.01                               | 4.1           | 17                              | 3.9            | 0.40                             | 0.14           |
| S 11                 | 12/6                 | 34.8             | 27.6 | 8.3                        | 25.4                       | 1.64            | 659.                       | 3632.                  | 0.28*      | 0.23          | 0.10            | 0.025                             | 0.01                               | 64            | 50                              | 2720.          | 8.3                              | 4.2            |
|                      | 1/4                  | 72.8             | 52.8 | 11.9                       | 21.2                       | 75.4            | 713.                       | 2981.                  | 0.18       | 0.10          | 1.3             | 0.005                             | 0.05                               | 52            | 26                              | 1620.          | 4.7                              | 2.3            |
|                      | 1/18                 | 66.4             | 52.8 | 12.0                       | 152.                       | 71.5            | 1150.                      | 2134.                  | 0.44       | 0.50          | 2.4             | 0.003                             | 0.04                               | 32            | 23                              | 1300.          | 5.6                              | 2.6            |
|                      | 3/12                 | 101              | 80.4 | 28.4                       | 240.                       | 144.            | 1094.                      | 4258.                  | 0.14       | 0.12          | 4.0             | 0.009                             | 0.01                               | 57            | 29                              | 2490.          | 4.2                              | 2.1            |
|                      | 4/9                  | 68.0             | 58.4 | 6.2                        | 8.3                        | 1.2             | —                          | 42.4                   | 0.08       | 11.1          | 0.06            | 0.008                             | 0.01                               | 3.8           | 19                              | 13.            | 0.47                             | 0.18           |

389

APPENDIX "C"

CLARK HILL RESERVOIR

SUMMARY OF INDIVIDUAL RESULTS - IRON, MANGANESE,  
TEMPERATURE, TURBIDITY AND DISSOLVED OXYGEN  
DETERMINATIONS; RESERVOIR DISCHARGE DATA.

(NOT FOR PUBLIC RELEASE)  
SAVANNAH RIVER STUDIES  
ANALYTICAL RESULTS

CLARK HILL RESERVOIR  
S-255  
Ga. Hwy #43 Bridge over Savannah River

| DATE    | DEPTH<br>FT. | MANG-<br>ANISE<br>ppm | IRON<br>ppm | TEMP.<br>DEG.C. | TURB.<br>ppm | DISS.<br>OXYGEN<br>ppm | DIS-<br>CHARGE<br>SEC-FT. | DATE | DEPTH<br>FT. | MANG-<br>ANISE<br>ppm | IRON<br>ppm | TEMP.<br>DEG.C. | TURB.<br>ppm | DISS-<br>OXYGEN<br>ppm | DIS-<br>CHARGE<br>SEC-FT. |
|---------|--------------|-----------------------|-------------|-----------------|--------------|------------------------|---------------------------|------|--------------|-----------------------|-------------|-----------------|--------------|------------------------|---------------------------|
| 4/30/52 | Total<br>35' |                       |             |                 |              |                        | 14,200                    |      |              |                       |             |                 |              |                        |                           |
| "       | 32'          | —                     | —           | 19'             | 90           | 8.2                    |                           |      |              |                       |             |                 |              |                        |                           |
| "       | 10'          | 0                     | 0.04        | 19              | 80           | 8.3                    |                           |      |              |                       |             |                 |              |                        |                           |
| 5/19/52 | Total<br>33' |                       |             |                 |              |                        | 4,050                     |      |              |                       |             |                 |              |                        |                           |
| "       | 30'          | 0.007                 | 0.20        | 24              | 3            | 8.9                    |                           |      |              |                       |             |                 |              |                        |                           |
| "       | 3'           | —                     | —           | 27              | 2            | 9.1                    |                           |      |              |                       |             |                 |              |                        |                           |
| 5/22/52 | Total<br>25' |                       |             |                 |              |                        | 4,080                     |      |              |                       |             |                 |              |                        |                           |
| "       | 23'          | 0.0013                | 0.04        | 24              | 4            | 5.6                    |                           |      |              |                       |             |                 |              |                        |                           |
| "       | 3'           | —                     | —           | 26              | 2            | 7.6                    |                           |      |              |                       |             |                 |              |                        |                           |

Preceding page blank

(NOT FOR PUBLIC RELEASE)

CLARK HILL RESERVOIR  
S-253, L1(S.C.)-3.2  
S.C. Hwy #43 Bridge over Little River, S.C.

SAVANNAH RIVER STUDIES  
ANALYTICAL RESULTS

| DATE    | DEPTH<br>FT. | NAME-<br>ANALYST | IRON<br>PPM | TEMP.<br>DEG.C. | TOTAL<br>PPM | DISS.<br>OXYGEN<br>PPM | DIS-<br>CHARGE<br>SEC.-FT. | DATE | DEPTH<br>FT. | NAME-<br>ANALYST | IRON<br>PPM | TEMP.<br>DEG.C. | TOTAL<br>PPM | DISS.<br>OXYGEN<br>PPM | DIS-<br>CHARGE<br>SEC.-FT. |
|---------|--------------|------------------|-------------|-----------------|--------------|------------------------|----------------------------|------|--------------|------------------|-------------|-----------------|--------------|------------------------|----------------------------|
| 4/30/52 | Total        |                  |             |                 |              |                        | 14,200                     |      |              |                  |             |                 |              |                        |                            |
| "       | 17'          |                  | 0           | 0.068           | 19           | 65                     | 7.6                        |      |              |                  |             |                 |              |                        |                            |
| "       | 5'           |                  | 0           | 0.068           | 20           | 60                     | 7.7                        |      |              |                  |             |                 |              |                        |                            |
| 5/19/52 | Total        |                  |             |                 |              |                        | 4,050                      |      |              |                  |             |                 |              |                        |                            |
| "       | 20           |                  | 0.007       | 0.32            | 23           | 5                      | 6.5                        |      |              |                  |             |                 |              |                        |                            |
| "       | 3'           |                  | —           | —               | 30           | 5                      | 9.5                        |      |              |                  |             |                 |              |                        |                            |
| 5/22/52 | Total        |                  |             |                 |              |                        | 4,080                      |      |              |                  |             |                 |              |                        |                            |
| "       | 15'          |                  | 0.0013      | 0.048           | 21           | 410                    | 6.1                        |      |              |                  |             |                 |              |                        |                            |
| "       | 3'           |                  | —           | —               | 25           | 300                    | 6.2                        |      |              |                  |             |                 |              |                        |                            |



CLARK HILL RESERVOIR  
8 - 246  
8 miles above Dam

| DATE   | DEPTH<br>FT. | MANG-<br>ANISE<br>ppm | IRON<br>ppm | TEMP.<br>DEG.C. | TURB.<br>ppm | DISS.<br>OXYGEN<br>ppm | DIS-<br>CHARGE<br>SEC.-FT. | DATE    | DEPTH<br>FT. | MANG-<br>ANISE<br>ppm | IRON<br>ppm | TEMP.<br>DEG.C. | TURB.<br>ppm | DISS.<br>OXYGEN<br>ppm | DIS-<br>CHARGE<br>SEC.-FT. |
|--------|--------------|-----------------------|-------------|-----------------|--------------|------------------------|----------------------------|---------|--------------|-----------------------|-------------|-----------------|--------------|------------------------|----------------------------|
|        |              |                       |             |                 |              |                        |                            |         |              |                       |             |                 |              |                        |                            |
| 5/6/52 | 030          |                       |             |                 |              |                        |                            | 5/26/52 | 750          |                       |             |                 |              |                        | 4,085                      |
| "      | 80'          | 0.867                 | 0.10        | 18              | 10           | 4.9                    |                            | "       | 72'          | 0.44                  | 0.048       | 21              | 6            | 0.27                   |                            |
| "      | 30'          | --                    | --          | 21              | 37           | 6.9                    |                            | "       | 47'          | 0.30                  | 0.02        | 22              | 41           | 2.6                    |                            |
| "      | 5'           | --                    | --          | 26              | 4            | 8.5                    |                            | "       | 22'          | --                    | --          | 25              | 41           | 7.1                    |                            |
|        |              |                       |             |                 |              |                        |                            | "       | 3'           | --                    | --          | 26              | 1            | 7.6                    |                            |

| 5/13/52 | Total |
|---------|-------|
| 751     | 7,100 |

|         |     |     |       |       |    |   |     |       |
|---------|-----|-----|-------|-------|----|---|-----|-------|
| 5/21/52 | 68° | 65° | 0.005 | 0.048 | 20 | 5 | 4.6 | 4,070 |
| "       | "   | 40° | 0.12  | 0.10  | 21 | 3 | 3.2 |       |
| "       | "   | 15° | —     | —     | 25 | 2 | 7.8 |       |
| "       | "   | 3°  | —     | —     | 26 | 1 | 7.9 |       |

[illegible]

**CLARK HILL RESERVOIR**  
1 mile above Dam - Junction of Little River.

CLARK HILL RESERVOIR

1 mile above Dam - Junction of Little River,  
Georgia with the Savannah River

8-239.1

1

[illegible]

8-239.1

## ANALYTICAL RESULTS

[illegible]

\* Alkalinity 22.9 ppm. Samples collected and analytical determinations made at location by Dr. D. C. Scott and U. of Ga. Biology students.

(NOT FOR PUBLIC RELEASE)

SAVANNAH RIVER STUDIES  
ANALYTICAL RESULTS

CLARK HILL RESERVOIR  
1 Mile above Dam - Junction of Little  
River Georgia with the Savannah River  
9-239.1

| DATE    | DEPTH<br>FT. | MASS-<br>ANALYSIS<br>ppm | IRON<br>ppm | TEMP.<br>DEG. F. | TURB.<br>ppm | DIS-<br>SOLVED<br>OXYGEN<br>ppm | DIS-<br>CHARGE<br>SEC-FT. | DATE | DEPTH<br>FT. | MASS-<br>ANALYSIS<br>ppm | IRON<br>ppm | TEMP.<br>DEG. F. | TURB.<br>ppm | DIS-<br>SOLVED<br>OXYGEN<br>ppm | DIS-<br>CHARGE<br>SEC-FT. |
|---------|--------------|--------------------------|-------------|------------------|--------------|---------------------------------|---------------------------|------|--------------|--------------------------|-------------|------------------|--------------|---------------------------------|---------------------------|
| 5/21/52 | Total        | 110                      |             |                  |              |                                 | 4,070                     |      |              |                          |             |                  |              |                                 |                           |
| "       | 107'         | 0.80                     | 0.08        | 17               | 30           | 0                               |                           |      |              |                          |             |                  |              |                                 |                           |
| "       | 82'          | 0.0013                   | 0.14        | 21               | 7            | 3.0                             |                           |      |              |                          |             |                  |              |                                 |                           |
| "       | 57'          | 0.0013                   | 0.06        | 20               | 3            | 4.6                             |                           |      |              |                          |             |                  |              |                                 |                           |
| "       | 32'          | —                        | —           | 22               | 3            | 6.5                             |                           |      |              |                          |             |                  |              |                                 |                           |
| "       | 3'           | —                        | —           | 25               | 4            | 7.9                             |                           |      |              |                          |             |                  |              |                                 |                           |

|         |       |       |       |    |    |     |       |  |  |  |  |  |  |  |  |
|---------|-------|-------|-------|----|----|-----|-------|--|--|--|--|--|--|--|--|
| 5/26/52 | Total | 115   |       |    |    |     | 4,085 |  |  |  |  |  |  |  |  |
| "       | 113'  | 1.12  | 0.12  | 19 | 15 | 0   |       |  |  |  |  |  |  |  |  |
| "       | 88'   | 0.24  | 0.032 | 20 | 4  | 2.0 |       |  |  |  |  |  |  |  |  |
| "       | 63'   | 0.033 | 0.08  | 21 | 4  | 4.0 |       |  |  |  |  |  |  |  |  |
| "       | 38'   | 0.024 | 0.04  | 22 | 2  | 4.9 |       |  |  |  |  |  |  |  |  |
| "       | 13'   | —     | —     | 24 | 1  | 7.4 |       |  |  |  |  |  |  |  |  |
| "       | 3'    | —     | —     | 25 | 1  | 7.7 |       |  |  |  |  |  |  |  |  |

(NOT FOR PUBLIC RELEASE)  
SAVANNAH RIVER STUDIES  
ANALYTICAL RESULTS

CLARK HILL RESERVOIR  
One Mile Above Dam-Junction of Little  
River, Georgia with the Savannah River

| DATE    | DEPTH<br>FT. | MANG-<br>ANISE<br>PPM | IRON<br>PPM | TEMP.<br>DEG.C. | TURB.<br>PPM | DISS.<br>OXYGEN<br>PPM | DISS-<br>CHARGE<br>SEC.FT. | DATE    | DEPTH<br>FT. | MANG-<br>ANISE<br>PPM | IRON<br>PPM | TEMP.<br>DEG.C. | TURB.<br>PPM | DISS.<br>OXYGEN<br>PPM | DISS-<br>CHARGE<br>SEC.FT. |
|---------|--------------|-----------------------|-------------|-----------------|--------------|------------------------|----------------------------|---------|--------------|-----------------------|-------------|-----------------|--------------|------------------------|----------------------------|
| 6/13/52 | 105          |                       |             | 16.4            | —            | —                      | 4.120                      | 7/28/52 | 110          |                       |             | 20.0            |              | 0 <sup>0</sup>         | 3.510                      |
|         | 100          |                       |             | 16.8            | —            | 0.14                   |                            |         | 100          |                       |             | 20.3            |              | 0 <sup>0</sup>         |                            |
|         | 90           |                       |             | 17.5            | <10          | 0.97                   |                            |         | 90           |                       |             | 20.6            |              | 0 <sup>0</sup>         |                            |
|         | 80           |                       |             | 18.5            | <10          | 4.2                    |                            |         | 80           |                       |             | 21.0            |              | 0 <sup>0</sup>         |                            |
|         | 70           |                       |             | 18.9            | <10          | 4.3                    |                            |         | 70           |                       |             | 21.6            |              | 0 <sup>0</sup>         |                            |
|         | 60           |                       |             | 19.6            | <10          | 3.0                    |                            |         | 60           |                       |             | 22.5            |              | 0 <sup>0</sup>         |                            |
|         | 50           |                       |             | 20.0            | <10          | 3.7                    |                            |         | 50           |                       |             | 23.9            |              | 0 <sup>0</sup>         |                            |
|         | 40           |                       |             | 20.8            | <10          | 2.7, 2.2               |                            |         | 40           |                       |             | 25.8            |              | 0 <sup>0</sup>         |                            |
|         | 35           |                       |             | 21.5            | <10          | —                      |                            |         | 30           |                       |             | 27.3            | 1.0          |                        |                            |
|         | 30           |                       |             | 22.1            | <10          | 3.2, 2.9               |                            |         | 20           |                       |             | 28.6            | 6.4          |                        |                            |
|         | 25           |                       |             | 23.3            | <10          | 4.3                    |                            |         | 10           |                       |             | 30.4            | 7.3          |                        |                            |
|         | 20           |                       |             | 27.8            | <10          | 7.6                    |                            |         | 0            |                       |             | 31.7            | 7.0          |                        |                            |
|         | 15           |                       |             | 28.1            | <10          | 7.6                    |                            |         |              |                       |             |                 |              |                        |                            |
|         | 10           |                       |             | 28.3            | <10          | 8.2                    |                            |         |              |                       |             |                 |              |                        |                            |
|         | 5            |                       |             | 28.6            | <10          | 8.2                    |                            |         |              |                       |             |                 |              |                        |                            |
|         | 0            |                       |             | 29.0            | <10          | 8.1                    |                            |         |              |                       |             |                 |              |                        |                            |

\*Noticeable odor of H<sub>2</sub>S present. The water leaving the reservoir gates had a distinct H<sub>2</sub>S odor which was noticeable 2 miles upwind.

Noticeable odor of H<sub>2</sub>S present. The water leaving the reservoir gates had a distinct H<sub>2</sub>S odor which was noticeable 2 miles upwind.

#samples collected and determinations made at location



PROVISIONAL DATA  
(NOT FOR PUBLIC RELEASE)  
SAVANNAH RIVER STUDIES  
ANALYTICAL RESULTS

CLARK HILL RESERVOIR

8-237.8

0.3 Mile below Clark Hill Dam

Sample taken from construction bridge below the dam

| DATE    | DEPTH<br>FT. | MANG-<br>ANES | IRON<br>PPM | TEMP.<br>DEG.C. | TURB.<br>PPM | DISS.<br>OXYGEN<br>PPM | DIS-<br>CHARGE<br>SEC-FT. | DATE | DEPTH<br>FT. | MANG-<br>ANES<br>PPM | IRON<br>PPM | TEMP.<br>DEG.C. | TURB.<br>PPM | DISS.<br>OXYGEN<br>PPM | DIS-<br>CHARGE<br>SEC-FT. |
|---------|--------------|---------------|-------------|-----------------|--------------|------------------------|---------------------------|------|--------------|----------------------|-------------|-----------------|--------------|------------------------|---------------------------|
| 5/26/52 | 3'           | 0.60          | 0.04        | 27              | 8            | 9.2                    | 4,085                     |      |              |                      |             |                 |              |                        |                           |

Preceding page blank

APPENDIX "D"

SUMMARY OF INDIVIDUAL ANALYTICAL  
RESULTS FOR THE SAVANNAH RIVER SECTION

S-23 to S-0

**SAVANNAH RIVER STUDIES**  
**Summary of Individual Analytical Results for the Savannah River Section, S-23 to S-2**  
**March 28, 1952**

| Station | High Tide          |     |              |             |      |     | Between Mean and High Tide |     |              |             | Mean Tide          |     |           |     | Low Tide          |             |      |     |           |     |        |            |     |
|---------|--------------------|-----|--------------|-------------|------|-----|----------------------------|-----|--------------|-------------|--------------------|-----|-----------|-----|-------------------|-------------|------|-----|-----------|-----|--------|------------|-----|
|         | 9:45 AM - 11:15 AM |     |              |             |      |     | 11:25 AM - 12:55 PM        |     |              |             | 12:25 PM - 2:30 PM |     |           |     | 4:00 PM - 5:49 PM |             |      |     |           |     |        |            |     |
|         | Chlorides          | ppm | Chlorides as | % Sea Water | D.O. | ppm | Chlorides                  | ppm | Chlorides as | % Sea Water | D.O.               | ppm | Chlorides | ppm | Chlorides as      | % Sea Water | D.O. | ppm | Chlorides | ppm | (M.O.) | Alkalinity | ppm |
|         |                    |     |              |             |      |     |                            |     |              |             |                    |     |           |     |                   |             |      |     |           |     |        |            |     |

**Surface Sample (3 feet below surface)**

| Station | Chlorides ppm | Chlorides as % Sea Water | D.O. ppm | D.O. % Saturation | Alkalinity (M.O.) ppm | Chlorides ppm | Chlorides as % Sea Water | D.O. ppm | D.O. % Saturation | Chlorides ppm | Chlorides as % Sea Water | D.O. ppm | D.O. % Saturation | Chlorides ppm | Chlorides as % Sea Water | D.O. ppm | D.O. % Saturation | Alkalinity (M.O.) ppm |
|---------|---------------|--------------------------|----------|-------------------|-----------------------|---------------|--------------------------|----------|-------------------|---------------|--------------------------|----------|-------------------|---------------|--------------------------|----------|-------------------|-----------------------|
| S-2     | 2,170         | 11                       | 5.0      | 50                | 29                    | 1,710         | 9                        | 5.6      | 56                | 1,770         | 9                        | 6.0      | 60                | 560           | 3                        | 6.3      | 62                | 28                    |
| S-5     | 1,150         | 6                        | 5.2      | 52                | 29                    | 930           | 5                        | 5.3      | 53                | 870           | 5                        | 6.1      | 60                | 155           | 0.8                      | 6.3      | 62                | —                     |
| S-8     | 760           | 4                        | 5.3      | 53                | 24                    | 340           | 2                        | 5.8      | 58                | 41            | 0.2                      | 6.6      | 65                | 21            | 0.1                      | 6.8      | 67                | 19                    |
| S-11    | 55            | 0.3                      | 7.0      | 69                | 20                    | 12            | <0.1                     | 7.2      | 71                | 9.7           | <0.1                     | 7.2      | 71                | 6.7           | <0.1                     | 7.3      | 72                | 16                    |
| S-14    | 4.4           | <0.1                     | 7.3      | 72                | 14                    | 4.1           | <0.1                     | 7.3      | 72                | 3.6           | <0.1                     | 7.3      | 72                | 4.7           | <0.1                     | 7.4      | 73                | 14                    |
| S-17    | 3.4           | <0.1                     | 7.5      | 74                | 14                    | 2.8           | <0.1                     | 7.5      | 74                | 3.3           | <0.1                     | 7.6      | 75                | 3.3           | <0.1                     | 7.6      | 75                | 13                    |
| S-20    | 2.4           | <0.1                     | 7.4      | 73                | 17                    | 2.6           | <0.1                     | 7.5      | 74                | 3.3           | <0.1                     | 7.6      | 75                | 3.8           | <0.1                     | 8.3      | 82                | 13                    |
| S-23    | 3.3           | <0.1                     | 7.6      | 74                | 13                    | 3.0           | <0.1                     | 7.7      | 76                | 3.3           | <0.1                     | 7.8      | 77                | 3.2           | <0.1                     | 7.7      | 75                | 13                    |

**Bottom Samples (3 feet above channel bottom, or top of sludge bank)**

| Station | Chlorides ppm | Chlorides as % Sea Water | D.O. ppm | D.O. % Saturation | Alkalinity (M.O.) ppm | Chlorides ppm | Chlorides as % Sea Water | D.O. ppm | D.O. % Saturation | Chlorides ppm | Chlorides as % Sea Water | D.O. ppm | D.O. % Saturation | Chlorides ppm | Chlorides as % Sea Water | D.O. ppm | D.O. % Saturation | Alkalinity (M.O.) ppm |
|---------|---------------|--------------------------|----------|-------------------|-----------------------|---------------|--------------------------|----------|-------------------|---------------|--------------------------|----------|-------------------|---------------|--------------------------|----------|-------------------|-----------------------|
| S-2     | 15,250        | 80                       | 6.9      | 80                | 93                    | 14,660        | 77                       | 6.9      | 79                | 10,430        | 55                       | 5.7      | 63                | 2,830         | 15                       | 5.2      | 53                | —                     |
| S-5     | 2,670         | 14                       | 4.7      | 48                | 25                    | 13,550        | 71                       | 5.7      | 65                | 4,610         | 24                       | 5.0      | 52                | 870           | 5                        | 5.9      | 59                | 96                    |
| S-8     | 900           | 5                        | 5.2      | 51                | 22                    | 6,200         | 33                       | 4.7      | 49                | 240           | 1                        | 6.1      | 60                | 86            | 0.5                      | 6.0      | 59                | 22                    |
| S-11    | 150           | 0.7                      | 6.4      | 63                | 20                    | 20            | 0.1                      | 7.4      | 73                | 13            | <0.1                     | 3.0      | 30                | 15            | <0.1                     | 6.5      | 64                | 17                    |
| S-14    | 4.7           | <0.1                     | 7.3      | 72                | 14                    | 4.2           | <0.1                     | 7.2      | 71                | 3.6           | <0.1                     | 7.4      | 73                | 4.2           | <0.1                     | 7.4      | 73                | 14                    |
| S-17    | 3.5           | <0.1                     | 7.6      | 75                | 19                    | 3.2           | <0.1                     | 7.4      | 73                | 3.3           | <0.1                     | 7.6      | 75                | 3.5           | <0.1                     | 7.6      | 75                | 12                    |
| S-20    | 5.5           | <0.1                     | 7.6      | 75                | 24                    | 3.0           | <0.1                     | 7.6      | 75                | 3.3           | <0.1                     | 7.6      | 75                | 3.9           | <0.1                     | 7.7      | 76                | 14                    |
| S-23    | 3.3           | <0.1                     | 7.7      | 76                | 16                    | 3.3           | <0.1                     | 7.6      | 75                | 3.3           | <0.1                     | 7.6      | 75                | 3.4           | <0.1                     | 8.3      | 82                | 14                    |

SAVANNAH RIVER STUDIES  
Summary of Individual Analytical Results for the Savannah River Section, S-23 to S-0  
June 23, 1952

| Time of Sampling                                                    | High Tide (Falling) |                  |                             |           |                 | Low Tide (Rising)  |                  |                             |           |                 |
|---------------------------------------------------------------------|---------------------|------------------|-----------------------------|-----------|-----------------|--------------------|------------------|-----------------------------|-----------|-----------------|
|                                                                     | 9:07 AM to 10:19 AM |                  |                             |           |                 | 3:25 PM to 5:04 PM |                  |                             |           |                 |
|                                                                     | Temperature<br>°C   | Chlorides<br>ppm | Chlorides as<br>% Sea Water | D.O., ppm | D.O. Saturation | Temperature<br>°C  | Chlorides<br>ppm | Chlorides as<br>% Sea Water | D.O., ppm | D.O. Saturation |
| Surface Samples (3 feet below surface)                              |                     |                  |                             |           |                 |                    |                  |                             |           |                 |
| S-0                                                                 | 30                  | 15,440           | 81                          | 3.8       | 58              | 30                 | 9,100            | 48                          | 5.2       | 76              |
| S-2                                                                 | 30                  | 12,340           | 65                          | 5.4       | 81              | 30                 | 6,280            | 33                          | 4.2       | 58              |
| S-5                                                                 | 30                  | 9,480            | 50                          | 5.0       | 72              | 30                 | 4,480            | 24                          | 3.5       | 48              |
| S-8                                                                 | 30                  | 6,480            | 34                          | 3.5       | 49              | 30                 | 2,350            | 12                          | 2.8       | 38              |
| S-11                                                                | 30                  | 4,220            | 22                          | 2.6       | 36              | 30                 | 1,560            | 8.2                         | 4.3       | 58              |
| S-14                                                                | 29                  | 1,570            | 8.3                         | 2.2       | 29              | 30                 | .80              | 0.4                         | 3.5       | 46              |
| S-17                                                                | 29                  | 390              | 2.1                         | 1.4       | 19              | 30                 | 7                | 0.04                        | 5.7       | 74              |
| S-20                                                                | 30                  | 11               | 0.1                         | 4.6       | 60              | 30                 | 5                | 0.03                        | 6.6       | 86              |
| S-23                                                                | 30                  | 5                | 0.03                        | 6.4       | 83              | 30                 | --               | --                          | 6.7       | 88              |
| Bottom Samples (3 feet above channel bottom, or top of sludge bank) |                     |                  |                             |           |                 |                    |                  |                             |           |                 |
| S-0                                                                 | 30                  | 17,930           | 94                          | 5.7       | 90              | 29                 | 16,700           | 88                          | 5.2       | 80              |
| S-2                                                                 | 30                  | 13,100           | 69                          | 5.7       | 86              | 29                 | 15,920           | 84                          | 4.9       | 75              |
| S-5                                                                 | 30                  | 16,560           | 87                          | 5.0       | 79              | 29                 | 14,200           | 75                          | 4.0       | 60              |
| S-8                                                                 | 30                  | 12,500           | 66                          | 4.2       | 63              | 29.5               | 8,570            | 45                          | 2.9       | 42              |
| S-11                                                                | 30                  | 11,460           | 60                          | 3.1       | 46              | 29                 | 2,670            | 14                          | 1.9       | 26              |
| S-14                                                                | 29                  | 3,710            | 20                          | 1.8       | 24              | 29                 | 910              | 5                           | 1.5       | 19              |
| S-17                                                                | 29                  | 1,650            | 8.7                         | 1.0       | 13              | 29.5               | 21               | 0.1                         | 5.6       | 73              |
| S-20                                                                | 29                  | 12               | 0.1                         | 4.0       | 52              | 29.5               | 7                | 0.04                        | 6.6       | 85              |
| S-23                                                                | 29                  | 5                | 0.03                        | 6.2       | 80              | 29.5               | --               | --                          | 6.7       | 87              |

Preceding page blank -15

APPENDIX "D"

DESCRIPTION OF THE PROCEDURE USED

FOR

ASSAYING THE RADIOACTIVITY IN FRESH WATER

AND FOR BIOTA

Obviously, when dealing with low level radioactive samples in the order of the minimum detectable amount, the observed count, because of statistical fluctuations, may be lower than the observed background. This gives rise to negative radioactive values. This is to be expected and indicates that the precision of the determination is poor.

#### FRESH WATER PROCEDURE

The procedure used for assaying the radioactivity in fresh water was as follow

##### 1. Suspended solids

- a. Filter 500 ml. of the well mixed water sample through a membrane filter \*/. Save the filtrate for Part 2. Allow the membrane plus the suspended solids to dry for several hours at room temperature before counting.
- b. Place the membrane filter in an aluminum dish. Insert an iron ring on the periphery of the filter to hold it flat. Insert sample into an internal proportional gasflow counter set at a voltage for alpha counting. Flush with P-10 gas (90 percent Argone-10 percent methane) and count for 16 minutes. Reset the instrument and count for an additional 16 minutes. If the 16 minute counts do not agree within the statistical error of counting, repeat the count. Repeat the counting procedure in an internal counter set at a voltage for beta counting. Correct for backscatter and geometry and convert the values to uuc/liter of water.

##### 2. Dissolved solids

- a. Place aluminum dishes (48 mm. in diameter and 15 ml. in capacity) in an oven at 103° C. for two hours. Cool in a desiccator and weigh.

\*/ Clark, H.F., Geldreich, E.D., Jeter, H.L., and Kabler, P.W. "Membrane Filter in Sanitary Bacteriology". Public Health Reports 66, 951 (1951).



- b. Evaporate duplicate 200 ml. portions of the filtered sample from Part 1 in duplicate weighed aluminum dishes on a hot plate which has been adjusted to keep the temperature slightly under the boiling point of the water.
- c. Dry the samples at 103-105°C. to constant weight.
- d. Place dish in internal counters and count for two 16 minute periods at a voltage for alpha counting and again at the beta counting voltage as for suspended solids. Convert the alpha and beta counts to  $\mu\text{mc/liter}$  of water.

#### BIOTA PROCEDURE

The laboratory processing procedure is as follows: \*/

##### 1. Fish Flesh

A composite sample of 10 fishes is dissolved in 50 ml. of 1:1 nitric acid. In the case of composite samples an aliquot equal to 1 to 2 grams of weighed fish, or in the case of an individual specimen 1 or 2 grams of weighed fish flesh is transferred to a 50 ml. Erlenmeyer flask with about 5 ml. of distilled water and 5 ml. of concentrated nitric acid. The sample is gradually heated to boiling on a hot plate and evaporated to dryness. About 10 ml. of

---

\*/ It should be noted that the procedures adopted for low level assay of biological samples may cause the loss of some radioactivity such as carbon, iodine and gases. The loss of small amounts of this radioactivity is considered to be relatively unimportant from a public health standpoint. The results include natural potassium-40 which might conceivably increase the fish flesh beta radioactivity by as much as 3  $\mu\text{mc/g.}$



concentrated nitric acid is added and the solution warmed to about 50°C. Hydrogen peroxide (30 per cent solution) is added dropwise, as required, to decolorize and promote wet oxidation. Digestion is continued at a gentle boiling temperature to dryness. The latter steps were repeated if necessary, until the residual solids in the flask are white and crystalline in nature. The acid digested mineral residue represents one to two per cent of the weight of the flesh. The residue is dissolved in a known volume of 0.5 milliequivalents of nitric acid and 7 to 10 ml. of water. It is convenient to prepare this clear digestate volume so that 2 ml. contains 0.2 or 0.4 gram of original tissue. A sample for counting is then prepared by compositing 2 ml. from each of two to five digestates in an aluminum dish and evaporating gently to dryness. The white crystalline solids are distributed fairly uniformly over the dish bottom. The sample is then counted for alpha and beta radioactivity as explained previously for water samples and reported in uuc/gram of original tissue. If the composite sample shows the presence of alpha or beta activity above a statistically normalized average, a sample of each of the digestates is deposited on an aluminum dish and counted separately.

## 2. Fish Bone

Bony tissue is removed from fish specimen, air dried, scraped free of flesh, and stored in vials at the field laboratory and

shipped to Cincinnati. Preliminary determinations of these fish bone samples indicate a wide fluctuation in the moisture content, probably due to variable preparation, shipping and storage conditions. Consequently, composite samples of 5 to 10 grams of bone are ignited in a tared crucible at 450° C. for at least four hours, cooled and weighed. The ignited bone sample is then ground to a powder in a mortar. One gram of bone powder is transferred to an Erlenmeyer flask and dissolved in about 2 ml. of warm one molar nitric acid. After solution, the volume is diluted with water so that each ml. contains 5.0 mg. of bone ash. Five to 10 ml. of solution is transferred to an aluminum dish, evaporated gently to dryness and counted. Small samples of about one gram of bone are similarly ignited in a tared crucible, weighed and dissolved without grinding.

### 3. Brush Box Organisms and Algae

These samples are treated similarly to fish flesh samples. In general, the samples required longer periods of acid digestion and the residues are frequently colored and difficult to dissolve in dilute nitric acid after wet oxidation. Suspensions of the residue are transferred to aluminum dishes and counted.

APPENDIX "F"

CHEMICAL METHODS OF WATER ANALYSIS

# MANGANESE - COLORIMETRIC DETERMINATION

Manganese is readily determined in small concentrations by the colorimetric determination of permanganate formed by oxidation of the manganese with periodate in acidic solution.

## 1. Apparatus

1.1 Evaporation dishes 500 - 1000 ml. capacity

1.2 Nessler tubes for visual color comparison

## 2. Reagents

2.1 Sulfuric acid, Concd. reagent grade

2.2 Perchloric acid 70-72% reagent grade

2.3 Sodium paraperiodate

2.4 Phosphoric acid 6M

2.5 Manganous Sulfate standard solution. Dissolve 0.1438 of  $KMnO_4$  in distilled water to which has been added 2-3 ml. of 2N  $H_2SO_4$ . Reduce by adding 0.4 gm. of  $NaHSO_3$ . Boil off excess  $SO_2$  and dilute to one liter with distilled water.

2.6 Sodium paraperiodate - sulfuric acid reagent. Prepare a  $H_2SO_4$  solu. by adding 120 ml. of conc.  $H_2SO_4$  to 1500 ml. of distilled water and diluting to 2 liters. Add 2.4 gm. of  $Na_3H_2IO_6$  heat to boiling and place for 30 minutes, in a boiling water bath.

2.7 Manganese color standard. Add 1.2 ml. of conc.  $H_2SO_4$  and 30 ml. of sodium paraperiodate-sulfuric acid reagent to exactly 20.0 ml. of standard  $MnSO_4$  solu. After thoroughly mixing, add 0.3 gm. of solid  $Na_3H_2IO_6$ , heat to boiling

and place for 30 minutes in a boiling water bath. Cool and dilute to one liter with  $H_2SO_4$  reagent. This standard contains 0.001 mg. of permanganate manganese per ml. Prepare a series of color standards in accordance with Table 10, page 57, 9th Edition, Standard Methods.

2.8 Nitric Acid, Conc. reagent grade

2.9 Aluminum hydroxide suspension

### 3. Preparation of Sample

Turbid and colored samples should be clarified with aluminum hydroxide in order to obtain values for manganese in solution only. Method of preparation of aluminum hydroxide appears on page 74, 9th Edition of Standard Methods for the Examination of Water and Sewage.

Aluminum hydroxide, at the rate of about 10 ml. per liter of sample, is added, well mixed, and then allowed to settle.

The suspension is filtered. If a clear and colorless solution is not obtained, the process is repeated.

### 4. Procedure

To a portion of sample containing 0 - 0.1 mg. of manganese, add 5 ml. of  $HNO_3$  and evaporate on a steam bath or on a hot plate at just below the boiling point to a volume of 15-20 ml. Transfer the solution, together with any solids remaining in the dish, to a 125-ml. conical flask, add 5 ml. additional  $HNO_3$  and 10 ml. of  $HClO_4$  70-72% and a few glass beads to prevent bumping. Evaporate on a hot plate until dense fumes



appear in the flask or until digestion is considered complete. The solution should be colorless. Repeat the addition of  $\text{HNO}_3$  and  $\text{HClO}_4$  if necessary. Cool the solution to room temperature, dilute to about 50 ml. with distilled water, boil to expel chlorine, then allow to cool. Filter through porcelain filter or paper into a 100-ml. volumetric flask, then make up to volume. Mix.

An aliquot of the acid-digested sample is placed into a 250-ml. conical flask and made up to a volume of about 50 ml. with distilled water. Add 20 ml.  $\text{H}_3\text{PO}_4$  5N, 1 ml.  $\text{H}_2\text{SO}_4$  conc. and 0.3 gm.  $\text{Na}_2\text{H}_2\text{IO}_6$ . Then heat the solution to boiling and keep just below boiling for 30 minutes. Cool to room temperature, then transfer the solution to a 100 ml. Nessler tube, make up to mark and compare with standards.

#### 5. Calculations

$$\text{ppm Mn} = \frac{\text{mg. Mn in the matched standard} \times 1000}{\text{ml. sample}}$$

### IRON - COLORIMETRIC DETERMINATION PHENANTHROLINE METHOD

Ortho-phenanthroline reacts with ferrous iron in dilute solution at a pH of 2.0 to 9.0 to give a pink to red color.

#### 1. Apparatus

1.1 Nessler tubes for visual color determination.

#### 2. Reagents

2.1 Hydroxylamine hydrochloride. Dissolve 10 gm.  $\text{NH}_2\text{OH}$  in

90 ml. water. Prepare fresh every few days.

2.2 Iron standard solution. Dissolve 0.7022 gm. of  $\text{FeSO}_4(\text{NH}_4)_2\text{SO}_4 \cdot 6\text{H}_2\text{O}$  in 50 ml. distilled water and 20 ml. concd.  $\text{H}_2\text{SO}_4$  and dilute to 1 liter. One ml. of the solution contains 0.1 mg. Fe.

2.3 Iron standard solution 0.001 mg. Fe per ml. Prepare shortly before use by diluting 10 ml. of 2.2 to 1000 ml.

2.4 1,10- phenanthroline. Dissolve 0.5 gr. of the monohydrate in 500 ml. of water heated to 80-90°C.

2.5 Sodium acetate. Dissolve 200 gr. of sodium acetate in 300 ml. of water.

### 3. Preparation of Sample

Turbid and colored samples should be clarified with aluminum hydroxide in order to obtain values for iron in solution only.

### 4. Procedure

Place an aliquot of the sample into a 100 ml. Nessler tube. Add 1 ml. of  $\text{NH}_2\text{OH}$  and 10 ml. sodium acetate solution. Dilute to about 75 ml. and add 10 ml. 1,10- phenanthroline solution. Dilute to volume, mix, and compare after 10 minutes with standards. The standards are stable for six months.

### 5. Calculations

$$\text{ppm Fe} = \frac{\text{mg Fe in the matched Std.} \times 1000}{\text{ml. sample}}$$

# 6. Suggested Standards

|       |          |
|-------|----------|
| 1 ml. | .001 mg. |
| 2     | .002     |
| 4     | .004     |
| 6     | .006     |
| 8     | .008     |
| 10    | .01      |
| 15    | .015     |
| 20    | .02      |

## Notes and Comments

1. Digestion fumes can be controlled by inserting a small funnel into neck of Erlenmeyer flask.
2. Substitute nitric acid for perchloric acid in digestion and proceed as follows  
 Acidify sample with 5 ml. nitric acid, evaporate to 15-20 ml., transfer, together with any solids remaining in the dish, to a 125 ml. Erlenmeyer Flask, add 5 ml. additional nitric acid and 5 ml. sulfuric acid. Evaporate on a hot plate until dense white fumes of  $\text{SO}_3$  appear.
3. Convenient sample size for digestion is 500 ml. Use the major portion for the determination of manganese and the remaining for iron.
4. Standards for both manganese and iron are stable for at least six months.

APPENDIX "C"

BIOLOGICAL METHODS AND TABLE

(COMPLETE LIST OF ORNITHINE  
FOUND IN BRUSH BOX SAMPLES)

# PLANKTON COLLECTION AND COUNTING METHODS

Plankton samples were collected with a two-liter Kemmerer bottle, and were concentrated by pouring them through a standard Wisconsin plankton net. The concentrates were washed from the bucket of the plankton net with distilled water and preserved in the field with formalin. Nannoplankton was sampled by saving 500 milliliters of the filtrate from the plankton net and preserving it in the field with formalin. The nannoplankton samples were later concentrated in the laboratory by use of a Foerst plankton centrifuge. The net plankton samples were counted by examining the entire contents of a one-milliliter Sedgwick-Rafter counting cell at a magnification of 100 diameters. The nannoplankton samples were counted by the use of the Sedgwick-Rafter counting cell and Whipple ocular micrometer at a magnification of 210 diameters. The number of Sedgwick-Rafter cells or Whipple fields counted for each sample depended upon the abundance of organisms in the sample. The net and nannoplankton counts were expressed as organisms per liter.

## BOTTOM ORGANISM COLLECTION METHODS

### Petersen Dredge

Samples were obtained with a Petersen Dredge lowered from a swivelled winch and boom mounted on a fourteen-foot boat. The Petersen Dredge samples an area of about .08 square meter, but the volume of materials taken varies greatly depending upon the character of the bottom and how completely the dredge fills. It was found most practicable to use the dredge without additional weights, even in swift water. Each sample consisted of 2 to 4 effective grabs with the dredge, taken

from as near the same spot as was possible while drifting the boat with the current. Incomplete samples, obtained when the jaws of the dredge were held apart by a stick or other object, were discarded. The bottom materials from each location were deposited in a wash tub and were washed through a series of nesting screens, the finest of which had a mesh size of 40 openings per inch. Each screen was washed thoroughly with a bucket pump and hose and the organisms retained by it were removed and preserved. No attempt was made to recover organisms which passed through the finest screen.

#### Brush Boxes

These consisted of 16" cubes of 1/2" mesh hardware cloth supported on the inside by a wooden frame. This size was chosen as one which would give an adequate sample, and would still fit into the top of a square wash tub. The boxes were floored inside with a piece of bronze window screening to prevent the escape of organisms as the boxes were being raised. A quantity of sticks, roots, and other natural debris, together with enough stones to make it sink, was placed inside the box. The top was wired down in such a manner that it could be easily opened again to remove the contents. Brush boxes were placed in situations from which samples were desired and allowed to remain for varying lengths of time. Upon removal from the water, the boxes were placed in tubs and washed thoroughly with a jet of water. The wash water was then strained through the same series of screens used for handling bottom samples and the organisms were treated in essentially the same manner as those from bottom samples. If convenient, the washing and screening was



done with tap water at the Augusta laboratory, but boxes from stations remote from the laboratory were washed and screened in the field with the aid of a bucket pump. In either case, when the fine mesh screen retained a very large amount of finely divided material and small organisms, these were preserved with formalin and carefully sorted in white enameled trays in the laboratory.

TABLE I -- Appendix "G"

COMPLETE LIST OF ORGANISMS

FOUND IN BRUSH BOX SAMPLES

TABLE I (Appendix "G") SAVANNAH RIVER STUDIES

July 1951-July 1952

COMPLETE LIST - Brush Box Sampling - Numbers of Organisms

| Location (River Mi.) | 8-223.15 | 8-219.0 |        |         |         |          |
|----------------------|----------|---------|--------|---------|---------|----------|
| Date Removed         | 8/24/51  | 7/30/51 | 8/6/51 | 8/13/51 | 8/20/51 | 10/16/51 |
| No. of days in place | 21       | 7       | 14     | 21      | 28      | 21       |

## PORIFERA

Spongilla fragilis

## TURBELLARIA

Dugesia tigrina

Undetermined

1

## ROTATORIA

Floscularia sp.

Pr.\*

Pr.

Pr.

## BRYOZOA

Plumatella repens

Pr.

## ANNELIDA: OLIGOCHAETA

Aulophorus furcatusPelosclex sp.

Undet. Naididae

Undet. Tubificidae

Undetermined

8

2

8

## ANNELIDA: HIRUDINEA

Undet. Glossiphoniidae

Undet. Erpobdellidae

Helobdella sp.

Undetermined

3

## ARTHROPODA: CRUSTACEA

Copepoda: Cyclops sp.Ostracoda: Cypris sp.,Cypridopsis sp.

145

6

10

43

Isopoda: Asellus militarisMancasellus sp.Amphipoda: Hyalella aztecaDecapoda: Macrobrachium ohionePalaemonetes paludosus

1

## ARTHROPODA: ARACHNIDA

Hydracarina: Albia sp. Arrenurus sp.Libertia sp. Neumania

Undetermined

13

\* = Present

(continued)

BLE I (Appendix "G")  
(continued)

## SAVANNAH RIVER STUDIES

July 1951-July 1952

COMPLETE LIST - Brush Box Sampling

- Numbers of Organisms

| S-215.5 |         |          | S-213.3 | S-203.3 | S-201.55 | S-172.0 |        |
|---------|---------|----------|---------|---------|----------|---------|--------|
| 8/27/51 | 9/19/51 | 10/16/51 | 9/19/51 | 8/27/51 | 8/28/51  | 7/27/52 | 9/4/51 |
| 21      | 21      | 21       | 21      | 21      | 22       | 165     | 21     |

Pr.

9

1

238

12

174

Pr.\*

Pr.

Pr.

Pr.

Pr.

Pr.

Pr.

Pr.

Pr.

50

3

6

12

3

8

396

22

13

1

3

9

1

3

16

79

1

20

3

3

5

4

3

\* = Present

(continued)

TABLE I (Appendix "G")  
(continued)

## SAVANNAH RIVER STUDIES

July 1951-July 1952

COMPLETE LIST - Brush Box Sampling

- Numbers of Organisms

| Location (River Mi.) | 8-171.7 | 8-171.6  | 8-171.52 | 8-171.19 | 8-165. |
|----------------------|---------|----------|----------|----------|--------|
| Date Removed         | 10/3/51 | 12/20/51 | 10/4/51  | 10/31/51 | 9/4/51 |
| No. of days in place | 21      | 37       | 22       | 26       | 21     |

## PORIFERA

Spongilla fragilis

Pr

## TURBELLARIA

Dugesia tigrina

1

1

4

Undetermined

## ROTATORIA

Floscularia sp.

Pr.

Pr.

## BRYOZOA

Plumatella repens

Pr.

Pr.

## ANNELIDA: OLIGOCHAETA

Aulophorus furcatusPelosclex sp.

Undet. Naididae

Undet. Tubificidae

1

1

Undetermined

60

## ANNELIDA: HIRUDINEA

Undet. Glossiphoniidae

1

Undet. Erpobdellidae

Helobdella sp.

Undetermined

## ARTHROPODA: CRUSTACEA

Copepoda: Cyclops sp.Ostracoda: Cypris sp.,

1

cypridopsis sp.Isopoda: Asellus militaris

3

Mancasellus sp.

1

Amphipoda: Hyaella aztecaDecapoda: Macrobrachium ohione

1

Palaemonetes paludosus

## ARTHROPODA: ARACHNIDA

Hydracarina: Albia sp. Arrenurus sp.Libertia sp. Neumania

Undetermined

1

5

1

\* = Present

(continued)

TABLE I (Appendix "G")  
(continued)

## SAVANNAH RIVER STUDIES

July 1951-July 1952

COMPLETE LIST - Brush Box Sampling

- Numbers of Organisms

|          |          |                    |          |          |          |         |         |
|----------|----------|--------------------|----------|----------|----------|---------|---------|
| 8-160.75 | 8-155.25 | 8-155.2<br>8-155.0 | 8-155.17 | 8-146.4  | 8-144.7  | 8-136.6 | 8-51.0  |
| 11/2/51  | 10/1/51  | 9/26/51            | 9/6/51   | 10/11/51 | 10/11/51 | 2/6/52  | 7/31/52 |
| 32       | 24       | 20                 | 22       | 21       | 21       | 53      | 160     |

1

7  
1

3

Pr.\*

Pr.

Pr.

3  
1  
2

1

4

2

5

1  
1

1

2

1

13

6

1

2

1

2

2

6

33

8

\* = Present

(continued)



TABLE I (Appendix "G")  
(Continued)

## SAVANNAH RIVER STUDIES

July 1951-July 1952

## COMPLETE LIST - Brush Box Sampling

## Numbers of Organisms

| Location (River Mi.) | S-223.15 |         |        |         |         |          | S-219.0 |
|----------------------|----------|---------|--------|---------|---------|----------|---------|
| Date Removed         | 8/24/51  | 7/30/51 | 8/6/51 | 8/13/51 | 8/20/51 | 10/16/51 |         |
| No. of days in place | 21       | 7       | 14     | 21      | 28      | 21       |         |

## ARTHROPODA: INSECTA

Plecoptera: Pteronarcys pictetiiTaeniopteryx litaNemoura sp.Allocaenia sp.Neoperla clymeneAcroneuria abnormisAcroneuria arenosaAcroneuria georgianaAcroneuria melaUndet. AcroneuriaPhasganophora capitataParagnetina kansensisUndet. PerlidaeIsoperla sp.Undet. Chloroperlidae

Undetermined

Ephemeroptera: Hexagenia sp.

3

Tortopus sp.Neopphemera youngiS. pulchellum gr. 1

3

17

23

41

50

45

Stenonema exiguum,S.interpunctatum,Undet. S.Heptagenia julia(?),H. pulla(?)Undet. H.Undet. HeptageniidaeUndet. AmetropinaeIsonychia spp.Paraleptophlebia sp.Ephemerella trilineata (?)Ephemerella invaria gr.nr.dorotheaUndet. EphemerellaCaenis spp.

21

1

2

10

Tricorythodes

1

29

25

10

29

1

albidineatus, 1 sp.Baetis spinosus

1

Baetis sp.

2

Undet. Baetinae

(continued)

TABLE I (Appendix "G")  
(continued)

## SAVANNAH RIVER STUDIES

July 1951-July 1952

COMPLETE LIST - Brush Box Sampling

-

Numbers of Organisms

| S-215.5 |         |          | S-213.3 | S-203.3 | S-201.55 | S-172.0 |        |
|---------|---------|----------|---------|---------|----------|---------|--------|
| 8/27/51 | 9/19/51 | 10/16/51 | 9/19/51 | 8/27/51 | 8/28/51  | 7/27/52 | 9/4/51 |
| 21      | 21      | 21       | 21      | 21      | 22       | 165     | 21     |

1

1

1

12

2

2

34

2

10

1  
5

1

68

1

1

1

1

1

(continued)

TABLE I (Appendix "G")  
(Continued)

## SAVANNAH RIVER STUDIES

July 1951-July 1952

## COMPLETE LIST - Brush Box Sampling

## Numbers of Organisms

| Location (River Mi.) | S-171.7 | S-171.6  | S-171.52 | S-171.15 | S-165.4               |
|----------------------|---------|----------|----------|----------|-----------------------|
| Date Removed         | 10/3/51 | 12/20/51 | 10/4/51  | 10/31/51 | 9/4/51 9/4/51 7/17/51 |
| No. of days in place | 21      | 37       | 22       | 26       | 21 21 139             |

## ARTHOPODA: INSECTA

Plecoptera: Pteronarcys pictetii

3

Taeniopteryx lita

5

1

Nemoura sp.Allocaonia sp.

2

Neoperla clymene

2

3

Acroneuria abnormis

14

1

8

9

7

3

Acroneuria arenosaAcroneuria georgianaAcroneuria mela

1

6

1

3

1

Undet. Acroneuria

1

4

Phasganophora capitata

2

Paragnetina kansensis

67

4

10

31

4

13

Undet. Perlidae

3

23

9

7

Isoperla sp.Undet. Chloroperlidae

## Undetermined

Ephemeroptera: Hexagenia sp.Tortopus sp.

2

Neoephemera youngi

34

S. pulchellum gr. 1

131

36

241

136

129

5

1

Stenonema exiguum,S. interpunctatum,Undet. S.Heptagenia julia(?).H. pulla(?)Undet. H.

6

1

25

Undet. Heptageniidae

95

21

Undet. AmetropinaeIsonychia spp.

1

5

2

Paraleptophlebia sp.Ephemerella trilineata(?)

2

4

Ephemerella invaria gr.nr. dorotheaUndet. Ephemerella

3

Caenis spp.

273

2

22

27

328

52

Tricorythodesalbidineatus, 1 sp. 72

22

6

62

10

Baetis spinosus

2

Baetis sp.Undet. Baetinae

1

(continued)

TABLE I (Appendix "G")  
(continued)

## SAVANNAH RIVER STUDIES

July 1951-July 1952

COMPLETE LIST - Brush Box Sampling

- Numbers of Organisms

| S-160.75 | S-155.25 | S-155.2<br>St-0.0 | S-155.17 | S-146.4  | S-144.7  | S-136.6 | S-51.0  |
|----------|----------|-------------------|----------|----------|----------|---------|---------|
| 11/2/51  | 10/1/51  | 9/26/51           | 9/6/51   | 10/11/51 | 10/11/51 | 2/6/52  | 7/31/52 |
| 32       | 24       | 20                | 22       | 21       | 21       | 53      | 160     |

|     |     |   |    |   |     |    |    |
|-----|-----|---|----|---|-----|----|----|
|     |     |   | 1  |   | 2   |    |    |
|     |     |   |    |   |     | 1  |    |
|     |     |   |    |   |     | 6  |    |
| 6   |     |   | 4  |   | 10  | 2  |    |
| 3   | 1   |   | 2  |   | 1   |    | 7  |
|     |     |   |    |   |     | 1  |    |
| 36  | 4   |   | 6  |   | 61  | 6  |    |
|     | 2   |   | 1  | 1 | 1   | 11 |    |
|     |     |   |    |   |     | 1  |    |
|     |     |   | 1  |   |     | 5  |    |
| 1   |     |   | 1  |   | 1   |    |    |
| 189 | 7   |   | 17 | 3 | 169 | 79 | 13 |
|     |     |   |    |   |     |    |    |
| 20  |     |   | 2  |   | 43  | 14 |    |
| 3   |     | 1 | 1  |   |     | 15 |    |
|     |     |   |    |   |     |    |    |
|     |     |   | 1  |   | 12  |    | 48 |
| 1   |     |   |    |   |     |    |    |
| 10  |     |   |    |   | 5   | 4  |    |
|     |     |   |    |   |     |    |    |
|     |     |   |    |   |     | 13 |    |
|     |     |   |    |   |     | 25 |    |
| 63  | 146 |   | 32 | 2 | 240 |    | 82 |
| 26  | 15  |   | 27 | 2 | 43  |    | 11 |
| 4   |     |   |    |   | 1   |    |    |
|     |     |   |    |   |     |    | 1  |

(continued)

TABLE I (Appendix "G")  
(Continued)

## SAVANNAH RIVER STUDIES

July 1951-July 1952

## COMPLETE LIST - Brush Box Sampling

## Numbers of Organisms

| Location (River Mi.) | 8-223.15 | 7-30/51 | 8/6/51 | 8/13/51 | 8/20/51 | 10/16/51 |
|----------------------|----------|---------|--------|---------|---------|----------|
| Date Removed         | 8/24/51  | 7/30/51 | 8/6/51 | 8/13/51 | 8/20/51 | 10/16/51 |
| No. of days in place | 21       | 7       | 14     | 21      | 28      | 21       |

|              |                                          |    |    |    |    |    |
|--------------|------------------------------------------|----|----|----|----|----|
| Odonata:     | <u>Gomphus dilatatus</u>                 |    |    |    |    |    |
|              | <u>Gomphus plagiatus</u>                 |    |    |    |    |    |
|              | <u>Gomphus spiniceps</u>                 |    |    |    |    |    |
|              | <u>Undet. Gomphus</u>                    |    |    |    | 2  | 1  |
|              | <u>Dromogomphus spinosus</u>             |    |    |    |    |    |
|              | <u>Dromogomphus spoliatus</u>            |    |    |    |    |    |
|              | <u>Undet. Dromogomphus ed</u>            |    |    |    |    |    |
|              | <u>Ophiogomphus sp.</u>                  |    |    |    |    |    |
|              | <u>Boyeria vinosa</u>                    |    |    |    |    |    |
|              | <u>Macromia sp.</u>                      |    |    | 1  | 1  | 1  |
|              | <u>Neurocordulia alabamensis</u>         |    |    |    |    |    |
|              | <u>N. molesta</u> , <u>N. virginiens</u> | 1  | 1  | 3  | 4  | 8  |
|              | <u>Undet. Cordulinae</u>                 |    |    | 1  |    | 3  |
|              | <u>Perithemis sp.</u>                    |    |    |    |    | 1  |
|              | <u>Agrion sp.</u>                        |    |    | 1  |    |    |
|              | <u>Hetaerina titia</u>                   |    | 1  |    | 1  | 2  |
|              | <u>Argia moesta putrida</u> ,            |    |    |    |    |    |
|              | <u>translata</u> , <u>Undet. A. spp.</u> | 29 | 9  | 19 | 20 | 43 |
|              | <u>Enallagma antennatum</u> ,            |    |    |    |    | 55 |
|              | <u>E. signatum</u>                       | 4  |    |    |    |    |
|              | <u>Undet. Coenagrionidae</u>             |    | 19 |    |    | 1  |
| Neuroptera:  | <u>Sialis sp.</u>                        | 1  |    |    |    |    |
|              | <u>Corydalis cornutus</u>                |    |    |    |    |    |
| Trichoptera: | <u>Dolophilodes</u>                      |    |    |    |    |    |
|              | <u>Chimarra aterrima</u>                 |    |    |    |    |    |
|              | <u>Phylocentropus sp.</u>                |    |    |    |    |    |
|              | <u>Neureclipsis sp.</u>                  |    | 1  | 10 | 12 | 12 |
|              | <u>Polycentropus sp.</u>                 |    |    |    |    | 6  |
|              | <u>Psychomyia sp.</u>                    |    |    |    |    |    |
|              | <u>Cyrnellus marginatus</u>              |    |    |    |    |    |
|              | <u>Psychomyiid genus A(Ross)</u>         |    |    |    | 2  | 1  |
|              | <u>Psychomyiid genus B(Ross)</u>         | 5  |    |    |    |    |
|              | <u>Undet. Psychomyiidae</u>              |    |    |    |    |    |
|              | <u>Potamyia flava</u>                    |    |    |    |    |    |
|              | <u>Hydropsyche frison</u>                |    |    |    |    |    |
|              | <u>Hydropsyche orris</u>                 |    |    |    |    |    |
|              | <u>Hydropsyche simulans</u>              |    |    |    |    |    |
|              | <u>Undet. Hydropsyche</u>                |    |    |    |    |    |
|              | <u>Cheumatopsyche sp.</u>                |    |    |    |    |    |
|              | <u>Macronemus carolina</u>               |    | 6  | 1  | 10 | 7  |
|              | <u>Undet. Hydropsychidae</u>             |    |    |    |    | 1  |
|              | <u>Ochrotrichia sp.</u>                  |    |    |    |    |    |
|              | <u>Orthotrichia sp.</u>                  |    |    |    |    |    |
|              | <u>Hydroptila sp.</u>                    |    |    |    |    |    |
|              | <u>Undet. Limnophilidae</u>              |    |    |    |    |    |
|              | <u>Leptocella candida</u> , L.           |    |    |    |    |    |
|              | <u>genus A(Ross)</u>                     | 5  | 3  |    | 1  | 1  |

(continued)

TABLE I (Appendix "G")  
(continued)

## SAVANNAH RIVER STUDIES

July 1951-July 1952

## COMPLETE LIST - Brush Box Sampling

## Numbers of Organisms

| S-215.5 |         |          | S-213.3 | S-203.3 | S-201.55 | S-172.0 |        |
|---------|---------|----------|---------|---------|----------|---------|--------|
| 8/27/51 | 9/19/51 | 10/16/51 | 9/19/51 | 8/27/51 | 8/28/51  | 7/27/52 | 9/4/51 |
| 21      | 21      | 21       | 21      | 21      | 22       | 165     | 21     |

6

1

3

1

3

1

1

5

4

7

10

9

5

2

8

1

1

1

6

13

15

54

28

66

12

32

1

2

1

1

5

66

1

8

1

4

14

1

124

38

10

21

2

6

1

6

5

25  
2

1

22

2  
2

(continued)



TABLE I (Appendix "G")

## SAVANNAH RIVER STUDIES

July 1951-July 1952

(Continued)

## COMPLETE LIST - Brush Box Sampling

## Numbers of Organisms

| Location (River Mi.)                     | S-171.7  | S-171.6  | S-171.52   | S-171.15 | S-165.4 |
|------------------------------------------|----------|----------|------------|----------|---------|
| Date Removed                             | 10/3/51  | 12/20/51 | 10/4/51    | 10/31/51 | 9/4/51  |
| No. of days in place                     | 21       | 37       | 22         | 26       | 21      |
| Odonata: <u>Gomphus dilatatus</u>        |          |          |            |          |         |
| <u>Gomphus plagiatus</u>                 |          |          |            |          |         |
| <u>Gomphus spiniceps</u>                 |          |          |            | 1        | 2       |
| Undet. <u>Gomphus</u>                    |          |          |            |          |         |
| <u>Dromogomphus spinosus</u>             |          |          |            |          |         |
| <u>Dromogomphus spoliatus</u>            |          |          |            | 1        |         |
| Undet. <u>Dromogomphus</u> ed            |          |          |            |          |         |
| <u>Ophiogomphus</u> sp.                  |          |          |            |          |         |
| <u>Boyeria vinosa</u>                    |          |          | 1          | 1        | 3       |
| <u>Macromia</u> sp.                      |          |          |            |          |         |
| <u>Neurocordulia alabamensis</u>         |          |          |            |          |         |
| <u>N. molesta</u> , <u>N. virginiana</u> | 21       |          | 12         | 3        | 8       |
| Undet. <u>Cordulinae</u>                 |          |          | 1          | 32       | 3       |
| <u>Perithemis</u> sp.                    |          |          |            |          |         |
| <u>Agrion</u> sp.                        |          |          | 1          | 1        |         |
| <u>Hetaerina titia</u>                   | 1        |          | 1          | 1        | 4       |
| <u>Argia moesta putrida</u> ,            |          |          |            |          |         |
| <u>translata</u> , Undet. <u>A. spp.</u> | 13       |          | 67         | 12       | 87      |
| <u>Enallagma antennatum</u> ,            |          |          |            |          |         |
| <u>E. signatum</u>                       |          |          | 4          |          |         |
| Undet. <u>Coenagrionidae</u>             |          |          | 1          |          | 1       |
| Neuroptera: <u>Sialis</u> sp.            |          |          |            |          |         |
| <u>Corydalu cornutus</u>                 | 11       | 2        | 2          | 2        | 13      |
| Trichoptera: <u>Dolophilodes</u>         |          |          |            |          |         |
| <u>Chimarra aterita</u>                  |          | 7        |            | 31       | 1       |
| <u>Phyllocentropus</u> sp.               |          |          |            |          |         |
| <u>Neureclipsis</u> sp.                  | 13       | 13       | 12         | 118      | 112     |
| <u>Polycentropus</u> sp.                 |          | 2        |            |          | 17      |
| <u>Psychomyia</u> sp.                    |          |          |            |          |         |
| <u>Cyrnellus marginatus</u>              |          |          |            |          |         |
| <u>Psychomyiid</u> genus A(Ross)         |          | 1        | 16         | 9        | 2       |
| <u>Psychomyiid</u> genus B(Ross)         |          |          |            |          |         |
| Undet. <u>Psychomyiidae</u>              |          |          |            |          | 2       |
| <u>Potamyia flava</u>                    |          |          |            |          | 46      |
| <u>Hydropsyche frison</u>                | 202      |          | 6, 1P**300 | 242      | 202     |
| <u>Hydropsyche orris</u>                 |          |          |            |          |         |
| <u>Hydropsyche simulans</u>              |          |          |            |          |         |
| Undet. <u>Hydropsyche</u>                | 80, 3P** | 8        | 7, 5P      | 253, 1P  | 226, 2P |
| <u>Cheumatopsyche</u> sp.                | 1        | 8        |            | 26       | 1       |
| <u>Macronemus carolina</u>               | 3107, 2P | 1        | 53         | 107      | 3484    |
| Undet. <u>Hydropsychidae</u>             |          |          |            |          | 2006    |
| <u>Ochrotrichia</u> sp.                  |          |          |            |          | 113     |
| <u>Orthotrichia</u> sp.                  |          |          |            |          |         |
| <u>Hydroptila</u> sp.                    |          |          |            | 1        |         |
| Undet. <u>Limnophilidae</u>              |          |          |            |          |         |
| <u>Leptocella candida</u> , L.           |          |          |            |          |         |
| genus A(Ross)                            | 2        |          | 27         | 22       | 5       |

\*\* = Pupa

(continued)

TABLE I (Appendix "G")  
(Continued)

## SAVANNAH RIVER STUDIES

July 1951-July 1952

| COMPLETE LIST - Brush Box Sampling |          |                      |          | Numbers of Organisms |          |         |          |
|------------------------------------|----------|----------------------|----------|----------------------|----------|---------|----------|
| 8-160.75                           | 8-155.25 | 8-155.2 <sup>2</sup> | 8-155.17 | 8-146.4              | 8-144.7  | 8-136.6 | 8-51.0   |
| 11/2/51                            | 10/1/51  | 9/26/51              | 9/6/51   | 10/11/51             | 10/11/51 | 2/6/52  | 7/31/52  |
| 32                                 | 24       | 20                   | 22       | 21                   | 21       | 53      | 160      |
|                                    |          |                      |          |                      |          |         | 3        |
|                                    | 2        |                      | 3        |                      |          |         | 1        |
|                                    |          |                      | 1        |                      |          |         |          |
|                                    |          |                      |          | 1                    |          |         |          |
| 12                                 | 4        |                      | 11       |                      |          |         | 19       |
|                                    | 1        |                      | 5        |                      |          |         |          |
|                                    |          |                      | 3        |                      |          |         | 1        |
| 19                                 | 27       | 17                   | 15       | 37                   | 2        | 2       | 12       |
|                                    |          | 7                    |          | 1                    |          |         |          |
| 7                                  |          |                      | 7        |                      | 1        | 1       | 2        |
| 21                                 |          |                      | 1        |                      |          | 1       |          |
| 564                                | 3        |                      | 21       |                      | 8        | 4       | 76       |
|                                    |          |                      | 1        |                      | 1        |         |          |
|                                    |          |                      |          | 7                    |          |         |          |
|                                    |          | 17                   |          |                      |          |         | 3        |
|                                    |          |                      |          |                      |          | 1       |          |
| 1                                  | 6        |                      | 19       |                      | 1        |         |          |
| 6                                  | 17       |                      | 87       |                      | 15       |         | 62       |
|                                    |          |                      |          |                      |          | 1       |          |
|                                    |          |                      |          |                      |          | 1       |          |
| 69                                 | 41       |                      |          |                      | 30       | 46      |          |
|                                    |          |                      |          |                      |          | 1       | 183      |
| 1293                               | 453      |                      | 797      | 1                    | 2144     | 15      | 447,3P** |
| 1425                               | 1        |                      | 110      |                      | 28       |         | 19       |
|                                    |          |                      |          |                      |          |         | 2        |
| 53                                 | 1        |                      | 5        |                      | 22       | 10      | 47,2P    |
| ** = Pupae                         |          |                      |          | (continued)          |          |         |          |

TABLE I (Appendix "G")  
(continued)

## SAVANNAH RIVER STUDIES

July 1951-July 1952

## COMPLETE LIST - Brush Box Sampling

## Numbers of Organisms

| Location (River Mi.)                    | 8-223.15 |         |             |         | 8-219.0 |          |
|-----------------------------------------|----------|---------|-------------|---------|---------|----------|
| Date Removed                            | 8/24/51  | 7/30/51 | 8/6/51      | 8/13/51 | 8/20/51 | 10/16/51 |
| No. of days in place                    | 21       | 7       | 14          | 21      | 28      | 21       |
| <u>Athripsodes</u> nr. <u>mentiosus</u> |          |         |             |         |         |          |
| <u>Athripsodes transversus</u> ,        |          |         |             |         |         |          |
| <u>A. sp.</u>                           |          |         |             |         |         |          |
| <u>Deletis inconspicua</u>              | 3        |         | 3           |         | 1       |          |
| <u>Traienodes</u> sp.                   |          |         |             | 2       |         |          |
| Undet. <u>Leptoceridae</u>              |          |         |             |         |         |          |
| <u>Brachycentrus numerosus</u>          |          |         |             |         |         |          |
| Undetermined                            |          |         | 4           |         |         |          |
| Hemiptera: Undet. <u>Veliidae</u>       |          |         |             |         |         |          |
| Coleoptera: <u>Dineutes</u> sp.         |          |         |             |         |         |          |
| <u>Peltodytes</u> sp.                   |          |         |             |         |         |          |
| <u>Hydrovatus</u> sp.                   |          |         |             |         |         |          |
| Undet. <u>Dytiscidae</u>                |          |         |             |         |         |          |
| <u>Hydrochus</u> sp.                    |          |         |             | 1       |         |          |
| <u>Berosus</u> sp.                      |          |         |             |         |         |          |
| <u>Dryops</u> sp.                       |          |         |             |         |         |          |
| <u>Helichus</u> sp.                     |          |         |             |         |         |          |
| <u>Ancyronyx variegatus</u>             |          |         | 3A*         | 1A      | 3A      | 1A       |
| <u>Macronychus glabratus</u>            | 1A       | 6A      | 1,20A       | 8A      | 24A     | 28A      |
| <u>Stenelmis convexula, decorata</u>    | 6A       | 6A      | 3A          | 1A      | 13A     | 1A       |
| <u>Simsonia</u> sp.                     |          |         |             |         |         |          |
| Undet. <u>Elmidae</u>                   |          |         |             |         |         |          |
| Undetermined                            |          |         |             |         |         |          |
| Diptera: <u>Atherix variegata</u>       |          |         |             |         |         |          |
| <u>Simulium</u> sp.                     |          |         |             |         |         |          |
| <u>Chaoborus punctipennis</u>           |          |         |             |         |         |          |
| <u>Ablabesmyia carnea</u>               |          |         |             |         |         |          |
| <u>Ablabesmyia monilis</u>              | 28       | 24      | 109,3P**122 |         | 169,4P  | 13       |
| <u>Ablabesmyia flavifrons</u> ,         |          |         |             |         |         |          |
| <u>P. melanops</u>                      | 3        |         | 1           |         |         |          |
| <u>Ablabesmyia pilosella</u>            |          |         |             |         |         |          |
| Undet. <u>Ablabesmyia</u>               |          |         |             |         |         |          |
| <u>Anatopynia dyari</u>                 | 1        |         |             |         |         |          |
| <u>Procladius</u> sp.                   |          |         |             |         |         |          |
| <u>Clinotanypus</u> sp.                 |          |         |             |         |         |          |
| <u>Coelotanypus</u> nr. <u>concinna</u> |          |         |             | 5       | 14      |          |
| <u>Corynoneura scutellata</u> , C.spp.  |          |         |             |         |         |          |
| <u>Thienemanniella</u> sp.              |          |         |             |         |         |          |
| <u>Brillia par</u>                      |          |         |             |         |         |          |
| <u>Spaniotoma</u> sens. lat. spp.       |          |         | 2           | 1       | 2       | 18,3P    |
| <u>Psectrocladius</u> sp.               |          |         |             |         |         |          |
| <u>Tanytarsus signatus flavellus</u>    |          |         | 1P          | 5P      | 3P      | 4P       |
| <u>Tanytarsus dissimilis</u>            |          |         |             |         |         |          |
| <u>Tanytarsus</u> sens. lat. spp.       |          | 1       | 12          | 18      | 9       | 9        |

\* A = Adult    \*\* P = Pupa

(continued)

TABLE I (Appendix "G")  
(continued)

## SAVANNAH RIVER STUDIES

July 1951-July 1952

## COMPLETE LIST - Brush Box Sampling

## Numbers of Organisms

| 8-215.5 |         |          | 8-213.3 | 8-203.3 | 8-201.55 | 8-172.0 |        |
|---------|---------|----------|---------|---------|----------|---------|--------|
| 8/27/51 | 9/19/51 | 10/16/51 | 9/19/51 | 8/27/51 | 8/28/51  | 7/27/52 | 9/4/51 |
| 21      | 21      | 21       | 21      | 21      | 22       | 165     | 21     |

|      |         |      |        |                    |        |        |    |
|------|---------|------|--------|--------------------|--------|--------|----|
| 1P** | 3       | 1    |        | 2                  | 16,28P | 1P     | 1  |
|      |         | 1    |        |                    |        |        |    |
|      |         | 1    |        |                    |        |        |    |
|      |         | 1    |        |                    |        |        |    |
|      |         | 1    |        |                    |        |        |    |
|      |         | 1    |        |                    |        |        |    |
|      |         | 1    |        |                    |        |        |    |
|      |         | 1    |        |                    |        |        |    |
|      |         | 1    |        |                    |        |        |    |
|      |         | 1    |        |                    |        |        |    |
| 2    |         | 3    | 39,8A* | 6A<br>10,21A<br>3A |        |        |    |
| 1A   | 1P      | 2A   | 1A     |                    | 17     |        |    |
| 2A   | 1A      | 2A   | 1A     |                    | 2      |        |    |
|      |         |      |        |                    | 2      |        |    |
|      |         |      |        |                    | 1      |        |    |
|      |         |      |        |                    |        |        |    |
|      |         |      |        |                    |        |        |    |
|      |         |      |        |                    |        |        |    |
|      |         |      |        |                    |        |        |    |
|      |         |      |        |                    |        |        |    |
| 9    | 394,19P | 12   | 987,3P | 81                 | 269,1P | 104,4P | 31 |
| 2    |         | 1    | 1      | 2                  |        |        |    |
| 2    |         |      | 18,1P  | 4                  | 5      | 23     |    |
| 2    |         |      | 5      | 4                  | 13     |        |    |
| 1    | 6,1P    | 8,3P | 2      | 17,1P              |        | 25,25P |    |

\* A = Adult

\*\* P = Pupa

(continued)

TABLE I (Appendix "G")  
(continued)

## SAVANNAH RIVER STUDIES

July 1951-July 1952

## COMPLETE LIST - Brush Box Sampling

## Numbers of Organisms

| Location (River Mi.)                                  | 8-171.7 | 8-171.6  | 8-171.52 | 8-171.15 | 8-165.4   |
|-------------------------------------------------------|---------|----------|----------|----------|-----------|
| Date Removed                                          | 10/3/51 | 12/20/51 | 10/4/51  | 10/31/51 | 9/4/51    |
| No. of days in place                                  | 21      | 37       | 22       | 26       | 21        |
|                                                       |         |          |          |          | 21        |
|                                                       |         |          |          |          | 139       |
| <u>Athripsodes</u> nr. <u>mentios</u>                 |         |          |          | 1        | 2         |
| <u>Athripsodes</u> <u>transversus</u> ,               |         |          |          |          |           |
| <u>A. sp.</u>                                         |         |          | 2        | 2        | 3         |
| <u>Deletis</u> <u>inconspicua</u>                     | 3       |          | 13       | 12       | 7         |
| <u>Traienodes</u> <u>sp.</u>                          |         |          |          | 7        | 7         |
| <u>Undet. Leptoceridae</u>                            |         |          |          |          |           |
| <u>Brachycentrus</u> <u>numerosus</u>                 |         | 14       |          | 2        | 1         |
| <u>Undetermined</u>                                   |         |          |          |          |           |
| Hemiptera: <u>Undet. Veliidae</u>                     |         |          |          |          |           |
| Coleoptera: <u>Dineutes</u> <u>sp.</u>                |         |          |          |          |           |
| <u>Peltodytes</u> <u>sp.</u>                          |         |          |          |          |           |
| <u>Hydrovatus</u> <u>sp.</u>                          |         |          |          |          |           |
| <u>Undet. Dytiscidae</u>                              |         |          |          |          |           |
| <u>Hydrochus</u> <u>sp.</u>                           |         |          |          | 8A       | 2A        |
| <u>Berosus</u> <u>sp.</u>                             |         |          |          |          |           |
| <u>Dryops</u> <u>sp.</u>                              |         |          |          |          |           |
| <u>Helichus</u> <u>sp.</u>                            |         |          |          |          | 1A        |
| <u>Ancyronyx</u> <u>variegatus</u>                    | 1,7A    | 3,7A     | 2A       |          |           |
| <u>Macronychus</u> <u>glabratus</u>                   | 1,27A   | 3A       | 2,10A    | 1,3A     | 6,8A 4A   |
| <u>Stenelmis</u> <u>convexula</u> ,                   |         |          | 4,95A    | 5,29A    | 6,53A 17A |
| <u>decorata</u>                                       | 9,62A   | 2,6A     | 1,406A   | 126A     | 8,14A 28A |
| <u>Simsonia</u> <u>sp.</u>                            |         |          |          |          | 1         |
| <u>Undet. Elmidae</u>                                 |         |          |          |          | 1         |
| <u>Undetermined</u>                                   |         |          |          |          |           |
| Diptera: <u>Atherix</u> <u>variegata</u>              |         |          |          | 1        |           |
| <u>Simulium</u> <u>sp.</u>                            |         | 11       | 1        | 3        |           |
| <u>Chaoborus</u> <u>punctipennis</u>                  |         | 1        |          | 6        | 3         |
| <u>Ablabesmyia</u> <u>carnea</u>                      |         |          |          |          | 1         |
| <u>Ablabesmyia</u> <u>monilis</u>                     | 6       | 1        | 1        | 3        | 88        |
| <u>Ablabesmyia</u> <u>flavifrons</u> ,                |         |          |          |          | 41        |
| <u>P. melanops</u>                                    | 45      |          | 4        | 8        | 14        |
| <u>Ablabesmyia</u> <u>pilosella</u>                   | 4       |          | 1        |          | 7         |
| <u>Undet. Ablabesmyia</u>                             |         |          |          |          | 7         |
| <u>Anatopynia</u> <u>dyari</u>                        |         |          |          |          | 8         |
| <u>Procladius</u> <u>sp.</u>                          |         |          |          |          | 7         |
| <u>Clinotanypus</u> <u>sp.</u>                        |         |          |          | 1        | 23        |
| <u>Coelotanypus</u> nr. <u>concinna</u>               | 1       |          |          |          | 1         |
| <u>Corynoneura</u> <u>scutellata</u> , <u>C. spp.</u> |         | 1        |          | 1        |           |
| <u>Thienmanniella</u> <u>sp.</u>                      |         |          |          |          |           |
| <u>Brillia</u> <u>par</u>                             |         | 19       |          | 1        | 1         |
| <u>Spaniotoma</u> sens. lat. spp.                     | 6       | 23       |          | 24, 4P** | 9         |
| <u>Psectrocladius</u> <u>sp.</u>                      |         |          |          |          | 26        |
| <u>Tanytarsus</u> <u>signatus</u> <u>flavellus</u>    |         |          |          | 5P       | 1P        |
| <u>Tanytarsus</u> <u>dissimilis</u>                   |         |          |          |          |           |
| <u>Tanytarsus</u> sens. lat. spp.                     | 250     |          | 101      | 154      | 121       |
|                                                       |         |          |          |          | 10        |
|                                                       |         |          |          |          | 6         |

\* A = Adult

\*\* P = Pupa

(continued)

TABLE I (Appendix "G")  
(continued)

## SAVANNAH RIVER STUDIES

July 1951-July 1952

## COMPLETE LIST - Brush Box Sampling

## Numbers of Organisms

| S-160.75 | S-155.25 | S-155.2 | S-155.17 | S-146.4  | S-144.7  | S-136.6 | S-51.0  |
|----------|----------|---------|----------|----------|----------|---------|---------|
| 11/2/51  | 10/1/51  | 9/26/51 | 9/6/51   | 10/11/51 | 10/11/51 | 2/6/52  | 7/31/52 |
| 32       | 24       | 20      | 22       | 21       | 21       | 53      | 160     |

2

1  
4

4

3

6

1  
11  
1

## 2A\*

|        |        |         |       |       |      |      |         |
|--------|--------|---------|-------|-------|------|------|---------|
| 1A     |        | 8A      | 15A   |       | 20A  |      | 31      |
| 7A     | 1A     | 15A     | 1,66A | 5A    | 23A  | 4,7A | 173,13A |
| 90A    | 3A     |         | 27A   |       | 106A | 58A  | 14,11A  |
| 6      | 6      | 1A<br>6 |       | 13,1A | 2    |      |         |
| 3<br>2 | 2      |         | 18    |       |      |      | 6       |
|        |        |         |       | 11    |      |      | 1<br>23 |
| 34     | 26     |         | 7     |       | 3    | 4    | 3<br>1  |
| 11     | 8<br>1 | 11      | 27    | 16    | 2    |      |         |
| 4      | 1      | 3       | 1     |       | 5    |      |         |
| 13     |        | 1       |       |       | 1    |      |         |
| 8      | 1      | 5       |       | 2     | 1    | 15   | 2       |
| 854    | 268    | 89      | 162   | 6     | 2050 | 2    | 21      |

\* A = Adult

(continued)



TABLE I (Appendix "G")  
(continued)

## SAVANNAH RIVER STUDIES

July 1951-July 1952

## COMPLETE LIST - Brush Box Sampling

## Numbers of Organisms

| Location (River Mi.) | S-223.15 |         |        |         | S-219.0 |          |
|----------------------|----------|---------|--------|---------|---------|----------|
| Date Removed         | 8/24/51  | 7/30/51 | 8/6/51 | 8/13/51 | 8/20/51 | 10/16/51 |
| No. of days in place | 21       | 7       | 14     | 21      | 28      | 21       |

|                                     |    |   |        |    |        |        |
|-------------------------------------|----|---|--------|----|--------|--------|
| <u>Rheotanytarsus exiguus</u>       |    |   |        |    |        |        |
| <u>Microspectra</u> sp.             |    |   |        |    |        | 1P**   |
| <u>Microtendipes pedellus</u>       |    |   |        |    |        |        |
| <u>Stenochironomus taeniopennis</u> | 36 |   |        |    |        |        |
| <u>Stictochironomus</u> nr.         |    |   |        |    |        |        |
| <u>flavicingula</u> johannsen       |    |   |        |    |        |        |
| <u>Polypedilum fallax</u>           | 2P |   | 2      | 5  | 2      | 1      |
| <u>Polypedilum illinoiense</u>      |    |   |        |    |        |        |
| Undet. <u>Polypedilum</u> spp.      | 13 | 4 | 42, 2P | 35 | 23     | 2      |
| <u>Endochironomus nigricans</u>     | 8  | 1 | 24, 1P | 17 | 49, 2P | 44, 1P |
| <u>Endochironomus tendens</u>       |    |   |        |    |        |        |
| <u>Endochironomus quadri-</u>       |    |   |        |    |        |        |
| <u>punctatus</u>                    |    |   |        |    |        |        |
| <u>Glyptotendipes lobiferus</u>     |    |   |        |    |        | 52     |
| <u>Cryptochironomus stylifer</u>    | 2  |   |        | 1  |        |        |
| <u>Cryptochironomus</u> sp.B.       |    |   |        |    |        |        |
| johannsen                           | 5  |   | 8      | 1  | 6      | 1      |
| Undet. <u>Cryptochironomus</u>      |    |   |        |    |        |        |
| <u>Limnochironomus modestus</u>     | 8  |   | 1      | 2  | 5      | 7, 1P  |
| <u>Parachironomus tenuicaudatus</u> |    |   |        |    |        |        |
| <u>Synchironomus</u> sp.            | 1  |   |        |    |        |        |
| <u>Chironomus plumosus</u> ,        |    |   |        |    |        |        |
| <u>Camptochironomus tentans</u>     |    |   |        |    |        |        |
| <u>Chironomus riparius</u>          |    |   |        |    |        |        |
| Undet. Chironomidae                 |    |   |        |    |        |        |
| (17 spp. larvae,                    |    |   |        |    |        |        |
| 9 spp. pupae)                       | 2  |   | 1, 5P  | 1P | 2, 1P  | 1, 2P  |
| <u>Bezzia setulosa</u>              |    |   |        |    |        |        |
| Undet. <u>Bezzia</u> sp.            |    |   |        |    |        |        |
| <u>Palpomyia</u> sp.                |    |   |        |    |        |        |
| <u>Stilobezzia</u> sp.              |    |   |        |    |        |        |
| Undet. Ceratopogonidae              |    |   |        | 3  | 1      |        |
| <u>Limnophora</u> sp.               |    |   |        |    |        |        |
| MOLLUSCA: GASTROPODA                |    |   |        |    |        |        |
| <u>Physa</u> sp.                    | 2  |   |        |    | 2      |        |
| <u>Gyraulus</u> spp.                |    |   |        |    | 4      | 1      |
| Undet. <u>Planorbidae</u>           | 2  |   |        |    |        |        |
| <u>Ferrissia</u> , sp.              | 1  |   | 1      |    | 7      |        |
| <u>Campeloma</u> sp.                |    |   |        |    |        |        |

\*\* P = Pupa

(continued)

TABLE I (Appendix "G")  
(continued)

## SAVANNAH RIVER STUDIES

July 1951-July 1952

## COMPLETE LIST - Brush Box Sampling

## Numbers of Organisms

| S-215.5 |         |          | S-213.3 | S-203.3 | S-201.55 | S-172.0 |        |
|---------|---------|----------|---------|---------|----------|---------|--------|
| 8/27/51 | 9/19/51 | 10/16/51 | 9/19/51 | 8/27/51 | 8/28/51  | 7/27/52 | 9/4/51 |
| 21      | 21      | 21       | 21      | 21      | 22       | 165     | 21     |

|      |        |    |       |      |         |         |    |
|------|--------|----|-------|------|---------|---------|----|
| 3    | 3,1P** |    | 3     | 6    | 5       | 23      | 12 |
| 2    | 39     | 1  | 36    | 181  | 33      | 11      |    |
|      | 3      |    | 5     | 7    |         | 2       |    |
|      |        |    | 2     |      | 7       |         |    |
| 32   | 4      |    | 2     | 29   | 1       | 9       | 3  |
| 8    | 53,7P  |    | 26,2P | 3    | 169,26P | 1       |    |
|      |        | 4P |       | 8    | 116     | 291,60P |    |
|      |        |    | 3     |      |         |         |    |
|      |        |    |       |      |         | 45      |    |
| 1,5P | 18,3P  | 4P | 22,7P | 1,1P | 15,12P  | 2,8P    | 4  |
|      |        |    |       |      |         |         |    |
|      |        |    |       |      |         | 7       |    |
| 9    | 31     | 29 | 94    | 1    | 9       | 28      |    |
| 17   | 54     | 69 | 20    |      |         | 4       |    |
|      |        |    | 7     |      |         | 17      |    |
| 1    |        | 3  |       |      | 1       | 2       |    |

\*\* P = Pupa

(continued)

TABLE I (Appendix "G")  
(continued)

## SAVANNAH RIVER STUDIES

July 1951-July 1952

## COMPLETE LIST - Brush Box Sampling

## Numbers of Organisms

| Location (River Mi.)                                            | S-171.7  | S-171.6  | S-171.52 | S-171.15 | S-165.4 |
|-----------------------------------------------------------------|----------|----------|----------|----------|---------|
| Date Removed                                                    | 10/3/51  | 12/20/51 | 10/4/51  | 10/31/51 | 9/4/51  |
| No. of days in place                                            | 21       | 37       | 22       | 26       | 21      |
|                                                                 |          |          |          |          | 21      |
|                                                                 |          |          |          |          | 139     |
| <u>Rheotanytarsus exiguus</u>                                   | 85P**    | 28       | 24P      | 76P      | 4P      |
| <u>Microspectra</u> sp.                                         |          | 1        |          |          |         |
| <u>Microtendipes pedellus</u>                                   |          |          |          |          |         |
| <u>Stenochironomus taeniopennis</u>                             |          |          |          |          |         |
| <u>Stictochironomus</u> nr.<br><u>flavicingula</u> Johannsen    |          |          |          |          |         |
| <u>Polypedilum fallax</u>                                       |          | 4        |          |          |         |
| <u>Polypedilum illinoiense</u>                                  |          |          |          | 24       | 64      |
| Undet. <u>Polypedilum</u> spp.                                  | 303, 14P | 7        | 15       | 63       | 141, 4P |
| <u>Endochironomus nigricans</u>                                 |          | 5        |          |          | 17      |
| <u>Endochironomus tendens</u>                                   |          |          |          |          | 1       |
| <u>Endochironomus quadri-</u><br><u>punctatus</u>               |          |          |          |          | 1       |
| <u>Glyptotendipes lobiferus</u>                                 |          |          |          |          |         |
| <u>Cryptochironomus stylifer</u>                                |          |          | 1        | 6        | 1       |
| <u>Cryptochironomus</u> sp. B.<br>Johannsen                     | 1        |          | 2        | 1        | 31      |
| Undet. <u>Cryptochironomus</u>                                  |          |          |          |          | 1       |
| <u>Limnochironomus modestus</u>                                 | 1        |          | 1, 1P    | 2, 2P    |         |
| <u>Parachironomus tenuicaudatus</u>                             |          |          |          |          |         |
| <u>Synchironomus</u> sp.                                        |          |          |          |          |         |
| <u>Chironomus plumosus</u> ,<br><u>Camptochironomus tentans</u> |          |          |          |          |         |
| <u>Chironomus riparius</u>                                      |          |          |          |          |         |
| Undet. Chironomidae<br>(17 spp. larvae,<br>9 spp. pupae)        | 2, 2P    | 2, 1P    | 1, 1P    | 13, 14P  | 17, 10P |
| <u>Bezzia setulosa</u>                                          |          |          |          | 1        | 3,      |
| Undet. <u>Bezzia</u> sp.                                        |          |          |          |          |         |
| <u>Palpomyia</u> sp.                                            |          |          |          | 1        |         |
| <u>Stilobezzia</u> sp.                                          |          |          |          |          |         |
| Undet. Ceratopogonidae                                          |          | 1        |          |          |         |
| <u>Limnophora</u> sp.                                           |          |          |          |          | 1       |
|                                                                 |          |          |          |          | 1       |
| MOLLUSCA: GASTROPODA                                            |          |          |          |          |         |
| <u>Physa</u> sp.                                                |          |          | 1        |          |         |
| <u>Gyraulus</u> spp.                                            |          |          |          |          | 3       |
| Undet. Planorbidae                                              |          |          |          |          |         |
| <u>Ferrissia</u> , sp.                                          |          |          |          |          |         |
| <u>Cameloma</u> sp.                                             |          |          |          |          |         |

\*\* P = Pupa

(continued)

TABLE I (Appendix "G")  
(continued)

## SAVANNAH RIVER STUDIES

July 1951-July 1952

## COMPLETE LIST - Brush Box Sampling

## Numbers of Organisms

| S-160.75 | S-155.25 | S-155.2 <sup>2</sup> | S-155.17 | S-146.4  | S-144.7  | S-136.6 | S-51.0  |
|----------|----------|----------------------|----------|----------|----------|---------|---------|
| 11/2/51  | 10/1/51  | 9/26/51              | 9/6/51   | 10/11/51 | 10/11/51 | 2/6/52  | 7/31/52 |
| 32       | 24       | 20                   | 22       | 21       | 21       | 53      | 160     |

1P\*\*

|    |     |   |     |    |     |   |      |
|----|-----|---|-----|----|-----|---|------|
| 70 | 106 |   | 105 |    | 269 | 1 | 8    |
| 3  |     |   |     | 4  |     | 4 |      |
| 2  |     | 3 |     | 32 | 1   | 2 | 6    |
| 1  |     |   |     |    |     |   |      |
|    |     |   |     |    |     |   | 1,1P |
| 11 |     | 1 | 2   |    |     |   |      |
| 1  |     |   |     |    |     |   |      |
|    |     |   |     |    |     |   |      |
| 4P | 24  | 7 | 2   | 1  |     | 6 | 5,1P |
|    | 3   |   |     | 1  |     |   |      |
|    |     |   |     |    |     |   | 1    |
|    |     | 1 |     | 2  | 9   |   |      |
| 3  |     |   |     |    |     |   |      |
| 1  |     |   | 1   |    |     |   |      |

\*\* P = Pupa

(continued)

TABLE I. (Appendix "G") SAVANNAH RIVER STUDIES July 1951-July 1952  
(Continued)  
COMPLETE LIST - Brush Box Sampling Numbers of Organisms

| Location (River Mi.) | S-223.15 |         |        |         | S-219.0 |          |
|----------------------|----------|---------|--------|---------|---------|----------|
| Date Removed         | 8/24/51  | 7/30/51 | 8/6/51 | 8/13/51 | 8/21/51 | 10/16/51 |
| No. of days in place | 21       | 7       | 14     | 21      | 28      | 21       |

|                        |   |   |  |   |   |    |
|------------------------|---|---|--|---|---|----|
| <u>Pleurocera</u> sp.  |   |   |  |   |   |    |
| <u>Somatogyrus</u> sp. |   |   |  |   |   | 1  |
| <u>Amnicola</u> spp.   |   | 3 |  |   | 7 | 12 |
| Undet. Amnicolidae     | 3 |   |  | 2 |   |    |
| Undetermined           |   |   |  |   |   |    |

## MOLLUSCA:PELECYPODA

|                      |  |  |  |  |   |  |
|----------------------|--|--|--|--|---|--|
| <u>Pisidium</u> sp.  |  |  |  |  |   |  |
| <u>Sphaerium</u> sp. |  |  |  |  |   |  |
| <u>Musculium</u> sp. |  |  |  |  |   |  |
| <u>Eupera</u> sp.    |  |  |  |  | 1 |  |
| Undet. Sphaeridae    |  |  |  |  |   |  |
| Undet. Unionidae     |  |  |  |  |   |  |

TOTAL ORGANISMS 365 133 337 332 536 386

(concluded below)

| Location (River Mi.) | S-171.7 | S-171.6  | S-171.52 | S-171.15 | S-165.4 |        |
|----------------------|---------|----------|----------|----------|---------|--------|
| Date Removed         | 10/3/51 | 12/20/51 | 10/4/51  | 10/31/51 | 9/4/51  | 9/4/51 |
| No. of days in place | 21      | 37       | 22       | 26       | 21      | 21     |

|                        |  |  |  |  |   |  |
|------------------------|--|--|--|--|---|--|
| <u>Pleurocera</u> sp.  |  |  |  |  |   |  |
| <u>Somatogyrus</u> sp. |  |  |  |  |   |  |
| <u>Amnicola</u> spp.   |  |  |  |  | 1 |  |
| Undet. Amnicolidae     |  |  |  |  |   |  |
| Undetermined           |  |  |  |  |   |  |

## MOLLUSCA:PELECYPODA

|                      |  |  |  |  |  |   |
|----------------------|--|--|--|--|--|---|
| <u>Pisidium</u> sp.  |  |  |  |  |  |   |
| <u>Sphaerium</u> sp. |  |  |  |  |  |   |
| <u>Musculium</u> sp. |  |  |  |  |  |   |
| <u>Eupera</u> sp.    |  |  |  |  |  |   |
| Undet. Sphaeridae    |  |  |  |  |  |   |
| Undet. Unionidae     |  |  |  |  |  | 1 |

TOTAL ORGANISMS 4851 270 1330 1491 5465 2943 237

TABLE I (Appendix "G") SAVANNAH RIVER STUDIES July 1951-July 1952  
(continued)

COMPLETE LIST - Brush Box Sampling

Numbers of Organisms

| S-215.5 |         |          | S-213.3 | S-203.3 | S-201.55 | S-172.0 |        |
|---------|---------|----------|---------|---------|----------|---------|--------|
| 8/27/51 | 9/19/51 | 10/16/51 | 9/19/51 | 8/27/51 | 8/28/51  | 7/27/52 | 9/4/51 |
| 21      | 21      | 21       | 21      | 21      | 22       | 165     | 21     |

7 10 3  
9 64 8 169 1 1 22

1 5 9 2  
6 67  
3  
1 2

61 870 353 1835 371 1074 1510 204

(concluded below)

| S-160.75 | S-155.25 | S-155.2<br>St-0.0 | S-155.17 | S-146.4  | S-144.7  | S-136.6 | S-51.0  |
|----------|----------|-------------------|----------|----------|----------|---------|---------|
| 11/2/51  | 10/1/51  | 9/26/51           | 9/6/51   | 10/11/51 | 10/11/51 | 2/6/52  | 7/31/52 |
| 32       | 24       | 20                | 22       | 21       | 21       | 53      | 160     |

13

6

3

1

5171 1186 227 1625 193 5366 386 1386